



**HAL**  
open science

## Germline mutations in the new E1' cryptic exon of the VHL gene in patients with tumours of von Hippel-Lindau disease spectrum or with paraganglioma

Alexandre Buffet, Bruna Calsina, Shahida Flores, Sophie Giraud, Marion Lenglet, Pauline Romanet, Elisa Deflorenne, Javier Aller, Isabelle Bourdeau, Brigitte Bressac-De Paillerets, et al.

### ► To cite this version:

Alexandre Buffet, Bruna Calsina, Shahida Flores, Sophie Giraud, Marion Lenglet, et al.. Germline mutations in the new E1' cryptic exon of the VHL gene in patients with tumours of von Hippel-Lindau disease spectrum or with paraganglioma. *Journal of Medical Genetics*, 2020, pp.1 - 8. 10.1136/jmedgenet-2019-106519 . hal-02484923

**HAL Id: hal-02484923**

**<https://ut3-toulouseinp.hal.science/hal-02484923>**

Submitted on 29 May 2021

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



OPEN ACCESS

ORIGINAL RESEARCH

# Germline mutations in the new E1' cryptic exon of the *VHL* gene in patients with tumours of von Hippel-Lindau disease spectrum or with paraganglioma

Alexandre Buffet <sup>1</sup>, Bruna Calsina,<sup>2</sup> Shahida Flores,<sup>3</sup> Sophie Giraud,<sup>4,5</sup> Marion Lenglet,<sup>6,7</sup> Pauline Romanet,<sup>8</sup> Elisa Deflorenne,<sup>1</sup> Javier Aller,<sup>9</sup> Isabelle Bourdeau,<sup>10</sup> Brigitte Bressac-de Paillerets,<sup>5,11</sup> María Calatayud,<sup>12</sup> Caroline Dehais,<sup>13</sup> Erwan De Mones Del Pujol,<sup>14</sup> Atanaska Elenkova,<sup>15</sup> Philippe Herman,<sup>16</sup> Peter Kamenický,<sup>17</sup> Sophie Lejeune,<sup>5,18</sup> Jean Louis Sadoul,<sup>19</sup> Anne Barlier,<sup>5,8</sup> Stephane Richard,<sup>5,6,20</sup> Judith Favier,<sup>1</sup> Nelly Burnichon,<sup>1,21</sup> Betty Gardie,<sup>6,7</sup> Patricia L Dahia,<sup>3</sup> Mercedes Robledo,<sup>2</sup> Anne-Paule Gimenez-Roqueplo<sup>1,5,21</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jmedgenet-2019-106519>).

For numbered affiliations see end of article.

## Correspondence to

Dr Alexandre Buffet, Université de Paris, PARCC, INSERM, Equipe Labellisée par la Ligue contre le Cancer, F-75015, Paris, France; alexandre.buffet@inserm.fr

BC, SF, SG and ML contributed equally.

Received 22 August 2019  
Revised 3 January 2020  
Accepted 6 January 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Buffet A, Calsina B, Flores S, et al. *J Med Genet* Epub ahead of print: [please include Day Month Year]. doi:10.1136/jmedgenet-2019-106519

## ABSTRACT

**Backgrounds** The incidence of germline mutations in the newly discovered cryptic exon (E1') of *VHL* gene in patients with von Hippel-Lindau (VHL) disease and in patients with paraganglioma or pheochromocytoma (PPGL) is not currently known.

**Methods** We studied a large international multicentre cohort of 1167 patients with a previous negative genetic testing. Germline DNA from 75 patients with a single tumour of the VHL spectrum ('Single VHL tumour' cohort), 70 patients with multiple tumours of the VHL spectrum ('Multiple VHL tumours' cohort), 76 patients with a VHL disease as described in the literature ('VHL-like' cohort) and 946 patients with a PPGL were screened for E1' genetic variants.

**Results** Six different genetic variants in E1' were detected in 12 patients. Two were classified as pathogenic, 3 as variants of unknown significance and 1 as benign. The rs139622356 was found in seven unrelated patients but described in only 16 patients out of the 31 390 of the Genome Aggregation Database ( $p < 0.0001$ ) suggesting that this variant might be either a recurrent mutation or a modifier mutation conferring a risk for the development of tumours and cancers of the VHL spectrum.

**Conclusions** *VHL* E1' cryptic exon mutations contribute to 1.32% (1/76) of 'VHL-like' cohort and to 0.11% (1/946) of PPGL cohort and should be screened in patients with clinical suspicion of VHL, and added to panels for Next Generation Sequencing (NGS) diagnostic testing of hereditary PPGL. Our data highlight the importance of studying variants identified in deep intronic sequences, which would have been missed by examining only coding sequences of genes/exons. These variants will likely be more frequently detected and studied with the upcoming implementation of whole-genome sequencing into clinical practice.

## INTRODUCTION

Von Hippel-Lindau (VHL) disease is an autosomal-dominant renal cancer predisposition syndrome<sup>1</sup>

responsible for the development in affected patients of renal cysts or clear cell carcinomas, and other features as retinal or central nervous system haemangioblastomas, pancreatic cysts or neuroendocrine tumours, endolymphatic sac tumours and pheochromocytomas and/or paragangliomas (PPGLs). A germline mutation (including gross deletion) is identified in one of the three exons of *VHL* in almost all affected patients.<sup>2</sup> Nevertheless, some patients with clinically diagnosed VHL disease, but without identified *VHL* germline mutation, have been reported.<sup>3</sup> One of the tumour types of the VHL tumour spectrum, PPGL, are rare neuroendocrine tumours with a great genetic heterogeneity and the highest heritability rate with about 40% of genetically determined forms.<sup>4,5</sup> Indeed, to date, approximately 17 susceptibility genes have been reported but two thirds of identified mutations are found in *SDHB*, *SDHD* and *VHL* genes.<sup>6,7</sup>

Recently, a cryptic exon of *VHL* gene, named E1', has been discovered. A germline mutation in the first intronic region which results in creation of a cryptic exon designated E1' was found in one large family with a typical VHL disease and without any alteration in the other *VHL* exons.<sup>8</sup> *VHL* gene is one of the major PPGL susceptibility genes but, to date, E1' exon has not been included in PPGL target gene panels.

Hence, our objective was to assess the prevalence of E1' germline mutations in two international cohorts of patients: first, in 221 patients with a single or multiple tumours suggesting a VHL disease and then in 946 with a PPGL but without an identified mutation in the three *VHL* exons or in the main PPGL susceptibility genes, respectively.

## METHODS

### Patient's selection

A total of 1167 patients were analysed, divided into four different groups:

**Table 1** Main clinical and tumour characteristics of the different cohorts

Total patients n=1167	
<i>VHL-like</i> , n=76 patients	
Age at first diagnosis mean (min-max)	45.4 (20–76)
Multiple haemangioblastomas	16 (21%)
Haemangioblastoma with another VHL tumour	28 (37%)
One VHL tumour and family history of VHL tumour	32 (42%)
<i>Multiple VHL tumours</i> , n=70 patients	
Age at first diagnosis mean, (min-max)	55 (11–81)
Three or more VHL tumours	3 (4%)
Two VHL tumours	67 (96%)
<i>Single VHL tumour</i> , n=75 patients	
Age at first diagnosis mean (min-max)	34.6 (11–78)
Clear cell renal cell carcinoma	3 (4%)
Cerebral haemangioblastoma	27 (36%)
Retinal haemangioblastoma	10 (13.3%)
Other tumours	35 (46.6%)
<i>PPGL</i> , n=946 patients	
Age at first diagnosis mean (min-max)	43 (8–94)
Benign PPGL	869 (92%)
Single benign PPGL	771 (82%)
Multiple benign PPGL	98 (10%)
Metastatic PPGL	77 (8%)
Single metastatic PPGL	67 (7%)
Multiple metastatic PPGL	10 (1%)
Familial PPGL	17 (2%)

PPGL, paraganglioma; VHL, von Hippel-Lindau.

- ▶ 946 patients with PPGL but without germline mutation in major PPGL susceptibility genes ('PPGL' cohort) (table 1 and online supplementary table S1).
- ▶ 76 patients with a VHL disease as defined in the literature,<sup>2,9</sup> that is, patients with multiple haemangioblastomas, or a single haemangioblastoma with another tumour of the VHL spectrum, or one tumour of the VHL spectrum (excepted epididymal and renal cysts) and family history of VHL tumour but no germline *VHL* gene mutation ('VHL-like' cohort).
- ▶ 70 patients with multiple tumours of the clinical spectrum of VHL disease but who did not fill the definition of a VHL disease and who had no germline *VHL* mutation ('Multiple VHL tumours' cohort).
- ▶ 75 patients with a single tumour of the VHL spectrum without *VHL* mutation occurring at a young age ('Single VHL tumour' cohort) (table 1 and online supplementary table S2).

Germline DNA from 'VHL-like', 'Multiple VHL tumours' and 'Single VHL tumour' cohorts had been previously tested for *VHL* gene by Sanger sequencing or Next Generation Sequencing and large rearrangements by MLPA or QMPSE. The procedures used for PPGL diagnosis were in accordance with international guidelines.<sup>10,11</sup>

Moreover, a control cohort of 198 European subjects without VHL manifestation was analysed in order to determine the frequency of variant in the general population.

Each patient signed a written informed consent for genetic analyses.

## Direct sequencing of the E1' cryptic exon of *VHL* on germline DNA

Sanger sequencing on germline DNA of E1' was performed as previously described.<sup>8</sup> Variants interpretation was performed by using different criterions: ACMG criteria,<sup>12</sup> allele frequency in databases, phenotype of patients and tumour analysis as described below.

## *VHL* gene analysis in tumour

Tumour DNA was extracted from frozen or paraffin embedded tumour by the QIAamp DNA minikit (Qiagen). Loss of heterozygosity (LOH) was evaluated by (1) Sanger sequencing of the E1' cryptic exon of *VHL* by mutation-specific primers and (2) microsatellite analysis on D3S1537, D3S1038, D3S1317 D3S3547, D3S3727 as previously described.<sup>13,14</sup> *VHL* gene deletion on tumour DNA was assessed with the *SALSA MLPA P016 VHL* probemix (MRC-Holland).

## CA9 immunochemistry

Immunohistochemistry was performed as previously described on 6 µm slides cut from paraffin-embedded tumours with anti CA9 antibody (1/1500, ab15086, Abcam).<sup>15</sup> Antigen retrieval was performed by boiling slides in Tris-EDTA buffer (pH9) for 45 min. Revelation was performed using Histogreen as a chromogen. Images were acquired with a Leica DM400B microscope with Leica Application Suite software V2.8.1 and a Leica DFC420C camera.

## *PNMT* and *VHL* RT-qPCR

RNA was extracted from paraffin embedded tumours of six control PPGL (3 *NF1*-related, 2 *RET*-related and 1 *TMEM127*-related PPGL), 5 *VHL*- related PPGL (all carrying a missense mutation in *VHL* gene) and patients #3 and #10 PPGL by using the Maxwell 16 LEV RNA FFPE Purification Kit (Promega). RNA was quantified and its purity assessed with a NanoDrop ND-1000 spectrophotometer (Labtech). RT PCR was performed on 1000 ng of RNA with iScript cDNA Synthesis Kit iScript (BioRad). Then, as described in,<sup>16</sup> pre amplification of *PNMT*, *VHL*, *GAPDH* and *18S* on complementary DNA was performed with SsoAdvanced PreAmp Supermix (BioRad). Because of RNA fragmentation, all primers were designed to amplify amplicons smaller than 100 bp. We used two *VHL* primer sets. The first set amplified the *VHL* transcript including exons 1 and 2 (E1-E2) (F: 5'-CATCCACAGCTACCGAGGTC-3' overlapping exons 1 and 2 and R: 5'-GTGTGTCCCTGCATCTCTGA-3' located on exon 2). The second set amplified the *VHL* transcript with exon 1 and the cryptic exon (E1-E1') (F: 5'-GCATCCACAGC-TACCGAGTC-3' overlapping exon 1 and the cryptic exon and R: 5'-AGTCTCCCCAGGAGGAATGT-3' located on the cryptic exon). Quantitative PCR was performed on *VHL* (E1-E2), *PNMT*, *GAPDH* and *18S* by SYBR Green Master MixSybrGreen (BioRad) on the C1000 Touch (BioRad) and *VHL* (E1-E1') was amplified by PCR in parallel with *GAPDH*. All experiments were performed in duplicate three times.

## Statistical analysis

Statistical analysis was carried out with GraphPad software. Differences between allele's frequency in gnomAD and our cohort of patients and relative risk were assessed by  $\chi^2$  tests. A  $p < 0.05$  was considered significant.

## RESULTS

We analysed the germline DNA of 1167 patients from France, Spain, Canada and the USA. We identified a rare germline

Table 2 Patients with a genetic variant in the E1' cryptic exon of VHL gene


Patient number	Phenotype (age at first diagnosis)	Follow-up duration (years)	Germline genetic variant	dbSNP ID	Allele frequency* (%)	LOH	Somatic VHL mutation	CAIX IHC	PMM1 and VHL (E1-E2) expression		Classification of VUS
									VHL (E1-E1) expression	VHL (E1-E2) expression	
#1	Carotid body PGL (47)	5	c.340+563C>T		NA	Not	ND	Neg	ND	ND	Benign
#2	Carotid body PGL (34)	1	c.340+578C>T	rs139622356	0.05	NA	NA	NA	NA	NA	VUS
#3	PCC (38)	7	c.340+578C>T	rs139622356	0.05	No	No	Neg	Normal	Increase	VUS
#4	Carotid body PGL (74)	1	c.340+578C>T	rs139622356	0.05	NA	NA	NA	NA	NA	VUS
#5	Carotid body PGL (56)	1	c.340+578C>T	rs139622356	0.05	NA	NA	NA	NA	NA	VUS
#6	PGL+ccRCC (81–82)	2	c.340+578C>T	rs139622356	0.05	NA	NA	NA	NA	NA	VUS
#7	bilccRCC (49–51)	NA	c.340+578C>T	rs139622356	0.05	NA	NA	NA	NA	NA	VUS
#8	ccRCC (39)	NA	c.340+578C>T	rs139622356	0.05	NA	NA	NA	NA	NA	VUS
#9	Multiple retinal HMB (36)	32	c.340+617C>G†		NA	NA	NA	NA	NA	NA	Pathogenic
#10	PCC (11)	15	c.340+682T>C		NA	No	c.482G>A p.R161Q	Pos	Decrease	Increase	Pathogenic
#11	Multiple HN PGL (12)	1	340+725A>T		NA	NA	NA	NA	NA	NA	VUS
#12	PCC (34)	10	c.340+866C>A	rs36631685	0.02	NA	NA	NA	NA	NA	VUS

\*Frequency in gnomAD or 1000 Genomes.

†Loss of the mutated allele.

‡Mutation described in Lenglet *et al.*<sup>8</sup>

bilccRCC, bilateral clear cell renal cell carcinoma; ccRCC, clear cell renal cell carcinoma; HMB, haemangioblastoma; HN, head and neck; LOH, loss of heterozygosity; NA, none available; ND, not done; Neg, negative immunohistochemistry; PCC, pheochromocytoma; PGL, paraganglioma; Pos, positive immunohistochemistry; VHL, von Hippel-Lindau; VUS, variant of uncertain significance.



**Figure 1** *VHL* E1' variants identified: mutations identified in patient #10 on the germline DNA and somatic DNA (A); location of germline variants found in *VHL* E1' (B).


genetic variant (minor allele frequency <1%) in the E1' *VHL* cryptic exon in 12 patients (1%). One of these patients was classified as 'VHL-like' (1 of 76 patients, 1.3%), 2 as 'Multiple VHL tumours' (2/70 patients, 2.9%), 1 as 'Single VHL tumour' (1/75 patients, 1.3%) and 8 belonged to the 'PPGL' cohort (8/946, 0.8%) (table 2). Among these 12 patients, we identified 6 different variants, 4 in the E1' and 2 at the intron-exon junction and we considered only two variants as pathogenic mutations (figure 1). None of these variants was found in a control cohort of 198 European subjects without VHL manifestations.

Seven patients (patients #2 to #8) (0.6%) carried the same rare variant of uncertain significance (VUS), c.340+578C>T which is referenced in dbSNP as rs139622356 and has been previously reported in the Genome Aggregation Database (gnomAD). The five remaining patients carried different E1' variants. One of them (patient #9) harbours the c.340+617C>G mutation previously described in the original paper.<sup>8</sup> Patient #12 carried the c.340+866C>A VUS, which is referenced in dbSNP (rs536631685) and 1000 Genomes, but not in the Genome Aggregation Database (gnomAD). Finally, three novel VUS of the E1' *VHL* cryptic exon were discovered in the three remaining patients. None of the four patients with a PPGL and an E1' VUS have developed VHL spectrum tumour(s) during their follow-up and none of them had family history of VHL disease (table 2); however, segregation analysis was only performed in patient #9. The proband's mother did carry the variant and had a pancreatic cyst and multiple vertebral body haemangiomas which both are evocating of VHL disease.<sup>17</sup>

Among the remaining 11 patients, 3 tumours were available, 2 as paraffin embedded samples (patients #1, #10) and 1 as a frozen tumour (patient #3). None of them presented a LOH at *VHL* locus and the mutated allele was lost as determined by

Sanger sequencing in tumour #1. In tumour DNA of patient #10, which harbours the c.340+682T>C variant, we identified a second variant in the exon 3 of *VHL* (c.482G>A; p.Arg161Gln), known to be pathogenic (figure 1). This somatic mutation was previously described in this patient.<sup>18</sup> In the absence of LOH, this exon 3 variant may function as the second *VHL* hit in this tumour. No other mutation of the *VHL* gene was identified in tumour DNA of patient #3.

To validate and classify these different VUS, we carried out different functional studies on available tumour tissues. We first performed immunohistochemistry to study the expression of CA9, known to be expressed at the membrane of tumour cells in case of VHL inactivation.<sup>19</sup> A membranous positive CA9 immunostaining has been previously reported in *VHL*-related PPGL, haemangioblastoma, endolymphatic tumours and ccRCC.<sup>15 20 21</sup> We observed a cluster of tumour cells with a positive membranous CA9 immunostaining only in the PPGL of patient #10 (figure 2A) which can be seen in *VHL*-related PPGL.<sup>15</sup> Then, we assessed the expression of *PNMT* gene, which is one of the most downregulated genes in *VHL*-related PPGL,<sup>22 23</sup> by RT-qPCR. As expected, the PPGL of patient #10 exhibited a significant low expression of *PNMT* mRNA, comparable to the *VHL*-tumours used as controls. On the contrary, the level of *PNMT* expression was equivalent to control tumours in the PPGL of patient #3, which produced both epinephrine and norepinephrine. Finally, we analysed the expression of *VHL* gene by RT-qPCR. We assessed the expression of two different *VHL* mRNA: the mRNA containing the exons 1 and 2 (E1-E2), which will lead with the exon 3 to the expression of the two main VHL proteins (pVHL213 and pVHL160) and *VHL* mRNA containing the exon 1 and E1' (E1-E1'), which was previously described as increased in tumour or in lymphoblastoid cell lines




**Figure 2** CA9 immunohistochemistry in patients tumours with a *VHL* E1' mutation showed a membranous immunostaining on a cluster of cells in patient #10 PPGL (A). *PNMT* and *VHL* (E1-E2 transcript) genes expression by RT-qPCR in patients #3 and #10 showed a low expression of the two genes in patient #10 tumours (B).

of patients with E1' mutation. In normal condition, this *VHL* E1-E1' mRNA is degraded by nonsense-mediated decay (NMD), and in this pathological condition, NMD may be overwhelmed. The PPGL of patients #3 and #10 showed expression of *VHL* E1-E1' mRNA which was absent in controls, suggesting that the two variants change the *VHL* mRNA splicing (figure 3). Moreover, the PPGL of patient #10 showed a low expression of *VHL* E1-E2 mRNA comparable to the *VHL*-related PPGL used as control (figure 2B). Altogether, these data provide evidence that this *VHL* E1' mutation (c.340+682T>C) is a pathogenic mutation that combined with the second mutation (c.482G>A; p.Arg161Gln) induce tumorigenesis.

Finally, 23 patients carry the c.340+648T>C (rs73024533) variant, previously described in dbSNP, at a heterozygous state. The allele frequency of rs73024533 in our cohort is comparable to that of the gnomAD database and of our control cohort of 198 European subjects (1.9% vs 1.3% and 1.3%, respectively,  $p=0.0536$ ).

## DISCUSSION

E1' mutations were previously described by Lenglet *et al* in eight families, either with erythrocytosis or VHL disease. These mutations led to an abnormal *VHL* mRNA with the insertion of the



**Figure 3** Expression of E1-E1' transcript of *VHL* gene showed expression only in patients #3 and #10.

E1' in the transcript and to mRNA degradation by NMD and to global defect in *VHL* protein expression.<sup>8</sup>

In our large international study, we identified four new germline variants in E1' *VHL* gene and we classified two of them as pathogenic, representing 1.3% of 'VHL-like' cohort (1/76 patients) and 0.11% of 'PPGL' cohort (1/946 patients). Our patients did not have all the manifestation of *VHL* disease. However, in the single patient in whom a familial genetic screening was performed (patient #9), the proband's mother had her first screening (cerebral and medullary MRI and abdominal CT scan) at the age of 70 years old, which diagnosed one pancreatic cyst and multiple vertebral body haemangiomas. Interestingly, multiple vertebral body haemangiomas are rare in *VHL* disease but have been described in patients with Chuvash polycythemia, a disease secondary to a recurrent germline biallelic mutation in *VHL* gene (c.598C>T, p.Arg200Trp).<sup>24</sup> Our data suggest incomplete penetrance of E1' *VHL* mutations, as it was previously described for the *SDHA* gene-another PPGL susceptibility gene- mutations that exhibit a relatively high allele frequency in gnomAD.<sup>25</sup>

We have identified the same variant c.340+578C>T (rs139622356) in seven patients, but our tumour analyses were not able to lead to the classification of this variant in a pathogenic variant. Indeed in one tumour with this variant, we identified the E1-E1' mRNA which suggest that the variant is pathogenic. However, epinephrine secretion and *PNMT* expression of this tumour are strong indicator against the diagnosis of *VHL*-related PPGL.<sup>26</sup> Moreover, we identified this variant in 0.6% of our

cohort, which is 10 times more frequent in our cohort than in reference databases. Indeed, this variant is described in 0.05% of gnomAD subjects (7/1167 vs 16/31 390,  $p < 0.0001$ ). It is noteworthy that in Tuscan and Iberian subjects reported in the 1000 Genomes project, the frequency of this rs139622356 is 0.9%. All these data suggest that this variant could be either a pathogenic variant that is not implicated in the PPGL of our patient because of the lack of LOH/second *VHL* mutation, or a modifier variant contributing potentially to an 8.5-fold risk (95% CI 4.4 to 14.3,  $p < 0.0001$ ) for development of PPGL or *VHL* tumours. Hence, more functional analyses and more tumours analyses will be required to achieve a definitive conclusion.

Our study demonstrates that E1' *VHL* variants are rare events in 'VHL-like' and 'PPGL' patients, but nearly as frequent as the *VHL* mutation rate in exons 1 and 2 in patients with PPGL (in the molecular genetic laboratory of Hôpital Européen Georges Pompidou-Paris-France *VHL* mutation rate in exon 1 has been reported to be 0.74% ( $p = 0.062$ ), in exon 2: 0.18% ( $p = 0.99$ ) and in exon 3: 0.92% ( $p = 0.0264$ ),<sup>27</sup> or as frequent as in exons of other PPGL susceptibility genes (for instance, the mutation rate in exon 1 of *SDHD* is 0.43%). However, because patients with well-established *VHL* pathogenic mutations were excluded from our cohort, the current frequency may be an underestimation. As the identification of *VHL* variants has important implications for management and follow-up of patients and relatives, we suggest that E1' cryptic exon should be added to NGS diagnostic panels. Considering the genetic heterogeneity of PPGLs and the high rate of detectable driver mutations in these tumours,<sup>10</sup> a low frequency of variants in any given new gene/exons is not unexpected. However, the interpretation of these E1' variants might be difficult and more functional analyses has to be designed in order to validate these variants. Finally, our study underlines the importance of variants identified in deep intronic sequences, which would have been missed by examining only coding sequences of genes/exomes. These variants will likely be more frequently detected and studied in the next future with the upcoming implementation of whole-genome sequencing into clinical practice.

#### Author affiliations

<sup>1</sup>Université de Paris, PARCC, INSERM, Equipe Labellisée par la Ligue contre le Cancer, F-75015, Paris, France

<sup>2</sup>Hereditary Endocrine Cancer Group, Spanish National Cancer Research Centre (CNIO), Madrid, Spain; and Centro de Investigación Biomédica en Red de Enfermedades Raras (CIBERER), Madrid, Spain

<sup>3</sup>Division of Hematology and Medical Oncology, Department of Medicine, Mays Cancer Center, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

<sup>4</sup>Genetics Department, Hospices Civils de LYON (HCL), University Hospital, East Biology and Pathology Center, B-A3, 59 Bld Pinel, 69677, Bron, France

<sup>5</sup>Réseau National pour Cancers Rares PREDIR labellisé par l'Institut National contre le Cancer, Hôpital Bicêtre, Assistance Publique-Hôpitaux de Paris, Le Kremlin-Bicêtre, France

<sup>6</sup>École Pratique des Hautes Études, PSL Research University, Paris, France

<sup>7</sup>L'Institut du Thorax, INSERM, Centre National de la Recherche Scientifique (CNRS), Université de Nantes, Nantes, France

<sup>8</sup>Aix Marseille Univ, APHM, INSERM, MMG, Laboratory of Molecular Biology Hospital La Conception, Marseille, France

<sup>9</sup>Endocrinology and Nutrition Service, Hospital Universitario Puerta de Hierro, 28222, Majadahonda, Spain

<sup>10</sup>Division of Endocrinology, Department of Medicine, Centre de recherche du Centre hospitalier de l'Université de Montréal (CRCHUM), Université de Montréal, Montréal, Quebec, Canada

<sup>11</sup>Gustave Roussy, Université Paris-Saclay, Département de Biopathologie and INSERM U1186, Villejuif, F-94805, France

<sup>12</sup>Department of Endocrinology and Nutrition, Hospital Universitario 12 de Octubre, 28041 Madrid, Spain

<sup>13</sup>Assistance Publique-Hôpitaux de Paris, Hôpitaux Universitaires La Pitié Salpêtrière-Charles Foix, Service de Neurologie 2-Mazarin, Paris, France

<sup>14</sup>Service ORL, Hôpital Pellegrin, CHU de Bordeaux, Bordeaux, France

<sup>15</sup>Department of Endocrinology, USHATE "Acad. Ivan Penchev", Medical University Sofia, Sofia, Bulgaria

<sup>16</sup>Assistance Publique, Hôpitaux de Paris, Service ORL-CCF, hôpital Lariboisière, université Paris VII, Paris, France

<sup>17</sup>Assistance Publique-Hôpitaux de Paris, Hôpital de Bicêtre, Service d'Endocrinologie et des Maladies de la Reproduction, Centre de Référence des Maladies Rares de l'Hypophyse, Le Kremlin Bicêtre, France

<sup>18</sup>Department of Clinical Genetics, Centre Hospitalier Régional et Universitaire de Lille, Lille, France

<sup>19</sup>Service d'Endocrinologie, Hôpital de L'Archet, CHU de Nice, Nice, France

<sup>20</sup>Génétique Oncologique EPHE, INSERM U1186, Gustave Roussy Cancer Campus, Villejuif, France

<sup>21</sup>Assistance Publique-Hôpitaux de Paris, Hôpital Européen Georges Pompidou, Service de Génétique, Paris, France

**Acknowledgements** We thank Catherine Tritscher for administrative assistance and Caroline Travers, Valentin Adamus and Astrid Ramahefasolo for technical assistance. We also thank Pr Pascal Pigny for his help and the PREDIR and TENGEN networks for their help in patient enrolment. The authors would like to express their gratitude to Sophie Devaux for her help.

**Contributors** APGR conceived and supervised the study. ABu, APGR, MR, PLD and BG designed the study and analysed the results. JF and NB participated to the data analysis. ABu., BC, SF, SG, ML and ED designed and performed the experiments. PR, JA, IB, BbP, MC, CD EdM, AE, PH, PK, SL, JLS, ABa, SR, BG, PLD, MR and APGR collected subjects and clinical parameters. ABu and APGR wrote the manuscript. ABu prepared the figures and the tables. All the authors discussed the results and commented the manuscript.

**Funding** ABu received a financial support from ITMO Cancer AVIESAN (Alliance Nationale pour les Sciences de la Vie et de la Santé, National Alliance for Life Sciences & Health) within the framework of the Cancer Plan and from la Fondation pour la Recherche Médicale (FDT20170436955). SF is currently supported by an NIH-NIGMS individual predoctoral fellowship grant (F31-GM131634-01) and, previously, by an NIH NRSA Institutional Predoctoral Training Grant T32CA148724 and NRSA F31-GM131634-01. PLD receives funding support from NIH-GM114102, Alex's Lemonade Stand Cancer Foundation (Innovation Award) and the NCATS- UL1 TR002645. The Genomic Sequencing Facility at the GCCRI is supported by the P30-CA54174 (CTRC at UTHSCSA) and NIH Shared Instrument grant 1S10OD021805-01 (S10 grant). BC is supported by Rafael del Pino Foundation. MR receives funding support from Instituto de Salud Carlos III (ISCIII), through the 'Acción Estratégica en Salud' (AES) (projects P117/01796), cofounded by the European Regional Development Fund (ERDF), and the Paradifference Foundation.

**Disclaimer** The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** All available data are in the article.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iD

Alexandre Buffet <http://orcid.org/0000-0002-4742-2708>

## REFERENCES

- Latif F, Tory K, Gnara J, Yao M, Duh FM, Orcutt ML, Stackhouse T, Kuzmin I, Modi W, Geil L. Identification of the von Hippel-Lindau disease tumor suppressor gene. *Science* 1993;260:1317–20.
- Lonser RR, Glenn GM, Walther M, Chew EY, Libutti SK, Linehan WM, Oldfield EH, disease vonH-L. Von Hippel-Lindau disease. *Lancet* 2003;361:2059–67.
- Binderup MLM, Galanakis M, Budtz-Jørgensen E, Kosteljanetz M, Luise Bisgaard M, Prevalence LBM. Prevalence, birth incidence, and penetrance of von Hippel-Lindau disease (vHL) in Denmark. *Eur J Hum Genet* 2017;25:301–7.
- Favier J, Amar L, Gimenez-Roqueplo A-P. Paraganglioma and pheochromocytoma: from genetics to personalized medicine. *Nat Rev Endocrinol* 2015;11:101–11.
- Gimenez-Roqueplo A-P, Dahia PL, Robledo M. An update on the genetics of paraganglioma, pheochromocytoma, and associated hereditary syndromes. *Horm Metab Res* 2012;44:328–33.
- Amar L, Bertherat J, Baudin E, Ajzenberg C, Bressac-de Paillerets B, Chabre O, Chamontin B, Delemer B, Giraud S, Murat A, Niccoli-Sire P, Richard S, Rohrer V, Sadoul J-L, Strompf L, Schlumberger M, Bertagna X, Plouin P-F, Jeunemaitre X, Gimenez-Roqueplo A-P. Genetic testing in pheochromocytoma or functional paraganglioma. *J Clin Oncol* 2005;23:8812–8.
- Currás-Freixes M, Piñero-Yañez E, Montero-Conde C, Apellániz-Ruiz M, Calsina B, Mancikova V, Remacha L, Richter S, Ercolino T, Rogowski-Lehmann N, Deuschlein T, Calatayud M, Guadalupe S, Álvarez-Escolá C, Lamas C, Aller J, Sastre-Marcos J, Lázaro C, Galofré JC, Patiño-García A, Meoro-Avilés A, Balmaña-Gelpi J, De Miguel-Novoa P, Balbín M, Matías-Guiu X, Letón R, Inglada-Pérez L, Torres-Pérez R, Roldán-Romero JM, Rodríguez-Antona C, Fliedner SMJ, Opocher G, Pacak K, Korpershoek E, de Krijger RR, Vroonen L, Mannelli M, Fassnacht M, Beuschlein F, Eisenhofer G, Cascón A, Al-Shahrour F, Robledo M. PheoSeq: a targeted next-generation sequencing assay for pheochromocytoma and paraganglioma diagnostics. *J Mol Diagn* 2017;19:575–88.
- Lenglet M, Robriquet F, Schwarz K, Camps C, Couturier A, Hoogewijs D, Buffet A, Knight SJL, Gad S, Couvé S, Chesnel F, Pacault M, Lindenbaum P, Job S, Dumont S, Besnard T, Cornec M, Dreau H, Pentony M, Kvikstad E, Deveaux S, Burnichon N, Ferlicot S, Vilaine M, Mazzella J-M, Airaud F, Garrec C, Heidet L, Irtan S, Mantadakis E, Bouchireb K, Debatin K-M, Redon R, Beziau S, Bressac-de Paillerets B, Teh BT, Girodon F, Randi M-L, Putti MC, Bours V, Van Wijk R, Göthert JR, Kattamis A, Janin N, Bento C, Taylor JC, Arlot-Bonnemains Y, Richard S, Gimenez-Roqueplo A-P, Cario H, Gardie B. Identification of a new *VHL* exon and complex splicing alterations in familial erythrocytosis or von Hippel-Lindau disease. *Blood* 2018;132:469–83.
- Maier ER, Neumann HP, Richard S. Von Hippel-Lindau disease: a clinical and scientific review. *Eur J Hum Genet* 2011;19:617–23.
- Toledo RA, Burnichon N, Cascon A, Benn DE, Bayley J-P, Welander J, Tops CM, Firth H, Dwight T, Ercolino T, Mannelli M, Opocher G, Clifton-Bligh R, Gimm O, Maher ER, Robledo M, Gimenez-Roqueplo A-P, Dahia PLM, NGS in PPGL (NGSNPPGL) Study Group. Consensus statement on next-generation-sequencing-based diagnostic testing of hereditary pheochromocytomas and paragangliomas. *Nat Rev Endocrinol* 2017;13:233–47.
- Lenders JWM, Duh Q-Y, Eisenhofer G, Gimenez-Roqueplo A-P, Grebe SKG, Murad MH, Naruse M, Pacak K, Young WF, Endocrine Society. Pheochromocytoma and paraganglioma: an endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2014;99:1915–42.
- Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, Grody WW, Hegde M, Lyon E, Spector E, Voelkerding K, Rehm HL, ACMG Laboratory Quality Assurance Committee. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of medical genetics and genomics and the association for molecular pathology. *Genet Med* 2015;17:405–23.
- Hoefling C, Schmidt H, Meinhardt M, Lohse A, Taubert H, Fuessel S, Schmidt U, Schuster K, Baretton G, Wirth MP, Meyer A. Comparative evaluation of microsatellite marker, AP-PCR and CGH studies in primary renal cell carcinoma. *Int J Mol Med* 2004;13:835–42.
- Girolami F, Passerini I, Gargano D, Frusconi S, Villari D, Nicta G, Torricelli F. Microsatellite analysis of chromosome 3p region in sporadic renal cell carcinomas. *Pathol Oncol Res* 2002;8:241–4.
- Favier J, Meatchi T, Robidel E, Badoual C, Sibony M, Nguyen AT, Gimenez-Roqueplo A-P, Burnichon N. Carbonic anhydrase 9 immunohistochemistry as a tool to predict or validate germline and somatic *VHL* mutations in pheochromocytoma and paraganglioma—a retrospective and prospective study. *Mod Pathol* 2020;33:57–64.
- Zeka F, Vanderheyden K, De Smet E, Cuvelier CA, Mestdagh P, Vandesompele J. Straightforward and sensitive RT-qPCR based gene expression analysis of FFPE samples. *Sci Rep* 2016;6:21418.
- van Asselt SJ, de Vries EG, van Dulleman HM, Brouwers AH, Walenkamp AM, Giles RH, Links TP. Pancreatic cyst development: insights from von Hippel-Lindau disease. *Cilia* 2013;2:3.
- Mircescu H, Wilkin F, Paquette J, Oligny LL, Decaluwe H, Gabouy L, Nolet S, Van Vliet G, Deal C. Molecular characterization of a pediatric pheochromocytoma with suspected bilateral disease. *J Pediatr* 2001;138:269–73.
- Ivanov SV, Kuzmin I, Wei MH, Pack S, Geil L, Johnson BE, Stanbridge EJ, Lerman MI. Down-Regulation of transmembrane carbonic anhydrases in renal cell carcinoma cell lines by wild-type von Hippel-Lindau transgenes. *Proc Natl Acad Sci U S A* 1998;95:12596–601.
- Pinato DJ, Ramachandran R, Toussi STK, Vergine M, Ngo N, Sharma R, Lloyd T, Meeran K, Palazzo F, Martin N, Khoo B, Dina R, Tan TM. Immunohistochemical markers of the hypoxic response can identify malignancy in pheochromocytomas and paragangliomas and optimize the detection of tumours with *VHL* germline mutations. *Br J Cancer* 2013;108:429–37.
- Stillebroer AB, Mulders PFA, Boerman OC, Oyen WJG, Oosterwijk E. Carbonic anhydrase IX in renal cell carcinoma: implications for prognosis, diagnosis, and therapy. *Eur Urol* 2010;58:75–83.
- Castro-Vega LJ, Letouzé E, Burnichon N, Buffet A, Disderot P-H, Khalifa E, Loriot C, Elarouci N, Morin A, Menara M, Lepoutre-Lussey C, Badoual C, Sibony M, Dousse B, Libé R, Zinzindohoue F, Plouin PF, Bertherat J, Amar L, de Reyniès A, Favier J, Gimenez-Roqueplo A-P. Multi-Omics analysis defines core genomic alterations in pheochromocytomas and paragangliomas. *Nat Commun* 2015;6:6044.



- 23 López-Jiménez E, Gómez-López G, Leandro-García LJ, Muñoz I, Schiavi F, Montero-Conde C, de Cubas AA, Ramires R, Landa I, Leskelä S, Maliszewska A, Inglada-Pérez L, de la Vega L, Rodríguez-Antona C, Letón R, Bernal C, de Campos JM, Díez-Tascón C, Fraga MF, Boulosa C, Pisano DG, Opocher G, Robledo M, Cascón A. Research resource: transcriptional profiling reveals different pseudohypoxic signatures in SDHB and VHL-related pheochromocytomas. *Mol Endocrinol* 2010;24:2382–91.
- 24 Gordeuk VR, Sergueeva AI, Miasnikova GY, Okhotin D, Voloshin Y, Choyke PL, Butman JA, Jedlickova K, Prchal JT, Polyakova LA. Congenital disorder of oxygen sensing: association of the homozygous Chuvash polycythemia VHL mutation with thrombosis and vascular abnormalities but not tumors. *Blood* 2004;103:3924–32.
- 25 van der Tuin K, Mensenkamp AR, Tops CMJ, Corssmit EPM, Dinjens WN, van de Horst-Schrivers ANA, Jansen JC, de Jong MM, Kunst HPM, Kusters B, Leter EM, Morreau H, van Nesselrooij BMP, Oldenburg RA, Spruijt L, Hes FJ, Timmers HJLM. Clinical aspects of SDHA-Related pheochromocytoma and paraganglioma: a nationwide study. *J Clin Endocrinol Metab* 2018;103:438–45.
- 26 Eisenhofer G, Lenders JWM, Timmers H, Mannelli M, Grebe SK, Hofbauer LC, Bornstein SR, Tiebel O, Adams K, Bratslavsky G, Linehan WM, Pacak K. Measurements of plasma methoxytyramine, normetanephrine, and metanephrine as discriminators of different hereditary forms of pheochromocytoma. *Clin Chem* 2011;57:411–20.
- 27 Ben Aim L, Pigny P, Castro-Vega LJ, Buffet A, Amar L, Bertherat J, Drui D, Guilhem I, Baudin E, Lussey-Lepoutre C, Corsini C, Chabrier G, Briet C, Faivre L, Cardot-Bauters C, Favier J, Gimenez-Roqueplo A-P, Burnichon N. Targeted next-generation sequencing detects rare genetic events in pheochromocytoma and paraganglioma. *J Med Genet* 2019;56:513–20.