



HAL
open science

Influence of substrate types and morphological traits on movement behavior in a toad and newt species

Audrey Trochet, Hugo Le Chevalier, Olivier Calvez, Alexandre Ribéron,
Romain R. Bertrand, Simon Blanchet

► **To cite this version:**

Audrey Trochet, Hugo Le Chevalier, Olivier Calvez, Alexandre Ribéron, Romain R. Bertrand, et al.. Influence of substrate types and morphological traits on movement behavior in a toad and newt species. PeerJ, 2019, 6, pp.e6053. 10.7717/peerj.6053 . hal-03032894

HAL Id: hal-03032894

<https://ut3-toulouseinp.hal.science/hal-03032894>

Submitted on 10 Dec 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Influence of substrate types and morphological traits on movement behavior in a toad and newt species

Audrey Trochet^{1,2}, Hugo Le Chevalier^{1,2}, Olivier Calvez², Alexandre Ribéron¹, Romain Bertrand^{2,3} and Simon Blanchet^{1,2}

¹ Laboratoire Evolution et Diversité Biologique, Université Paul Sabatier (Toulouse III), Toulouse, France

² Station d'Ecologie Théorique et Expérimentale, CNRS, Moulis, France

³ Center for Biodiversity Theory and Modelling, CNRS, Moulis, France

ABSTRACT

Background: Inter-patch movements may lead to genetic mixing, decreasing both inbreeding and population extinction risks, and is hence a crucial aspect of amphibian meta-population dynamics. Traveling through heterogeneous landscapes might be particularly risky for amphibians. Understanding how these species perceive their environment and how they move in heterogeneous habitats is an essential step in explaining metapopulation dynamics and can be important for predicting species' responses to climate change and for conservation policy and management.

Methods: Using an experimental approach, the present study focused on the movement behavior (crossing speed and number of stops) on different substrates mimicking landscape components (human-made and natural substrates) in two amphibian species contrasting in locomotion mode: the common toad (*Bufo bufo*), a hopping and burrowing anuran and the marbled newt (*Triturus marmoratus*), a walking salamander. We tested the hypothesis that species reaction to substrate is dependent on specific ecological requirements or locomotion modes because of morphological and behavioral differences.

Results: In both species, substrate type influenced individual crossing speed, with individuals moving faster on soil than on concrete substrate. We also demonstrated that long-legged individuals moved faster than individuals with short legs.

In both species, the number of stops was higher in females than in males. In common toads, the number of stops did not vary between substrates tested, whereas in marbled newts the number of stops was higher on concrete than on soil substrate.

Discussion: We highlighted that concrete substrate (mimicking roads) negatively affect the crossing speed of both studied species, with an effect potentially higher in marbled newts. Our findings corroborate negative effects of such heterogeneous landscapes on movement behavior of two amphibian species, which may have implications for the dynamics of metapopulations.

Submitted 2 July 2018
Accepted 29 October 2018
Published 9 January 2019

Corresponding author
Audrey Trochet,
audrey.trochet@sete.cnrs.fr

Academic editor
Donald Kramer

Additional Information and
Declarations can be found on
page 10

DOI 10.7717/peerj.6053

© Copyright
2019 Trochet et al.

Distributed under
Creative Commons CC-BY 4.0

OPEN ACCESS

Subjects Animal Behavior, Biodiversity, Conservation Biology

Keywords Matrix permeability, Inter-patch movements, Fragmented landscapes, Roads, Common toads, Bufonidae, Marbled newts, Connectivity, Salamandridae

INTRODUCTION

Inter-patch movements are key processes for maintaining gene flow among populations (Kareiva & Wennergren, 1995; Ronce, 2007) with strong consequences for metapopulation dynamics and population persistence (Clobert *et al.*, 2001, 2012; Bowler & Benton, 2005). For decades, the architecture of landscapes has profoundly changed due to human activities. Anthropogenic practices, such as agriculture, urbanization, or the expansion of road networks, have led to discontinuities in the habitat matrix: formerly continuous habitats become smaller and more isolated from each other, resulting in the well-known habitat fragmentation pattern (Collinge, 2009; Wilson *et al.*, 2016). Fragmentation and land use change could create inhospitable habitat within the current mobility range of individuals, which may force them to adapt their movement behavior (Arendt, 1988; Andreassen & Ims, 1998; Kuefler *et al.*, 2010) to ensure sufficient connectivity among populations in spite of these environmental changes. This change in movement behavior may increase the costs associated with mobility, for instance by exposing individuals to higher mortality rate during the transience phase (when crossing roads, e.g.; Carr, Pope & Fahrig, 2002). Elucidating how individuals react and adapt their movement patterns in disturbed landscapes might improve our knowledge of metapopulation dynamics across a changing landscape, increase the realism of predictions of species' responses to global change, and support the implementation of pertinent conservation plans.

One-third of the amphibian species are currently threatened worldwide, with 43% of species having declined over the last few decades (Stuart *et al.*, 2004). Due to ecological requirements, pond- and stream-breeding amphibians are exposed to different habitat types (i.e., both terrestrial and aquatic) throughout their life cycle, often in patchy and heterogeneous landscapes (Marsh & Trenham, 2001). During the terrestrial phase, individual movements are risky, due to through predator and UV-B exposure (Kats *et al.*, 2000) and desiccation risk (Rothermel & Luhring, 2005; Pittman *et al.*, 2013; Pittman, Osbourn & Semlitsch, 2014). Many studies have considered the multiple effects of habitat fragmentation—and their related landscape components—on amphibian populations, both at the individual and population levels (see Cushman, 2006 for a review). In particular, urban areas and human-made infrastructure, such as roads, negatively affect these species in two ways: directly, through fatal collisions with vehicles (Fahrig *et al.*, 1995; Rytwinski & Fahrig, 2015) and indirectly by fragmenting habitat and subsequent reduction of gene flow and colonization events (DeMaynadier & Hunter, 2000; Mazerolle, 2004; Marsh *et al.*, 2005; Cosentino, Schooley & Phillips, 2011; Youngquist *et al.*, 2016; Lenhardt *et al.*, 2017). Amphibians are more susceptible to fatal collisions when crossing roads during migration (Glista, DeVault & DeWoody, 2008) because they often become immobile when facing an approaching vehicle (Mazerolle, Huot & Gravel, 2005; Bouchard *et al.*, 2009). Also, the mobility of crossing individuals could be affected by the nature of the road's substrate. In this vein, several studies investigated the effects of different substrates on individual movements at the intra-specific level (Ims & Yoccoz, 1997; Wiens, Schooley & Weeks, 1997; Wiens, 2001; Stevens *et al.*, 2006; Semlitsch *et al.*, 2012). For instance in *Plethodon metcalfi*, the individual crossing speed was

higher on asphalt than on grass (Semlitsch et al., 2012) with various consequences on mobility success. Some amphibian species can also benefit from anthropogenic landscape elements. For example, *Rhinella marina* (cane toads) seemed to use roads as dispersal corridors in Australia (Brown et al., 2006). Despite the crucial importance of inter-patch movements in altered landscapes, little is known about the multi-species interactions among individual movements and different substrates (but see Rothermel & Semlitsch, 2002). Indeed, habitat-species interactions are complex and highly specific (Kolozsvary & Swihart, 1999; Trochet et al., 2016a). Heterogeneous landscapes and associated landscape parameters could differently alter movement of organisms with strong divergence between species (Rothermel & Semlitsch, 2002) resulting in contrasting conservation needs and species suitability under global change.

The costs associated with inter-patch movement can be high (Van Dyck & Baguette, 2005) and could lead to high selective pressures on mobility and associated phenotypic traits (Bonte et al., 2012). According to this expectation, many studies have focused on the correlation between movement and phenotypic traits. At the intra-specific level, phenotypic differences related to dispersal ability between individuals have been reported. For instance, larger and/or longer individuals are generally expected to be dispersers, because they should benefit from their high levels of competition to disperse farther (Léna et al., 1998). Evidence for a relationship between body size and movement has been described in many taxa (insects: Anholt, 1990; Legrand et al., 2015; mammals: Gundersen, Andreassen & Ims, 2002; Holekamp & Sherman, 1989; O'Riain, Jarvis & Faulkes, 1996; reptiles: Léna et al., 1998; birds: Barbraud, Johnson & Bertault, 2003; Delgado et al., 2010; fishes: Radinger & Wolter, 2014; amphibians: Ousterhout & Semlitsch, 2018). For walking and/or hopping animals, selection for efficient displacement might lead to leg elongation. As a result, morphological adaptation to movement can be deduced from estimates of leg length (Moya-Laraño et al., 2008). This correlation between movement and leg length (i.e., hind-limb length, hereafter HLL) has been demonstrated in some species (reptiles: Losos, 1990; spiders: Moya-Laraño et al., 2008; amphibians: Bennett, Garland & Else, 1989; Choi, Shim & Ricklefs, 2003; Phillips et al., 2006), but still remains rare.

Understanding how amphibians perceive their environment and how they actually move in heterogeneous habitats is an essential step for understanding metapopulation structure, and can be important for improving the realism of predictive models of species' responses to global change. The present study focused on the movement behavior (crossing speed) on two different substrates mimicking landscape components (a human-made substrate and a natural substrate) in two contrasting amphibian species, the common toad (*Bufo bufo*), a hopping and burrowing anuran, and the marbled newt (*Triturus marmoratus*), a walking salamander. Each species has specific ecological requirements: common toads have an explosive breeding season and spend a large proportion of their life on terrestrial habitats (i.e., forests, bushlands, or urban areas) whereas the breeding season of marbled newts is longer, and this species inhabits relatively small well-vegetated ponds surrounded by woodlands (Jehle & Arntzen, 2000; AmphibiaWeb, 2012). Contrary to common toads that generally avoid this kind of habitat

type, marbled newts also occur in more open areas like heathens and agricultural landscapes. Common toads and marbled newts can live within the same habitat, such as grassland or woodland, and could therefore face the same environmental pressures during terrestrial movements ([Daverson, Muths & Bosch, 2012](#); [Trochet et al., 2017](#)). Also, both the contrasting morphology and modes of locomotion induce different muscular contractions during displacement, leading to different mobility ability between the two species ([Smith & Green, 2005](#)) and therefore explain the selection of various habitats encounter during terrestrial phase.

Our study aimed at testing the influence of different substrate types on two amphibian species in order to highlight if and how substrates can alter movement in these sympatric species. Considering these species allowed testing the hypothesis that species could react differently to substrate nature, depending on specific ecological requirements, mobility abilities, or locomotion modes because of morphological and behavioral differences. Identifying such differences could help improving our understanding about species-specific interactions within a human-made environment. Consequently, it would be interesting to consider those interactions to improve predictive models of species' responses to climate change and to propose efficient conservation management plans.

MATERIALS AND METHODS

Studied species

The common toad is one of the most widely distributed and abundant anuran species in Europe ([Gasc et al., 1997](#)). Usually, reproduction occurs in February, and large numbers of toads migrate to breeding sites (i.e., large ponds, ditches, or lakes with relatively clear water, quite variable in area and depth; [AmphibiaWeb, 2012](#)) where the males compete for mating. After an explosive breeding season, toads leave ponds and return to terrestrial habitats ([Gittins, 1983](#)) where they spend a large proportion of their life. This species occupies not only various terrestrial habitats over three to four km from the breeding site ([Smith & Green, 2005](#)), such as coniferous, mixed and deciduous forests, bushlands, but also urban areas such as gardens and parks ([Nollert & Nollert, 2003](#)). Common toads hibernate singly or in groups from September to February, on land and occasionally in streams and springs.

The marbled newt is a large-bodied urodele species from Western Europe, found in France, Spain, and Portugal ([Sillero et al., 2014](#)). The reproduction period is longer than common toads (from the beginning of March to July) and takes place in different aquatic habitats, including well-vegetated ponds, pools, ditches, and streams ([AmphibiaWeb, 2017](#)). After breeding, adults leave water bodies by walking, and deciduous or mixed woodland, where they find refuges under dead and rotting wood and other hiding places ([Jehle & Arntzen, 2000](#)). Displacement distances of newts around their breeding sites are shorter than anurans and in the range of several hundred meters ([Trochet et al., 2017](#)).

Sampling and morphological measurements

Our work complies with the international animal care guidelines of the Association for the Study of Animal Behaviour, and all required French permits relating to authorization of

capture, marking, transport, detention, use, and release of protected amphibian species have been obtained (permit nos. 09-2014-14 and 32-2014-07; animal experimentation accreditation no. A09-1) from the DREAL Occitanie (“Direction Régionale de l’Environnement, de l’Aménagement et du Logement”). Ethical approval was included under the protected species handling permit from the DREAL Occitanie. The project was approved by the “Conseil National de la Protection de la Nature” on September 14, 2014 and by the “Conseil Scientifique Régional du Patrimoine Naturel (CSRPN)” of the region Midi-Pyrénées on October 14, 2014.

In total, 83 common toads (68 males and 15 non-gravid females) and 46 marbled newts (23 males and 23 non-gravid females) were captured in the south of France at the end of breeding season (for toads: from March 13, 2015 to March 20, 2015; for newts: from April 28, 2015 to May 6, 2015) within and near two different ponds to reduce potential impact on populations (geographical coordinates of pond 1 43.671781°N, 0.504308°E and pond 2: 43.076347°N, 1.351639°E). Individuals were then brought to the lab for experimentation and released between June and July 2015. During experiments, animals were housed at the Station d’Ecologie Théorique et Expérimentale (Moulis, France) in same-species groups of four to six individuals in semi-aquatic terraria of 60 × 30 × 30 cm at 20 °C. They were fed ad libitum with live mealworms and tubifex worms. For unambiguous identification, all individuals were PIT-tagged (RFID Standards ISO 11784 & 11785 type FDX-B, 1.4 × 8 mm, 134.2 khz from BIOLOG-ID, France; animal experimentation accreditation no. A09-1) immediately after capture and before the experiments following the protocol developed in *Le Chevalier et al. (2017)*. We then measured snout-to-vent length (SVL) and HLL to the nearest one mm and body weight (mass) to the nearest 0.01 g. To limit the effects of stress on behavioral responses, all individuals were kept in captivity for several days without any manipulation until the experimental tests.

Movement tests

All tests were performed in June and July 2015, after the breeding season when all individuals were in the terrestrial phase. In order to test the crossing capacities of both species, we made them move along two tracks (200 cm long × 10 cm wide × 20 cm high), each filled with a different substrate: smooth concrete slab (human-made) or soil (natural). Tracks were not moistened so that the substrate was unfavorable. During the experiments, all individuals were chased down the tracks and forced to move by gently poking their back with a finger after each stop (3 s between pokes if needed). Only one individual was tested at a time and we recorded the number of stops (*stops*) and the *crossing speed* (in cm/s; including stops) to the nearest 0.1 s to travel 200 cm from departure to arrival line. In order to provide reliable estimates of crossing capacity using a repeated-measure design while minimizing stress, every individual was tested three times on each substrate. In toads, tests were randomly spread over 2 days (three trials on day 1 and three trials on day 2; from March 18, 2015 to March 31, 2015) while newts, for which we kept animals during a longer period for another experiment not detailed

here, tests were randomly performed over 27.9 ± 8.5 days (mean \pm SD; from April 28, 2015 to June 2, 2015). Outside of test periods, animals were returned to the semi-aquatic aquaria. Because locomotion in amphibians is influenced by temperature (Herrel & Bonneaud, 2012; James et al., 2012; Šamajová & Gvoždík, 2010), all tests were performed in a greenhouse under controlled-temperature conditions (mean \pm SD: 25 ± 1 °C) with light conditions similar to nature.

Some individuals ($n = 32$) did not complete the 200 cm run (stopping completely—three successive pokes without moving—or turning back); these individuals and their replicates were removed from the analyses for statistical reasons. We therefore included 77 toads (77 toads \times 3 replicates \times 2 substrates = 462 tests) and 20 marbled newts (20 newts \times 3 replicates \times 2 substrates = 120 tests) in the analyses.

Statistical analyses

For each species, the influence of substrate (concrete and soil) on crossing speed was first tested using Wilcoxon rank-sum tests and then modeled using linear mixed-effect models (LMMs). Mass and HLL were strongly related to SVL ($r_s = 0.866$, $P < 0.001$, and $r_s = 0.691$, $P < 0.001$, respectively). To avoid collinearity in our model, we used the scaled mass index (M_i) of condition using the following equation:

$$M_i = M \times (\text{SVL}_0/\text{SVL})^{b_{\text{SMA}}}$$

where M and SVL are the body mass and the snout–vent length of the individual, respectively. SVL_0 is the mean SVL of the population, and b_{SMA} is the standardized major axis slope from the OLS regression of log-transformed body mass on log-transformed SVL divided by Pearson's correlation coefficient (Peig & Green, 2009). We also used the relative size of the HLL (hereafter *leg*) estimated as the residuals of the linear regression between HLL and SVL.

We built linear mixed-effect models (LMMs) using the *crossing speed* (log-transformed) as response variable, individual as a random factor and M_i , *leg*, *substrate*, *sex*, *species* and first-order interactions between *species* and other variables as fixed effects. Because the *crossing speed* was strongly related to the number of stops in both species (Spearman correlations: *T. marmoratus*: $r_s = -0.543$, $P < 0.001$; *B. bufo*: $r_s = -0.767$, $P < 0.001$), we also added *stops* as covariate in our models. We then built a second LMM using *stops* as response variable, individual as a random factor and M_i , *leg*, *substrate*, *sex*, *species* and first-order interactions between *species* and other variables as fixed effects. LMMs were performed using the lme4 R-package (Bates et al., 2017).

Model selection was performed using the Akaike information criteria (AIC; Burnham & Anderson, 2002). If several best models were retained ($\Delta\text{AIC} < 2$), we used a model averaging procedure among all possible models to determine the relative importance of each selected variable. Two parameters from this averaging procedure were retained to test the importance of each variable on the *crossing speed*: the confidence interval of the averaged estimated slope of the selected term (high effects had confidence intervals that did not include zero) and the relative weight of the term (i.e., the relative Akaike weight of the top-ranked models ($\Delta\text{AIC} < 2$) in which the term appeared). All statistical analyses were performed using R v.3.1.0 (R Development Core Team, 2014).

Table 1 Summary of the averaged linear mixed-effects model showing the influence of significant variables on the crossing speed (in cm/s) for the studied species the marbled newt (*Triturus marmoratus*; $N = 20$) and the common toad (*Bufo bufo*; $N = 77$).

	Estimate	<i>P</i>	95% CI of estimate	Weight
(Intercept)	2.2793	<0.001***	(2.2153–2.3433)	
Substrate (soil)	0.0722	<0.001***	(0.0378–0.1067)	1.00
Species (<i>Triturus marmoratus</i>)	−1.0940	<0.001***	(−1.3533 to −0.8347)	1.00
<i>leg</i>	0.0998	0.0045**	(0.0309–0.1687)	0.62
<i>stops</i>	−0.2526	<0.001***	(−0.2754 to −0.2298)	1.00

Notes:

leg: relative hind-limb length. *stops*: number of stops. Weight: relative Akaike weight of the top-ranked models ($\Delta AIC < 2$) in which the term appeared.

*** $P < 0.001$,

** $P < 0.01$.




Figure 1 Mean crossing speed (in cm/s) on concrete and soil substrates in the common toad and the marbled newt. Crossing speed differed significantly between both substrates in toads (Wilcoxon rank-sum test: $P = 0.0101$) and in newts (Wilcoxon rank-sum test: $P < 0.001$). Error bars represent standard error. Notes: *** $P < 0.001$, ** $P < 0.01$. [Full-size !\[\]\(4c0374441789aff1845b5172a6ac367a_img.jpg\) DOI: 10.7717/peerj.6053/fig-1](https://doi.org/10.7717/peerj.6053/fig-1)

RESULTS

After model selection, the best models explaining variation in *crossing speed* retained substrates (soil and concrete), species, *legs*, and *stops* variables (Table 1; Supplementary Material 1). *Crossing speed* was significantly lower in marbled newts than in common toads, and both species moved faster on soil than on concrete (Table 1; Fig. 1). *Crossing speed* was also correlated with morphological traits. Long-legged individuals moved faster than individuals with short legs (Table 1).

Table 2 Summary of the best linear mixed-effect model showing the influence of significant variables on the number of stops to travel 200 cm during the experimental test for the marbled newt (*Triturus marmoratus*; $N = 20$) and the common toad (*Bufo bufo*; $N = 77$).

	Estimate	P
(Intercept)	0.7227	<0.001 ***
Substrate (soil)	-0.1212	0.0744
Species (<i>Triturus marmoratus</i>)	-0.9422	<0.001 ***
Sex (male)	-0.5453	0.0009 ***
Substrate (soil)* species (<i>Triturus marmoratus</i>)	-0.3937	0.0086 **

Notes:

*** $P < 0.001$,

** $P < 0.01$.

The variable *stops* was also influenced by species and substrates, and by sex and the interaction between substrates and species (Table 2). In both species, the number of stops was higher in females than in males. In common toads, *stops* did not vary between substrate types, whereas in marbled newts the number of stops was significantly higher on concrete than on soil (Table 2).

DISCUSSION

Our results demonstrated that both species were affected by substrate types, moving significantly slower on a human-made (concrete) than on a natural (soil) substrate. Crossing speed was also related to a morphological trait, with long-legged individuals moving faster than individuals with shorts legs. In addition, we found that the number of stops to cross 200 cm was influenced by sexes, substrates, and species, which indicates that substrates could differently affect the mobility of two sympatric species living into similar habitats.

Influence of substrate type on crossing speed

Inter-patch movement is expected to depend on the nature of the substrate crossed. Some landscape features may be associated with high resistance to movement while others facilitate movement (low resistance). In a previous study, *Stevens et al. (2006)* experimentally demonstrated that the natterjack toad (*B. calamita*) significantly preferred substrates mimicking forest and soil than those mimicking agricultural lands. Another similar study revealed that in *P. metcalfi*, individual crossing speed was higher on asphalt than on grass and soil (*Semlitsch et al., 2012*) with various consequences on mobility success.

In our experiment, the concrete substrate represented linear roads, both in its nature (mixture of cement and gravel) and length (two m wide road). Roads constitute a very hostile environment for amphibians (dry and warm substrate that could induce a desiccation risk, collisions with vehicles and/or habitat fragmentation; *Fahrig et al., 1995*; *DeMaynadier & Hunter, 2000*; *Mazerolle, 2004*; *Marsh et al., 2005*; *Cosentino, Schooley & Phillips, 2011*; *Youngquist et al., 2016*; *Lenhardt et al., 2017*). According to our hypotheses, our results showed that substrate type influenced the movement behavior of both species. Contrary to previous studies suggesting that terrestrial amphibians moved

more quickly and directly through unfavorable areas in which they are physiologically stressed (*Semlitsch et al., 2012; Peterman et al., 2014*), we revealed that both common toads and marbled newts moved faster (i.e., higher crossing speed) on soil than on concrete (Fig. 1; Table 1). We then suggest that both species could be more exposed to traffic, and suffer more from both desiccation and mortality risks on roads than on soil (*Petronilho & Dias, 2005; Santos et al., 2007; Sillero, 2008; Bouchard et al., 2009; Elzanowski et al., 2009; Matos, Sillero & Argaña, 2012*).

In both species, the number of stops was higher in females than in males, independently of substrate type (Table 2). Such differences between sexes may be driven by divergent breeding benefits, which could lead to a trade-off between movement and high energetic costs of reproduction in females, and the well-known sex-biased dispersal (see *Trochet et al., 2016b*). Indeed, some studies found male-biased dispersal in amphibian species, in anurans as well as in urodela, suggesting that females are philopatric in these organisms (*Austin et al., 2003; Lampert et al., 2003; Liebgold, Brodie & Cabe, 2011*). In our study, males seem more likely to disperse because they stopped less often than females. As a result, males could also have lower mortality risks induced by collisions with vehicles than females when crossing roads, which may have strong consequences on population dynamics. Our results also demonstrated that the number of stops did not vary between substrates in common toads, whereas marbled newts stopped more often on concrete than on soil (Table 2). This last result shows that marbled newts may be even more at-risk than toads on pavement (and then on roads) because both their crossing speed and their number of stops increased in this substrate type.

In the context of a fragmented landscape, our results corroborate the negative effects of urbanization and human-made infrastructures such as roads on amphibians, leading to an increase in population extinction risk (*Fahrig et al., 1995; Carr, Pope & Fahrig, 2002; Sotiropoulos et al., 2013*). Here, we highlight a direct influence of the substrate on the displacement of two amphibian species with divergent ecological requirements and locomotion modes living within similar habitats. Our findings are relevant for increased realism of dispersal in predictive modeling, notably by informing landscape permeability for species, body size distribution, and sex-biased dispersal. They also emphasize the importance of road-crossing structure and landscape management at a small spatial scale for amphibian conservation.

Movement-related traits in both species

According to our expectations, our results showed an influence of a morphological trait (*leg*) on the crossing speed in both species. Various morphological variables enable organisms to be adapted for ecologically effective movement (*Bennett, Garland & Else, 1989; Losos, 1990; Choi, Shim & Ricklefs, 2003; Phillips et al., 2006; Moya-Laraño et al., 2008*), such as HLL. A recent meta-analysis among several anuran species actually demonstrate that jumping performances were strongly correlated to HLL after correcting by SVL (*Gomes et al., 2009*). According to this finding, we also demonstrated that movement behavior (defined here as crossing speed) was related to the limb length in both species (see also *Bennett, Garland & Else, 1989; Choi, Shim & Ricklefs, 2003*;

Phillips et al., 2006). Indeed, long-legged individuals moved faster than individuals with short legs ([Table 1](#)), which corroborates the idea that limb length may be tightly associated with movement behavior adaptations in amphibians. Longer legs could facilitate more rapid or longer-distance displacement (*Phillips et al., 2006*), as well as generating other advantages such as improved predator evasion and simplifying the negotiation of barriers and obstacles. As a consequence, the mortality risk of longer-legged individuals could be lower than individuals with short legs.

Limitations of the study

Our experimental protocol is easy-to-use and repeatable but, because of its simplicity, may fail to identify all aspects of the complexity of amphibian's terrestrial movements. Indeed, our results highlight the impact of substrate nature of human-made infrastructures on amphibian movements at a small spatial scale, but in order to validate and understand the consequences of such an impact, field studies (using capture-mark-recapture or telemetry monitoring) are needed. Also, our studies focus on adults, while most dispersal events are likely to be ensured by juveniles in amphibian species (*Semlitsch, 2008*). Also, it could be very interesting to test the variation of crossing speed in these organisms under several temperature and humidity conditions. Studying terrestrial displacements of amphibian species on different life stages (larvae, juveniles, and adults) and phases of their life cycle (breeding migrations, dispersal events, etc.) would provide a better identification of the impact of human-made infrastructures on the ecology of amphibian communities, and therefore improve the efficiency of management and conservation efforts.

CONCLUSIONS

Inter-patch movement is a multifaceted process, subject to internal and external biotic and abiotic factors. Our findings demonstrate effects of substrates to cross them on the movement behavior in two contrasting amphibian species living within similar habitats. In particular, individuals moved slower on concrete, making them more vulnerable on roads. In both species, we also showed significant relationship between a morphological trait (*leg*) and crossing speed, suggesting that long-legged amphibians could cross human-made infrastructures faster, which could reduce mortality risk. Comparing the potential influence of various substrates individual movements are essential for explaining and predicting the dynamics of metapopulation living in strongly altered landscapes, which is a prerequisite for developing appropriate conservation management plans.

ADDITIONAL INFORMATION AND DECLARATIONS

Funding

Financial support was provided by a post-doctoral fellowship to Audrey Trochet from the Fondation de France. This work was supported by the French Laboratory of Excellence project "TULIP" (ANR-10-LABX-41; ANR-11-IDEX-0002-02) and benefited from both

“Investissement d’Avenir” grants managed by Agence Nationale de la Recherche (AnaEE, ANR-11-INBS-0001AnaEE-Services). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Grant Disclosures

The following grant information was disclosed by the authors:

Fondation de France.

French Laboratory of Excellence project “TULIP”: ANR-10-LABX-41; ANR-11-IDEX-0002-02.

“Investissement d’Avenir” grants managed by Agence Nationale de la Recherche: AnaEE, ANR-11-INBS-0001AnaEE-Services.

Competing Interests

The authors declare that they have no competing interests.

Author Contributions

- Audrey Trochet conceived and designed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Hugo Le Chevalier conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, approved the final draft.
- Olivier Calvez contributed reagents/materials/analysis tools, approved the final draft.
- Alexandre Ribéron conceived and designed the experiments, approved the final draft.
- Romain Bertrand analyzed the data, contributed reagents/materials/analysis tools, approved the final draft.
- Simon Blanchet conceived and designed the experiments, contributed reagents/materials/analysis tools, approved the final draft.

Animal Ethics

The following information was supplied relating to ethical approvals (approving body and any reference numbers):

The project was approved by the “Conseil National de la Protection de la Nature” on September 14, 2014 and by the “Conseil Scientifique Régional du Patrimoine Naturel (CSRPN)” of the region Midi-Pyrénées on October 14, 2014. Ethical approval was included under the protected species handling permit from the DREAL Occitanie (“Direction Régionale de l’environnement, de l’Aménagement et du Logement”). Permit nos.: 09-2014-14 and 32-2014-07.

Field Study Permissions

The following information was supplied relating to field study approvals (i.e., approving body and any reference numbers):

Field experiments were approved by the DREAL Occitanie (“Direction Régionale de l’environnement, de l’Aménagement et du Logement”). French permits relating to an

authorization of capture, marking, transport, detention, use and release of protected amphibian species have been obtained (permit nos. 09-2014-14 and 32-2014-07).

Data Availability

The following information was supplied regarding data availability:

Raw data is provided in a [Supplementary File](#).

Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.6053#supplemental-information>.

REFERENCES

- AmphibiaWeb. 2012.** *Bufo bufo*: Common Toad. Berkeley: University of California. Available at <http://amphibiaweb.org/species/127> (accessed 8 August 2018).
- AmphibiaWeb. 2017.** *Triturus marmoratus*: Marbled Newt. Berkeley: University of California. Available at <http://amphibiaweb.org/species/4300> (accessed 8 August 2018).
- Andreassen HP, Ims RA. 1998.** The effects of experimental habitat destruction and patch isolation on space use and fitness parameters in female root vole *Microtus oeconomus*. *Journal of Animal Ecology* **67**(6):941–952 DOI [10.1046/j.1365-2656.1998.6760941.x](https://doi.org/10.1046/j.1365-2656.1998.6760941.x).
- Anholt B. 1990.** Size-biased dispersal prior to breeding in a damselfly. *Oecologia* **83**(3):385–387 DOI [10.1007/BF00317564](https://doi.org/10.1007/BF00317564).
- Arendt WJ. 1988.** Range expansion of the cattle egret (*Bubulcus ibis*) in the Greater Caribbean basin. *Colonial Waterbirds* **11**(2):252–262 DOI [10.2307/1521007](https://doi.org/10.2307/1521007).
- Austin JD, Davila JA, Loughheed SC, Boag P. 2003.** Genetic evidence for female-biased dispersal in the bullfrog, *Rana catesbeiana* (Ranidae). *Molecular Ecology* **12**(11):3165–3172 DOI [10.1046/j.1365-294X.2003.01948.x](https://doi.org/10.1046/j.1365-294X.2003.01948.x).
- Barbraud C, Johnson AR, Bertault G. 2003.** Phenotypic correlates of post-fledging dispersal in a population of greater flamingos: the importance of body condition. *Journal of Animal Ecology* **72**(2):246–257 DOI [10.1046/j.1365-2656.2003.00695.x](https://doi.org/10.1046/j.1365-2656.2003.00695.x).
- Bates D, Maechler M, Bolker B, Walker S. 2017.** *lme4: Linear mixed-effects models using Eigen and S4 classes*. R Package version 1.1-13. Available at <https://github.com/jjallaire/lme4>.
- Bennett AF, Garland T, Else PL. 1989.** Individual correlation of morphology, muscle mechanics, and locomotion in a salamander. *American Journal of Physiology* **256**(6):R1200–R1208 DOI [10.1152/ajpregu.1989.256.6.R1200](https://doi.org/10.1152/ajpregu.1989.256.6.R1200).
- Bonte D, Van Dyck H, Bullock JM, Coulon A, Delgado M, Gibbs M, Lehouck V, Matthysen E, Mustin K, Saastamoinen M, Schtickzelle N, Stevens VM, Vandewoestijne S, Baguette M, Barton K, Benton TG, Chaput-Bardy A, Clobert J, Dytham C, Hovestadt T, Meier CM, Palmer SCF, Turlure C, Travis JM. 2012.** Costs of dispersal. *Biological Reviews* **87**(2):290–312 DOI [10.1111/j.1469-185X.2011.00201.x](https://doi.org/10.1111/j.1469-185X.2011.00201.x).
- Bouchard J, Ford AT, Eigenbrod FE, Fahrig L. 2009.** Behavioral responses of northern leopard frogs (*Rana pipiens*) to roads and traffic: implications for population persistence. *Ecology and Society* **14**(2):23 DOI [10.5751/ES-03022-140223](https://doi.org/10.5751/ES-03022-140223).
- Bowler D, Benton T. 2005.** Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. *Biological Reviews* **80**(2):205–225 DOI [10.1017/S1464793104006645](https://doi.org/10.1017/S1464793104006645).
- Brown GP, Phillips BL, Webb JK, Shine R. 2006.** Toad on the road: use of roads as dispersal corridors by cane toads (*Bufo marinus*) at an invasion front in tropical Australia. *Biological Conservation* **133**(1):88–94 DOI [10.1016/j.biocon.2006.05.020](https://doi.org/10.1016/j.biocon.2006.05.020).

- Burnham K, Anderson D. 2002.** *Model selection and multimodel inference: a practical information-theoretic approach*. Second Edition. New York: Springer Verlag.
- Carr L, Pope S, Fahrig L. 2002.** Impacts of landscape transformation by roads. In: Gutzwiller KJ, ed. *Applying Landscape Ecology in Biological Conservation*. New York: Springer Verlag, 225–243.
- Choi I, Shim JH, Ricklefs RE. 2003.** Morphometric relationships of take-off speed in anuran amphibians. *Journal of Experimental Zoology* **299A(2)**:99–102
DOI [10.1002/jez.a.10293](https://doi.org/10.1002/jez.a.10293).
- Clobert J, Baguette M, Benton TG, Bullock JM. 2012.** *Dispersal ecology and evolution*. Oxford: Oxford University Press.
- Clobert J, Danchin E, Dhondt AA, Nichols JD. 2001.** *Dispersal*. Oxford: Oxford University Press.
- Collinge SK. 2009.** *Ecology of fragmented landscapes*. Baltimore: Johns Hopkins University Press.
- Cosentino BJ, Schooley RL, Phillips CA. 2011.** Spatial connectivity moderates the effect of predatory fish on salamander metapopulation dynamics. *Ecosphere* **2(8)**:1–14
DOI [10.1890/ES11-00111.1](https://doi.org/10.1890/ES11-00111.1).
- Cushman S. 2006.** A effects of habitat loss and fragmentation on amphibians: a review and prospectus. *Biological Conservation* **128(2)**:231–240 DOI [10.1016/j.biocon.2005.09.031](https://doi.org/10.1016/j.biocon.2005.09.031).
- Daversa DR, Muths E, Bosch J. 2012.** Terrestrial movement patterns of the Common Toad (*Bufo bufo*) in Central Spain reveal habitat of conservation importance. *Journal of Herpetology* **46(4)**:658–664 DOI [10.1670/11-012](https://doi.org/10.1670/11-012).
- Delgado M, Penteriani V, Revilla E, Nams VO. 2010.** The effect of phenotypic traits and external cues on natal dispersal movements. *Journal of Animal Ecology* **79(3)**:620–632
DOI [10.1111/j.1365-2656.2009.01655.x](https://doi.org/10.1111/j.1365-2656.2009.01655.x).
- DeMaynadier PG, Hunter ML Jr. 2000.** Road effects on amphibian movements in a forested landscape. *Natural Areas Journal* **20**:56–65.
- Elzanowski A, Ciesiołkiewicz J, Kaczor M, Radwańska J, Urban R. 2009.** Amphibian road mortality in Europe: a meta-analysis with new data from Poland. *European Journal of Wildlife Research* **55(1)**:33–43 DOI [10.1007/s10344-008-0211-x](https://doi.org/10.1007/s10344-008-0211-x).
- Fahrig L, Pedlar JH, Pope SE, Taylor PD, Wegner JF. 1995.** Effect of road traffic on amphibian density. *Biological Conservation* **73(3)**:177–182 DOI [10.1016/0006-3207\(94\)00102-V](https://doi.org/10.1016/0006-3207(94)00102-V).
- Gasc JP, Cabela A, Crnobrnja-Isailovic J, Dolmen D, Grossenbacher K, Haffner P, Lescure J, Martens H, Martinez-Rica JP, Maurin H, Oliveira ME, Sofianidou TS, Veith M, Zuiderwijk A. 1997.** *Atlas of amphibians and reptiles in Europe*. Paris: Muséum National d'Histoire Naturelle.
- Gittins SP. 1983.** The breeding migration of the Common Toad (*Bufo bufo*) to a pond in mid-Wales. *Journal of Zoology* **199(4)**:555–562 DOI [10.1111/j.1469-7998.1983.tb05106.x](https://doi.org/10.1111/j.1469-7998.1983.tb05106.x).
- Glista DJ, DeVault TL, DeWoody JA. 2008.** Vertebrate road mortality predominantly impacts amphibians. *Herpetological Conservation and Biology* **3**:77–87.
- Gomes FR, Rezende EL, Grizante MB, Navas CA. 2009.** The evolution of jumping performance in anurans: morphological correlates and ecological implications. *Journal of Evolutionary Biology* **22(5)**:1088–1097 DOI [10.1111/j.1420-9101.2009.01718.x](https://doi.org/10.1111/j.1420-9101.2009.01718.x).
- Gundersen G, Andreassen HP, Ims RA. 2002.** Individual and population-level determinants of immigration success on local habitat patches: an experimental approach. *Ecology Letters* **5(2)**:294–301 DOI [10.1046/j.1461-0248.2002.00320.x](https://doi.org/10.1046/j.1461-0248.2002.00320.x).
- Herrel A, Bonneaud C. 2012.** Temperature dependence of locomotor performance in the tropical clawed frog, *Xenopus tropicalis*. *Journal of Experimental Biology* **215(14)**:2465–2470
DOI [10.1242/jeb.069765](https://doi.org/10.1242/jeb.069765).

- Holekamp K, Sherman P. 1989.** Why male ground squirrels disperse. *American Scientist* 77:232–239.
- Ims RA, Yoccoz NG. 1997.** Studying transfer processes in metapopulations: emigration, migration, and colonization. In: Hanski IA, Gilpin ME, eds. *Metapopulation Biology: Ecology, Genetics, and Evolution*. London: Academic Press, 247–265.
- James RS, Tallis J, Herrel A, Bonneaud C. 2012.** Warmer is better: thermal sensitivity of both maximal and sustained power output in the iliotibialis muscle isolated from adult *Xenopus tropicalis*. *Journal of Experimental Biology* 215(3):552–558 DOI 10.1242/jeb.063396.
- Jehle R, Arntzen JW. 2000.** Post-breeding migrations of newts (*Triturus cristatus* and *T. marmoratus*) with contrasting ecological requirements. *Journal of Zoology* 251(3):297–306 DOI 10.1111/j.1469-7998.2000.tb01080.x.
- Kareiva P, Wennergren U. 1995.** Connecting landscape patterns to ecosystem and population processes. *Nature* 373(6512):299–302 DOI 10.1038/373299a0.
- Kats LB, Kiesecker JM, Chivers DP, Blaustein AR. 2000.** Effects of UV-B on antipredator behavior in three species of amphibians. *Ethology* 106(10):921–932 DOI 10.1046/j.1439-0310.2000.00608.x.
- Kolozsvary MB, Swihart RK. 1999.** Habitat fragmentation and the distribution of amphibians: patch and landscape correlates in farmland. *Canadian Journal of Zoology* 77(8):1288–1299 DOI 10.1139/z99-102.
- Kuefler D, Hudgens B, Haddad NM, Morris WF, Thurgate N. 2010.** The conflicting role of matrix habitats as conduits and barriers for dispersal. *Ecology* 91(4):944–950 DOI 10.1890/09-0614.1.
- Lampert KP, Rand AS, Mueller UG, Ryan MJ. 2003.** Fine-scale genetic pattern and evidence for sex-biased dispersal in the túngara frog, *Physalaemus pustulosus*. *Molecular Ecology* 12(12):3325–3334 DOI 10.1046/j.1365-294X.2003.02016.x.
- Le Chevalier H, Calvez O, Martinez-Silvestre A, Picard D, Guérin S, Isselin-Nondedeu F, Ribéron A, Trochet A. 2017.** Marking techniques in the Marbled Newt (*Triturus marmoratus*): PIT-Tag and tracking device implant protocols. *Acta Herpetologica* 12:79–88 DOI 10.13128/Acta_Herpetol-19158.
- Legrand D, Trochet A, Moulherat S, Calvez O, Stevens VM, Ducatez S, Clobert J, Baguette M. 2015.** Ranking the ecological causes of dispersal in a butterfly. *Ecography* 38(8):822–831 DOI 10.1111/ecog.01283.
- Léna JP, Clobert J, De Fraipont M, Lecomte J, Guyot G. 1998.** The relative influence of density and kinship on dispersal in the common lizard. *Behavioral Ecology* 9(5):500–507 DOI 10.1093/beheco/9.5.500.
- Lenhardt PP, Brühl CA, Leeb C, Theissinger K. 2017.** Amphibian population genetics in agricultural landscapes: does viticulture drive the population structuring of the European common frog (*Rana temporaria*)? *PeerJ* 5:e3520 DOI 10.7717/peerj.3520.
- Liebgold EB, Brodie ED 3rd, Cabe PR. 2011.** Female philopatry and male-biased dispersal in a direct-developing salamander, *Plethodon cinereus*. *Molecular Ecology* 20(2):249–257 DOI 10.1111/j.1365-294X.2010.04946.x.
- Losos JB. 1990.** Ecomorphology, performance capability, and scaling of West Indian *Anolis* lizards: an evolutionary analysis. *Ecological Monographs* 60(3):369–388 DOI 10.2307/1943062.
- Marsh DM, Milam GS, Gorham N, Beckham NG. 2005.** Forest roads as partial barriers to terrestrial salamander movement. *Conservation Biology* 19:2004–2008 DOI 10.1111/j.1523-1739.2005.00238.x.

- Marsh DM, Trenham PC. 2001.** Metapopulation dynamics and amphibian conservation. *Conservation Biology* 15:40–49 DOI [10.1111/j.1523-1739.2001.00129.x](https://doi.org/10.1111/j.1523-1739.2001.00129.x).
- Matos C, Sillero N, Argaña E. 2012.** Spatial analysis of amphibian road mortality levels in northern Portugal country roads. *Amphibia-Reptilia* 33(3–4):469–483 DOI [10.1163/15685381-00002850](https://doi.org/10.1163/15685381-00002850).
- Mazerolle MJ. 2004.** Amphibian road mortality in response to nightly variations in traffic intensity. *Herpetologica* 60(1):45–53 DOI [10.1655/02-109](https://doi.org/10.1655/02-109).
- Mazerolle MJ, Huot M, Gravel M. 2005.** Behavior of amphibians on the road in response to car traffic. *Herpetologica* 61(4):380–388 DOI [10.1655/04-79.1](https://doi.org/10.1655/04-79.1).
- Moya-Laraño J, Macías-Ordóñez R, Blanckenhorn WU, Fernández-Montraveta C. 2008.** Analysing body condition: mass, volume or density? *Journal of Animal Ecology* 77(6):1099–1108 DOI [10.1111/j.1365-2656.2008.01433.x](https://doi.org/10.1111/j.1365-2656.2008.01433.x).
- Nollért A, Nollért C. 2003.** *Guide des amphibiens d'Europe: biologie, identification, répartition*. Lonay: Delachaux et Niestlé.
- O'Riain M, Jarvis J, Faulkes C. 1996.** A dispersive morph in the naked mole-rat. *Nature* 380(6575):619–621 DOI [10.1038/380619a0](https://doi.org/10.1038/380619a0).
- Ousterhout BH, Semlitsch RD. 2018.** Effects of conditionally expressed phenotypes and environment on amphibian dispersal in nature. *Oikos* 127(8):1142–1151 DOI [10.1111/oik.05276](https://doi.org/10.1111/oik.05276).
- Peig J, Green AJ. 2009.** New perspectives for estimating body condition from mass/length data: the scaled mass index as an alternative method. *Oikos* 118(12):1883–1891 DOI [10.1111/j.1600-0706.2009.17643.x](https://doi.org/10.1111/j.1600-0706.2009.17643.x).
- Peterman WE, Connette GM, Semlitsch RD, Eggert LS. 2014.** Ecological resistance surfaces predict fine scale genetic differentiation in a terrestrial woodland salamander. *Molecular Ecology* 23(10):2402–2413 DOI [10.1111/mec.12747](https://doi.org/10.1111/mec.12747).
- Petronilho JMS, Dias CMM. 2005.** Impact of two forest roads upon wildlife after a road pavement change in a coastal area in the center of Portugal. *Wildlife Biology in Practice* 1:128–139.
- Phillips BL, Brown GP, Webb JK, Shine R. 2006.** Invasion and the evolution of speed in toads. *Nature* 439(7078):803 DOI [10.1038/439803a](https://doi.org/10.1038/439803a).
- Pittman SE, Osbourn MS, Drake DL, Semlitsch RD. 2013.** Predation of juvenile ringed salamanders (*Ambystoma annulatum*) during initial movement out of ponds. *Herpetological Conservation and Biology* 8:681–687.
- Pittman SE, Osbourn MS, Semlitsch RD. 2014.** Movement ecology of amphibians: a missing component for understanding population declines. *Biological Conservation* 169:44–53 DOI [10.1016/j.biocon.2013.10.020](https://doi.org/10.1016/j.biocon.2013.10.020).
- Radinger J, Wolter C. 2014.** Patterns and predictors of fish dispersal in rivers. *Fish and Fisheries* 15(3):456–473 DOI [10.1111/faf.12028](https://doi.org/10.1111/faf.12028).
- R Development Core Team. 2014.** *R: a language and environment for statistical computing*. Vienna: The R Foundation for Statistical Computing. Available at <https://www.R-project.org/>.
- Ronce O. 2007.** How does it feel to be like a rolling stone? Ten questions about dispersal evolution. *Annual Review of Ecology, Evolution, and Systematics* 38(1):231–253 DOI [10.1146/annurev.ecolsys.38.091206.095611](https://doi.org/10.1146/annurev.ecolsys.38.091206.095611).
- Rothermel BB, Luhring TM. 2005.** Burrow availability and desiccation risk of mole salamanders (*Ambystoma talpoideum*) in harvested versus unharvested forest stands. *Journal of Herpetology* 39(4):619–626 DOI [10.1670/251-04A.1](https://doi.org/10.1670/251-04A.1).

- Rothermel BB, Semlitsch RD. 2002. An experimental investigation of landscape resistance of forest versus old-field habitats to emigrating juvenile amphibians. *Conservation Biology* 16(5):1324–1332 DOI 10.1046/j.1523-1739.2002.01085.x.
- Rytwinski T, Fahrig L. 2015. The impacts of roads and traffic on terrestrial animal populations. In: Van Der Ree R, Smith DJ, Grilo C, eds. *Handbook Road Ecology*. Chichester: John Wiley & Sons, Inc., 237–246.
- Šamajová P, Gvoždík L. 2010. Inaccurate or disparate temperature cues? Seasonal acclimation of terrestrial and aquatic locomotor capacity in newts. *Functional Ecology* 24(5):1023–1030 DOI 10.1111/j.1365-2435.2010.01720.x.
- Santos X, Llorente GA, Montori A, Carretero MA, Franch M, Garriga N, Richter-Boix A. 2007. Evaluating factors affecting amphibian mortality on roads: the case of the Common Toad *Bufo bufo*, near a breeding place. *Animal Biodiversity and Conservation* 1:97–104.
- Semlitsch RD. 2008. Differentiating migration and dispersal processes for pond-breeding amphibians. *Journal of Wildlife Management* 72(1):260–267 DOI 10.2193/2007-082.
- Semlitsch RD, Ecrement S, Fuller A, Hammer K, Howard J, Krager C, Mozelyz J, Ogle J, Shipman N, Speier J, Walker M, Walkers B. 2012. Natural and anthropogenic substrates affect movement behavior of the Southern Graycheek Salamander (*Plethodon metcalfi*). *Canadian Journal of Zoology* 90(9):1128–1135 DOI 10.1139/z2012-079.
- Sillero N. 2008. Amphibian mortality levels on Spanish country roads: descriptive and spatial analysis. *Amphibia-Reptilia* 29(3):337–347 DOI 10.1163/156853808785112066.
- Sillero N, Campos J, Bonardi A, Corti C, Creemers R, Crochet PA, Crnobrnja Isailović J, Denoël M, Ficetola GF, Gonçalves J, Kuzmin S, Lymberakis P, De Pous P, Rodriguez A, Sindaco R, Speybroeck J, Toxopeus B, Vieites DR, Vences M. 2014. Updated distribution and biogeography of amphibians and reptiles of Europe. *Amphibia-Reptilia* 35(1):1–31 DOI 10.1163/15685381-00002935.
- Smith MA, Green DM. 2005. Dispersal and the metapopulation paradigm in amphibian ecology and conservation: are all amphibian populations metapopulations? *Ecography* 28(1):110–128 DOI 10.1111/j.0906-7590.2005.04042.x.
- Sotiropoulos K, Eleftherakos K, Tsaparis D, Kasapidis P, Giokas S, Legakis A, Kotoulas G. 2013. Fine scale spatial genetic structure of two syntopic newts across a network of ponds: implications for conservation. *Conservation Genetics* 14(2):385–400 DOI 10.1007/s10592-013-0452-4.
- Stevens VM, Le Boulengé E, Wesselingh RA, Baguette M. 2006. Quantifying functional connectivity: experimental assessment of boundary permeability for the Natterjack toad *Bufo calamita*. *Oecologia* 150(1):161–171 DOI 10.1007/s00442-006-0500-6.
- Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues ASL, Fischman DL, Waller RW. 2004. Status and trends of amphibian declines and extinctions worldwide. *Science* 306(5702):1783–1786 DOI 10.1126/science.1103538.
- Trochet A, Courtois E, Stevens VM, Baguette M, Chaine A, Schmeller DS, Clobert J. 2016b. Evolution of sex-biased dispersal. *Quarterly Review of Biology* 91(3):297–320 DOI 10.1086/688097.
- Trochet A, Dechartre J, Le Chevalier H, Baillat B, Calvez O, Blanchet S, Ribéron A. 2016a. Effects of habitat and fragmented-landscape parameters on amphibian distribution at a large spatial scale. *Herpetological Journal* 26:73.
- Trochet A, Le Chevalier H, Calvez O, Barthe L, Isselin-Nondedeu F, Picard D, Debelgarric M, Pégourié N, Rocher R, Ribéron A. 2017. Post-breeding movement correlated with morphology

in the marbled newt in forest and agricultural lands: a comparative radio-tracking study. *Herpetologica* 73(1):1–9 DOI 10.1655/Herpetologica-D-15-00072.

Van Dyck H, Baguette M. 2005. Dispersal behaviour in fragmented landscapes: routine or special movements? *Basic and Applied Ecology* 6(6):535–545 DOI 10.1016/j.baae.2005.03.005.

Wiens JA. 2001. The landscape context of dispersal. In: Clobert J, Danchin E, Dhondt AA, Nichols JD, eds. *Dispersal*. Oxford: Oxford University Press, 96–109.

Wiens JA, Schooley RL, Weeks RD. 1997. Patchy landscapes and animal movements: do beetles percolate? *Oikos* 78(2):257–264 DOI 10.2307/3546292.

Wilson MC, Chen XY, Corlett RT, Didham RK, Ding P, Holt RD, Holyoak M, Hu G, Hugues AC, Jiang L, Laurance WF, Liu J, Pimm SL, Robinson SK, Russo SE, Si X, Wilcove DS, Wu J, Yu M. 2016. Habitat fragmentation and biodiversity conservation: key findings and future challenges. *Landscape Ecology* 31(2):219–227 DOI 10.1007/s10980-015-0322-1.

Youngquist MB, Inoue K, Berg DJ, Boone MD. 2016. Effects of land use on population presence and genetic structure of an amphibian in an agricultural landscape. *Landscape Ecology* 32(1):147–162 DOI 10.1007/s10980-016-0438-y.