



HAL
open science

Is COVID-19 infection more severe in kidney transplant recipients?

Sophie Caillard, Nathalie Chavarot, Hélène Francois, Marie Matignon, Clarisse Greze, Nassim Kamar, Philippe Gatault, Olivier Thaunat, Tristan Legris, Luc Frimat, et al.

► To cite this version:

Sophie Caillard, Nathalie Chavarot, Hélène Francois, Marie Matignon, Clarisse Greze, et al.. Is COVID-19 infection more severe in kidney transplant recipients?. *American Journal of Transplantation*, 2021, 21 (3), pp.1295-1303. 10.1111/ajt.16424 . hal-03371189

HAL Id: hal-03371189

<https://ut3-toulouseinp.hal.science/hal-03371189v1>

Submitted on 18 Oct 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PROFESSOR SOPHIE CAILLARD (Orcid ID : 0000-0002-0525-4291)
DR NATHALIE CHAVAROT (Orcid ID : 0000-0001-6361-1392)
DR MARIE MATIGNON (Orcid ID : 0000-0001-7704-427X)
DR NASSIM KAMAR (Orcid ID : 0000-0003-1930-8964)
PROFESSOR OLIVIER THAUNAT (Orcid ID : 0000-0002-3648-8963)
DR RENAUD SNANOUDJ (Orcid ID : 0000-0002-6657-6235)
DR DOMINIQUE BERTRAND (Orcid ID : 0000-0002-8766-4859)
DR CHRISTOPHE MASSET (Orcid ID : 0000-0002-7442-2164)
DR LIONEL COUZI (Orcid ID : 0000-0002-9213-6196)
DR JONATHAN MAURICE CHEMOUNY (Orcid ID : 0000-0001-6309-3986)
PROFESSOR ANTOINE DURRBACH (Orcid ID : 0000-0003-0385-8726)
PROFESSOR DANY ANGLICHEAU (Orcid ID : 0000-0001-5793-6174)
DR ILIES BENOTMANE (Orcid ID : 0000-0001-9113-2479)
PROFESSOR YANNICK LE MEUR (Orcid ID : 0000-0001-8913-415X)

Article type : B - Brief Communication

Is Covid-19 infection more severe in kidney transplant recipients?

Sophie Caillard¹ MD PhD, Nathalie Chavarot² MD, H el ene Francois³ MD PhD, Marie Matignon⁴, MD, Clarisse Greze⁵ MD, Nassim Kamar⁶ MD PhD, Philippe Gatault⁷ MD PhD, Olivier Thauat⁸ MD PhD, Tristan Legris⁹ MD, Luc Frimat¹⁰ MD PhD, Pierre Francois Westeel¹¹ MD, Valentin Goutaudier¹² MD, Renaud Snanouj¹³ MD PhD, Charlotte Colosio¹⁴ MD, Antoine Sicard¹⁵ MD PhD, Dominique Bertrand¹⁶ MD, Christiane Mousson¹⁷ MD, PhD, Jamal Bamoulid¹⁸ MD PhD, Christophe Masset¹⁹ MD, Antoine Thierry²⁰, MD PhD, Lionel Couzi²¹ MD PhD, Jonathan M. Chemouny²² MD, Agnes Duveau²³ MD, Valerie Moal⁹ MD PhD, Gilles Blancho¹⁹ MD PhD, Philippe Grimbert⁴ MD PhD, Antoine Durrbach⁴ MD PhD, Bruno Moulin¹ MD PhD, Dany Anglicheau² MD PhD, Yvon Ruch²⁴ MD, Charlotte Kaeuffer²⁴ MD, Ilies Benotmane¹, MD

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/AJT.16424](https://doi.org/10.1111/AJT.16424)

Morgane Solis²⁵PharmD, Yannick Le Meur²⁶ MD PhD, Marc Hazzan^{27*} MD PhD, Francois Danion^{24*} MD – on behalf of the French SOT COVID Registry**

¹Department of Nephrology and Transplantation, Strasbourg University Hospital, INSERM, UMR-S 1109, Strasbourg, France

²Department of Nephrology and Transplantation, Hôpital Universitaire Necker – APHP Center – Université de Paris INEM INSERM U 1151 – CNRS UMR 8253, Paris, France

³AP-HP (Assistance Publique-Hôpitaux de Paris), Department of Nephrology and Transplantation, Hopital Tenon, Paris, France

⁴AP-HP, Nephrology and Renal Transplantation Department, Institut Francilien de Recherche en Néphrologie et Transplantation (IFRNT), Groupe Hospitalier Henri-Mondor/Albert-Chenevier. Université Paris-Est-Créteil, (UPEC), DHU (Département Hospitalo-Universitaire) VIC (Virus-Immunité-Cancer), IMRB (Institut Mondor de Recherche Biomédicale), Equipe 21, INSERM U 955, Créteil, France

⁵Department of Nephrology and Transplantation, Hôpital Bichat, Paris, France

⁶Department of Nephrology and Transplantation, University of Toulouse, Toulouse, France

⁷Department of Nephrology and Transplantation, University of Tours, Tours, France

⁸Department of Transplantation, Nephrology and Clinical Immunology, Hôpital Edouard Herriot, Hospices Civils de Lyon, Université Claude Bernard Lyon 1, Lyon, France

⁹Aix Marseille Université, Hôpitaux Universitaires de Marseille, Hôpital Conception, Centre de Néphrologie et Transplantation Rénale, Marseille, France

¹⁰University of Lorraine, CHRU-Nancy, Department of Nephrology, Vandoeuvre, France
INSERM CIC-EC CIE6, Nancy, France

¹¹Department of Nephrology and Transplantation, University of Amiens, Amiens, France

¹²Department of Nephrology and Transplantation, University of Montpellier, Montpellier, France

¹³Department of Nephrology and Transplantation, Hôpital Bicêtre, Le Kremlin-Bicêtre, France

¹⁴Department of Nephrology and Transplantation, University of Reims, Reims, France

¹⁵Department of Nephrology, Dialysis, and Transplantation, Hopital Pasteur 2, C.H.U. de Nice, Unité de Recherche Clinique Côte d'Azur (UR2CA), Université Côte d'Azur, Nice, France

¹⁶Department of Nephrology and Transplantation, University of Rouen, Rouen, France

¹⁷Department of Nephrology and Transplantation, University of Dijon, Dijon, France

¹⁸Department of Nephrology, University of Besançon, Besançon, France

¹⁹Department of Nephrology and Transplantation, Center Hospitalier Universitaire de Nantes, Nantes, France

²⁰Department of Nephrology, University of Poitiers, Poitiers, France

²¹Department of Nephrology-Transplantation-Dialysis-Apheresis, Hôpital Pellegrin, CHU de Bordeaux Pellegrin. Unité Mixte de Recherche “ImmunoConcEpT” 5164 – Université de Bordeaux, France

²²University of Rennes, CHU Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) UMR_S 1085, CIC – P 1414, Rennes, France

²³Department of Nephrology and Transplantation, University of Angers, Angers, France

²⁴Department of Infectious Diseases, Strasbourg University Hospital, Strasbourg, France

²⁵Department of Virology, Strasbourg University Hospital, INSERM, UMR-S 1109, Strasbourg, France

²⁶Department of Nephrology, CHU de Brest, UMR1227, Lymphocytes B et Autoimmunité, Université de Brest, Inserm, Labex IGO, Brest, France

²⁷Department of Nephrology and Transplantation, University of Lille, Lille, France

*These authors contributed equally to this work

Corresponding author: Sophie Caillard,

Email: Sophie.caillard@chru-strasbourg.fr

ABSTRACT

There are no studies which have compared the risk of severe Covid-19 and related mortality between transplant recipients and non-transplant patients. We enrolled two groups of patients hospitalized for Covid-19, i.e., kidney transplant recipients from the French Registry of Solid Organ Transplant (n=306) and a single-center cohort of non-transplant patients (n=795). An analysis was performed among subgroups matched for age and risk factors for severe Covid-19 or mortality. Severe Covid-19 was defined as admission (or transfer) to an intensive care unit, need for mechanical ventilation, or death. Transplant recipients were younger and had more comorbidities compared to non-transplant patients. They presented with higher creatinine levels and developed more episodes of acute kidney injury. After matching, the 30-day cumulative incidence of severe Covid-19 did not differ between KTR and non-transplant patients; however, 30-day Covid-19-related mortality was significantly higher in KTR (17.9% *versus* 11.4%, respectively, p=0.038). Age >60 years, cardiovascular disease, dyspnea, fever, lymphopenia, and C-reactive protein (CRP) were associated with severe Covid-19 in univariate analysis, whereas transplant status and serum creatinine levels were not. Age >60 years, hypertension, cardiovascular disease, diabetes, CRP >60 mg/L, lymphopenia, kidney transplant status (HR=1.55), and creatinine level >115 $\mu\text{mol/L}$ (HR=2.32) were associated with Covid-19-related mortality in univariate analysis. In multivariable analysis, cardiovascular disease, dyspnea, and fever were associated with severe disease, whereas age >60 years, cardiovascular disease, dyspnea, fever, and creatinine level >115 $\mu\text{mol/L}$ retained their independent associations with mortality. Kidney transplant recipients had a higher Covid-19-related mortality compared to non-transplant hospitalized patients.

Trial Registration: The French SOT COVID Registry (Clinicaltrials.gov identifier, NCT04360707).

Introduction

Prior experience with respiratory viruses in patients who had undergone solid organ transplantation revealed how recipients have greater susceptibility, more rapid progression to pneumonia, greater disease severity, and prolonged viral shedding compared with non-transplant hosts. In light of past coronavirus outbreaks,^{1,2} Covid-19 poses a significant threat for immunocompromised patients, and transplant physicians are particularly concerned about the impact of this new infection on this frail population. Single-center studies have reported a high mortality rate in kidney transplant recipients (KTR) with Covid-19.³⁻⁵ There is also evidence that at least part of Covid-19's severity is linked to the "cytokine storm", which is a disproportionate hyperinflammatory reaction occurring in infected patients.⁶ In this scenario, immunosuppressive drugs may be clinically useful in reducing this dysfunctional immune response by attenuating the positive feedback loop typical of the cytokine release syndrome (CRS). Nonetheless, the question as to whether KTR would actually exhibit a higher risk of severe Covid-19 or – alternatively – immunosuppression would protect them from CRS and critical forms of the disease remains unanswered.

Chronic kidney disease and acute kidney injury (AKI) have been reported to affect the prognosis of patients hospitalized for Covid-19.⁷ Notably, KTR are in an immunosuppressed state with concurrent chronic kidney disease and are particularly susceptible to AKI. Starting from these premises, this research was undertaken to determine how these factors may influence the clinical outcomes of KTR with Covid-19. We also compared the prognosis of Covid-19 in KTR and non-transplant patients by using data from a French nationwide registry.

Patients and Methods

A cohort of KTR hospitalized for Covid-19 was identified from a multicenter nationwide French Registry – termed French SOT COVID – between March 1 and April 30, 2020. Inclusion criteria were age >18 years at the diagnosis of Covid-19 and presence of a functioning graft. The control group consisted of non-transplant adult patients with confirmed Covid-19 who were hospitalized at the Strasbourg University Hospital between March 1 and March 31, 2020. Cases with a history of immunosuppression (previous transplantation; patients on high-dose steroids, immunosuppressive drugs, or biological therapies in the month preceding hospitalization; those with primary immune deficiency; and those with previous splenectomy) were excluded from the control group. The diagnostic criteria for Covid-19 were as follows: 1) severe acute respiratory

syndrome coronavirus-2 (SARS-CoV-2) infection determined by reverse transcriptase-polymerase chain reaction (RT-PCR) testing of nasopharyngeal swab specimens, or 2) presence of typical respiratory symptoms associated with evocative pulmonary lesions on low-dose chest computed tomography (CT) when RT-PCR yielded negative results. AKI was defined according to the Kidney Disease Improving Global Outcomes guidelines. Severe Covid-19 was defined as admission (or transfer) to an intensive care unit (ICU), need for mechanical ventilation, or death. All other patients were considered as non-severe cases. Ethical approval for the creation of the French SOT COVID Registry was obtained from the Institutional Review Board of the Strasbourg University (approval number 02.26). The study was registered at clinicaltrials.gov (NCT04360707). While the requirement for informed consent was waived, all patients were informed about their inclusion in the registry. The study protocol for non-transplant patients was approved by the Human Investigation Review Committee of the Strasbourg University Hospital (approval number CE-2020-51). Patients who declined to participate were deemed ineligible.

Statistical analysis

Discrete variables are presented as counts and percentages, whereas continuous data are summarized as medians and interquartile ranges (IQRs) upon verification of their skewed distribution. The composite endpoint of severe COVID-19 was considered as a time-dependent variable since the onset of the symptoms. Cumulative event curves (for severe Covid-19 or Covid-19-related mortality) of transplant recipients and non-transplant patients were plotted with the Kaplan-Meier method and compared using the log-rank test. Cox proportional hazards univariable and multivariable models were constructed using a backward-conditional selection procedure to identify predictors of the study endpoints. The optimal model was selected according to the highest concordance (Harrel's C statistic) value. Results are expressed as hazard ratios (HRs) with their 95% confidence intervals (CIs). Matching was performed by selection of nearest neighbor best control matches for each individual in the KTR group.⁸ Patients were matched in a 1:1 ratio using the logit of the estimated propensity of being in the transplant group as the distance metric. Age, BMI, cardiovascular and respiratory diseases, cancer, and diabetes were included as covariates in the propensity score model because these variables are the main risk factors for COVID-19.^{9, 10, 11} A caliper (0.3) was set for age only. There were 33 transplant recipients who could not be matched to a non-transplant patient. Tabular data for matched cohorts are reported as standardized mean differences with their 95% CIs. All analyses were undertaken in the R

environment (R Foundation for Statistical Computing, Vienna, Austria). A value of $P < 0.05$ (two-tailed) was considered statistically significant.

Results

A total of 306 KTR were included in the SOT COVID Registry at the time of this analysis. The median recipient age was 62 years (IQR: 52–69 years) and 67.6% were men. The median time between transplantation and Covid-19 diagnosis was 74.6 months (IQR: 27.8–140.6 months), and only 12% of all KTR were in the first post-transplant year at the time of Covid-19 diagnosis. Immunosuppressive drugs used at baseline and management of immunosuppression at the time of Covid-19 diagnosis are summarized in Supplementary Table 1. The control cohort consisted of 795 non-transplant patients with Covid-19 (median age: 69 years, IQR: 57–79 years; 58.6% men). The characteristics of the two study groups are reported in Supplementary Table 2. KTR were younger, more commonly male, and had lower body mass index (BMI) but had more comorbidities (hypertension, cardiovascular diseases, respiratory diseases, and diabetes). They less frequently exhibited dyspnea during admission for Covid-19, but more commonly had fever and diarrhea than non-transplant patients. Of note, the median time from symptom onset to admission was shorter among KTR than non-transplant patients (5 *versus* 7 days, respectively, $p=0.006$). KTR displayed a less severe inflammatory syndrome, a more profound lymphopenia, and a higher creatinine level at admission (176 $\mu\text{mol/L}$ *versus* 75 $\mu\text{mol/L}$, respectively, $p<0.001$). Infection management was slightly different, with antibiotics and azithromycin more frequently used in non-transplant patients, in contrast to more antifungal drugs (4.6% *versus* 2.1%, respectively, $p=0.028$), fewer specific antivirals (lopinavir/ritonavir 5.2% *versus* 21.8%, respectively, $p<0.01$), and more frequent tocilizumab (5.6% *versus* 1%, respectively, $p<0.001$) in KTR. Moreover, KTR were less frequently in need of vasopressor support but were significantly more likely to develop AKI (46.1% *versus* 11.2%, respectively, $p<0.001$) that would require dialysis (12.7% *versus* 8.1%, respectively, $p=0.023$). The 30-day cumulative incidence of severe Covid-19 and death were similar in the two groups (43.8% *versus* 41.2%, $p=0.21$, and 17% *versus* 16.6%, $p=0.46$, in non-transplant patients and KTR, respectively; Supplementary Figures 1A and 1B). Owing to the significant differences in age and comorbidities, a further analysis was performed on matched KTR ($n=273$) and non-transplant patients ($n=273$). The median follow-up time from admission was 64 days (IQR: 55–71 days) for the entire matched cohort. Specifically, it was 58 days (IQR: 48–67 days) and 67 days (IQR: 62–73 days) for transplant recipients and non-

transplant patients, respectively. The characteristics and outcomes of the matched groups are shown in Table 1. The univariable analysis showed that the 30-day cumulative incidence of severe disease was similar in both groups (Figure 1A; 42.2% *versus* 42.1% in KTR and non-transplant patients, respectively; $p=0.6$), whereas the 30-day mortality was significantly higher among KTR (Figure 1B; 17.9% *versus* 11.4%, respectively; $p=0.038$). Risk factors for severe COVID-19 were age > 60 years, cardiovascular disease, dyspnea, fever, lymphopenia, and C-reactive protein >60 mg/L (Table 2). Furthermore, age >60 years, hypertension, cardiovascular disease, diabetes, lymphopenia, being a KTR (HR=1.55, 1.02–2.35), and having a creatinine level >115 $\mu\text{mol/L}$ (HR=2.32, 1.45–3.70) were associated with mortality (Table 3). In multivariable analysis, cardiovascular disease, dyspnea, and fever were independent risk factors for severe disease. Age >60 years, cardiovascular disease, having dyspnea and fever at admission, and a serum creatinine >115 $\mu\text{mol/L}$ were also independently associated with mortality, whereas being a kidney transplant recipient was not (Table 4). Because the two matched groups were not well balanced in terms of hypertension, we constructed a different model in which this variable was included for matching. However, the results of multivariable analysis were entirely consistent with those reported in the model that did not include hypertension as a matching variable (data not shown).

Discussion

The present study compared for the first time hospitalized KTR with Covid-19 to a cohort of hospitalized non-transplant patients in order to determine if they would have different outcomes and a higher mortality rate. First, we demonstrated that the entire cohort of KTR hospitalized for Covid-19 exhibited significant differences compared to the non-transplant cohort. Accordingly, KTR were younger (by 7 years) and had a higher burden of comorbidities. As expected, non-transplant patients had a better renal function at admission. This could reflect either the presence of a preexisting chronic kidney disease or an AKI in KTR – who were frequently admitted with diarrhea and high fever. Moreover, subsequent AKI and renal replacement therapy occurred more frequently among KTR than in non-transplant patients (46.1% and 11.2%, respectively) during hospitalization. AKI was observed in 5.1% of patients hospitalized with Covid-19 in Cheng et al's report ⁷, and 4.5% of patients in the meta-analysis published by Yang et al.¹² The etiology of AKI during Covid-19 is multifactorial. In addition to SARS-CoV-2's direct attack of tubular cells via ACE2 receptors, other factors that may contribute to kidney injury include hypoxia, CRS, and a

hypercoagulable state.¹³ The susceptibility of KTR to dehydration, nephrotoxic drugs, and hemodynamic instability can also explain the high frequency of renal dysfunction in this cohort. Given the differences between the transplant and non-transplant cohorts' baseline characteristics, we performed a matched analysis after adjusting for known risk factors of severe Covid-19 and Covid-19-related death^{9, 10, 11} (i.e., age, BMI, cardiovascular and respiratory diseases, cancer and diabetes) to minimize the effects of potential confounders. Our results validate the findings from previous studies in non-transplant patients with respect to known risk factors for severe Covid-19 and Covid-19-related death.^{14,15} However, being a KTR was not associated with a more frequent need for ICU admission in our study. This could be explained by the shorter time from symptom onset to hospitalization (5 *versus* 7 days in KTR and non-transplant patients, respectively) and the lower incidence of pulmonary involvement at admission (dyspnea: 45.1 *versus* 63.7% in KTR and non-transplant patients, respectively).

However, the comparison of matched cohorts also showed that KTR had a two-fold higher risk of Covid-19-related death compared to non-transplant patients after adjustment for age, BMI, and major comorbidities. While previous studies have shown that transplantation is a risk factor for mortality, a direct matched comparison between transplanted and non-transplant patients had never been performed. Data from a very large cohort of 17 million patients indicated that organ transplant recipients had an adjusted 3.55-fold higher risk of death, whereas those with glomerular filtration rates below 30 mL/min had a 2.5-fold increased risk.¹⁶ In Cheng et al's cohort,⁷ the incidence of in-hospital death in patients with increased baseline serum creatinine was 33.7%, which was higher than in those with normal creatinine levels (13.2%). Notably, this difference persisted after adjusting for age and comorbidities. In prior studies focusing on respiratory viruses (H1N1, SARS, and MERS-CoV),^{17,18} kidney injury was also associated with an increased risk of death. It remains uncertain whether the higher mortality rate observed in KTR is caused by immunosuppression and/or the increased rate of renal dysfunction. Multivariable analysis revealed that being a KTR was not independently associated with mortality, whereas a serum creatinine >115 µmol/L was retained in the model as an independent risk factor for death. These results indicate a prominent role for renal failure as a driver of Covid-19-related mortality.

Our findings need to be interpreted in the context of some limitations. First, one may argue that the comparability between a multicenter French nationwide transplant cohort and a single-center control group is low. Nevertheless, the single-center control group was large and representative of a different range of settings (i.e., medical, surgical and ICU departments). Second, we are aware

that KTR and control patients were managed differently – which can represent a potential source of confounding when their clinical outcomes are analyzed. While baseline lymphocyte count and creatinine concentrations were available for KTR, the majority of non-transplant patients had a negative clinical history before the onset of Covid-19. Therefore, we were unable to provide data on these parameters in non-transplant patients. Finally, all of the study patients were hospitalized. Thus, the question as to whether our findings are generalizable to an outpatient setting remains answered. These caveats notwithstanding, this study is the largest to date to comprehensively compare the clinical features and outcomes of Covid-19 in KTR with respect to non-transplant patients.

In summary, our study shows that, after adjustment for potential confounders, KTR with Covid-19 had a higher mortality rate than non-transplant patients, despite a similar occurrence of severe disease. While preexistent chronic kidney disease or AKI might have a greater prognostic impact than the immunosuppression state, further research is needed to shed more light on this issue.

Acknowledgements

This study was supported by the Strasbourg University Hospital (COVIS-HUS Study- HUS number 7760)

Conflicts of Interest

The authors of this manuscript have no conflicts of interest to declare as described by the *American Journal of Transplantation*.

Data Availability Statement

The data that support the findings of this study are available from the corresponding author upon reasonable request.

****French SOT COVID Registry**

Sophie CAILLARD, Bruno MOULIN, Service de Néphrologie et Transplantation, Hôpitaux Universitaires de Strasbourg, STRASBOURG ; Samira FAFI-KREMER, Laboratoire de Virologie, Hôpitaux Universitaires de Strasbourg, STRASBOURG; Marc HAZZAN, Service de Néphrologie, Hôpital Huriez, LILLE ; ANGLICHEAU Dany, Service de Néphrologie et Transplantation Adultes, AP-HP, Hôpital Necker, PARIS ; Alexandre HERTIG, Jérôme TOURRET, Benoit BARROU, Service de Néphrologie, AP-HP, Hôpital La Pitié Salpêtrière, PARIS ; Emmanuel MORELON, Olivier THAUNAT, Service de Néphrologie, Hôpital Edouard Herriot, LYON ; Lionel COUZI, Pierre MERVILLE, Service de Néphrologie – Transplantation – Dialyse, Hôpital Pellegrin, BORDEAUX ; Valérie MOAL, Tristan LEGRIS, Service de Néphrologie et Transplantation, AP-HM, Hôpital de la Conception, MARSEILLE ; Pierre-François WESTEEL, Maïté JAUREGUY, Service de Néphrologie, CHU Amiens Picardie, AMIENS ; Luc FRIMAT, Service de Néphrologie, CHRU Nancy, VANDOEUVRE ; Didier DUCLOUX, Jamal BAMOULID, Service de Néphrologie, Hôpital Jean-Minjoz, BESANCON; Dominique BERTRAND, Service de Néphrologie, CHU de Rouen, ROUEN ; Michel TSIMARATOS, Florentine GARAIX-GILARDO, Service de Pédiatrie multidisciplinaire, Hôpital La Timone, MARSEILLE ; Jérôme DUMORTIER, Service d'Hépatogastroentérologie, Hôpital Edouard Herriot, LYON ; Sacha MUSSOT, Antoine ROUX, Centre chirurgical Marie Lannelongue, LE PLESSIS ROBINSON ; Laurent SEBBAG, Service d'insuffisance cardiaque, Hôpital Louis Pradel, BRON ; Yannick LE MEUR, Service de Néphrologie, Hôpital de la cavale blanche, BREST ; Gilles BLANCHO, Christophe MASSET, Service de Néphrologie – Transplantation, Hôtel Dieu, NANTES ; Nassim KAMAR, Service de Néphrologie et Transplantation, Hôpital Rangueil, TOULOUSE ; Hélène FRANCOIS, Eric Rondeau, Service de Néphrologie, Dialyse et Transplantation, AP-HP, Hôpital Tenon, PARIS ; Nicolas BOUVIER, Service de Néphrologie, Dialyse, Transplantation rénale, CHU, CAEN ; Christiane MOUSSON, Service de Néphrologie, DIJON ; Matthias BUCHLER, Philippe GATAULT, Service de Néphrologie, TOURS ; Jean-François AUGUSTO, Agnès DUVEAU, Service de Néphrologie, Dialyse, Transplantation, CHU Angers, ANGERS ; Cécile VIGNEAU, Marie-Christine MORIN, Jonathan CHEMOUNY, Leonard GOLBIN, Service de Néphrologie, CHU de Rennes, RENNES; Philippe GRIMBERT, Marie MATIGNON, Antoine DURRBACH, Service de Néphrologie, Hôpital Henri-Mondor, CRETEIL ; Clarisse GREZE, Service de Néphrologie, AP-HP, Hôpital Bichat Claude Bernard, PARIS ; Renaud SNANOUDJ, Service de Néphrologie, Hôpital Foch, ,

Service de Néphrologie et Transplantation Hôpital du Kremlin Bicêtre, LE KREMLIN BICETRE ; Charlotte COLOSIO, Betoul SCHVARTZ, Service de Néphrologie, Hôpital Maison Blanche, REIMS ; Paolo MALVEZZI, Service de Néphrologie, Hémodialyse, Transplantation rénale, Hôpital La Tronche, GRENOBLE ; Christophe MARIAT, Service de Néphrologie, CHU de Saint Etienne, SAINT ETIENNE ; Antoine THIERRY, Service de Néphrologie, Hémodialyse et Transplantation rénale, Hôpital Jean Bernard, POITIERS ; Moglie LE QUINTREC, Service de Néphrologie-Transplantation-Dialyse, CHU Lapeyronie, MONTPELLIER ; Antoine SICARD, Service de Néphrologie, Hôpital Pasteur, NICE ; Jean Philippe REROLLE, Service de Néphrologie, CHU Dupuytren, LIMOGES ; Anne-Élisabeth Heng, Cyril GARROUSTE, Service de Néphrologie, CHU Gabriel Montpied, CLERMONT-FERRAND ; Henri VACHER COPONAT, Service de Néphrologie, CHU de La Réunion, SAINT DENIS ; Éric EPAILLY, service de cardiologie, Hôpitaux Universitaires de Strasbourg, STRASBOURG ; Olivier BRUGIERE, Service d'hépatologie, Hôpital Foch, SURESNES ; Sébastien DHARANCY, Service d'hépatologie, Hôpital Huriez, LILLE ; Éphrem SALAME, service de chirurgie hépatique, Hôpital Universitaire de Tours, TOURS ; Faouzi SALIBA, service d'hépatologie, Centre hépato-biliaire Paul Brousse, VILLEJUIF.

References

1. Kumar D, Tellier R, Draker R, Levy G, Humar A. Severe Acute Respiratory Syndrome (SARS) in a liver transplant recipient and guidelines for donor SARS screening. *Am J Transplant.* 2003;3(8):977-981.
2. AlGhamdi M, Mushtaq F, Awn N, Shalhoub S. MERS CoV infection in two renal transplant recipients: case report. *Am J Transplant.* 2015;15(4):1101-1104.
3. Akalin E, Azzi Y, Bartash R, et al. Covid-19 and kidney transplantation. *N Engl J Med.* 2020;382(25):2475- 2477.
4. Pereira MR, Mohan S, Cohen DJ, et al. Covid-19 in solid transplant organ recipients: initial report of the US epicenter. *Am J Transplant.* 2020;20(7):1800-1808.
5. Alberici F, Delbarba E, Manenti C, et al.
A single center observational study of the clinical characteristics and short-term outcome of 20 kidney transplant patients admitted for SARS-CoV2 pneumonia
Kidney Int. 2020;97(6):1083-1088.

6. Blanco-Melo D, Nilsson-Payant BE, Liu WC, et al. Imbalanced host response to SARS-CoV-2 drives development of COVID-19. *Cell*. 2020;181(5):1036-1045.e9.
7. Cheng Y, Luo R, Wang K et al. Kidney disease is associated with in-hospital death of patients with COVID-19. *Kidney Int*. 2020;97(5):829-838.
8. Ho D, Imai K, King G, Stuart EA. MatchIt: Nonparametric preprocessing for parametric causal inference. *Journal of Statistical Software*. 2011;42(8):1-28.
9. Shi Y, Yu X, Zhao H, Wang H, Zhao R, Sheng J. Host susceptibility to severe COVID-19 and establishment of a host risk score: findings of 487 cases outside Wuhan. *Crit Care*. 2020 Mar 18;24(1):108.
10. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020 Mar 28;395(10229):1054-1062.
11. Harrison SL, Fazio-Eynullayeva E, Lane DA, Underhill P, Lip GYH. Comorbidities associated with mortality in 31,461 adults with COVID-19 in the United States: A federated electronic medical record analysis. *PLoS Med*. 2020 Sep 10;17(9):e1003321.
12. Yang X, Jin Y, Li R, Zhang Z, Sun R, Chen D. Prevalence and impact of acute renal impairment on COVID-19: a systematic review and meta-analysis. *Critical Care*. 2020;24(1):356.
13. Naicker S, Yang CW, Hwang SJ, Liu BC, Chen JH, Jha V. The novel coronavirus 2019 epidemic and kidneys. *Kidney Int*. 2020;97(5):824-828.
14. Deng G, Yin M, Chen X, Zeng F. Clinical determinants for fatality of 44,672 patients with COVID-19. *Critical Care* 2020;24(1):179.
15. Grasselli G, Zangrillo A, Zanella A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy Region, Italy. *JAMA*. 2020;323(16):1574-1581.
16. Williamson EJ, Walker AJ, Bhaskaran K. OpenSAFELY: factors associated with COVID-19 death in 17 million patients [published online 8 July, 2020]. *Nature*. Doi: doi.org/10.1038/s41586-020-2521-4.
17. Chu KH, Tsang WK, Tang CS, et al. Acute renal impairment in coronavirus-associated severe acute respiratory syndrome. *Kidney Int*. 2005;67(2):698-705.

18. Jung JY, Park BH, Hong SB, et al. Acute kidney injury in critically ill patients with pandemic influenza A pneumonia 2009 in Korea: a multicenter study. *J Crit Care.* 2011;26(6):577-585.

Table 1. Clinical characteristics, management, and outcomes of matched non-transplant patients and kidney transplant recipients hospitalized for Covid-19

	Non-Transplant <i>N</i> =273	Transplant <i>N</i> =273	SMD* [95%CI]	N
Baseline characteristics				
Median age [IQR], years	63.0 [48.0-74.0]	62.0 [53.0-69.0]	0.042 [-0.126;0.210]	546
Age > 60 years, n (%)	147 (53.8%)	159 (58.2%)	0.089 [-0.079;0.256]	546
Men, n (%)	173 (63.4%)	181 (66.3%)	0.061 [-0.106;0.229]	546
Median BMI [IQR], kg/m ²	27.0 [23.0-30.0]	26.0 [24.0-30.0]	0.024 [-0.144;0.192]	546
BMI > 25 kg/m ² , n (%)	181 (66.3%)	177 (64.8%)	0.031 [-0.137;0.199]	546
Hypertension, n (%)	136 (49.8%)	232 (91.3%)	1.024 [0.842;1.205]	527
RAS blockers, n (%)	94 (34.4%)	121 (48.8%)	0.294 [0.122;0.467]	521
Cardiovascular disease, n (%)	106 (38.8%)	106 (38.8%)	0.000 [-0.168;0.168]	546
Respiratory disease, n (%)	45 (16.5%)	38 (13.9%)	0.071 [-0.096;0.239]	546
Diabetes, n (%)	98 (35.9%)	101 (37.0%)	0.023 [-0.145;0.191]	546
Cancer, n (%)	26 (9.5%)	34 (12.5%)	0.094 [-0.074;0.262]	546
Smoking, n (%)	12 (4.4%)	28 (12.7%)	0.301 [0.123;0.480]	493
Clinical presentation				
Anosmia, n (%)	24 (11.5%)	34 (14.1%)	0.077 [-0.109;0.262]	449
Cough, n (%)	158 (57.9%)	162 (66.1%)	0.171 [-0.002;0.343]	518
Dyspnea, n (%)	174 (63.7%)	123 (45.1%)	0.382 [0.213;0.551]	546
Fever, n (%)	201 (73.6%)	199 (80.6%)	0.166 [-0.007;0.338]	520
Headache, n (%)	47 (17.2%)	41 (18.3%)	0.028 [-0.148;0.205]	497
Diarrhea, n (%)	63 (23.1%)	93 (36.3%)	0.293 [0.122;0.465]	529
Time from diagnosis to admission [IQR] , days	7.0 [3.0-9.0]	5.0 [3.0-8.0]	0.088 [-0.086;0.262]	511
Laboratory data				
Median CRP [IQR], mg/L	80 [33-148]	62 [27-118]	0.143 [-0.044;0.330]	451
Median lymphocyte count [IQR], G/L	0.88 [0.65-1.29]	0.70 [0.41-0.96]	0.382 [0.193;0.571]	450

	Non-Transplant N=273	Transplant N=273	SMD* [95%CI]	N
Median platelet count [IQR], G/L	200 [159-268]	180 [146-238]	0.173 [-0.014;0.360]	453
Thrombopenia < 150 G/L, n (%)	57 (22%)	53 (27%)	0.129 [-0.058;0.315]	453
Median creatinine [IQR], μmol/L	76 [59-99]	176 [132-259]	0.945 [0.759;1.131]	495
Drug treatment				
Azithromycin, n (%)	123 (45.1%)	66 (24.2%)	0.450 [0.280;0.620]	546
Other antibiotics, n (%)	204 (74.7%)	179 (65.6%)	0.201 [0.033;0.369]	546
Antifungal drugs, n (%)	7 (2.6%)	12 (4.4%)	0.100 [-0.068;0.268]	546
Remdesivir, n (%)	0 (0.0%)	2 (0.7%)	0.121 [-0.046;0.289]	546
Lopinavir/Ritonavir, n (%)	71 (26.0%)	15 (5.5%)	0.587 [0.416;0.758]	546
Oseltamivir, n (%)	2 (0.7%)	6 (2.2%)	0.122 [-0.046;0.290]	546
Hydroxychloroquine, n (%)	55 (20.1%)	63 (23.1%)	0.071 [-0.097;0.239]	546
Tocilizumab, n (%)	3 (1.1%)	15 (5.5%)	0.248 [0.080;0.416]	546
Outcomes				
Bacterial coinfection, n (%)	169 (61.9%)	54 (19.8%)	0.948 [0.772;1.135]	546
Viral coinfection, n (%)	1 (0.4%)	9 (3.3%)	0.220 [0.052;0.388]	546
Fungal coinfection, n (%)	2 (0.7%)	11 (4.0%)	0.218 [0.029;0.386]	546
Oxygen therapy, n (%)	156 (71.2%)	170 (73.9%)	0.060 [-0.125;0.245]	449
Mechanical ventilation, n (%)	96 (35.2%)	78 (28.6%)	0.142 [-0.026;0.310]	546
Vasopressor support, n (%)	69 (25.3%)	36 (13.2%)	0.310 [0.142;0.479]	546
Acute kidney injury, n (%)	36 (13.2%)	125 (45.8%)	0.766 [0.592;0.939]	546
Renal replacement therapy, n (%)	27 (9.9%)	36 (13.2%)	0.103 [-0.065;0.271]	546

Abbreviations: IQR, interquartile range; BMI, body mass index; ICU, intensive care unit; RAS, renin-angiotensin system; CRP, C-reactive protein.

*SMD: standardized mean difference, CI, confidence interval.

Data are expressed as medians (IQRs) or counts (percentages), as appropriate.

Patients were matched (1:1 ratio) for age, BMI, cardiovascular and respiratory diseases, cancer, and diabetes.

Accepted Article

Table 2. Risk factors for severe Covid-19 (ICU admission or mechanical ventilation or death) in univariable analysis in matched transplant and non-transplant cohorts (n=546)

	No event <i>N=314</i>	Event <i>N=232</i>	HR	p.ratio	N
Baseline characteristics					
Renal Transplantation - no.(%)	157 (50.0%)	116 (50.0%)	0.91 [0.71;1.18]	0.498	546
Median age [IQR] - yr	60.0 [48.0-70.0]	65.0 [55.8-73.0]	1.01 [1.01;1.02]	0.002	546
Age > 60 yr - no.(%)	159 (50.6%)	147 (63.4%)	1.46 [1.12;1.91]	0.005	546
Male - no.(%)	194 (61.8%)	160 (69.0%)	1.30 [0.99;1.72]	0.063	546
BMI > 25 kg/m ² - no.(%)	200 (63.7%)	158 (68.1%)	1.20 [0.91;1.59]	0.187	546
Hypertension - no.(%)	207 (68.5%)	161 (71.6%)	1.05 [0.78;1.40]	0.763	527
RAS blockers - no.(%)	126 (42.0%)	89 (40.3%)	0.92 [0.70;1.20]	0.534	521
Cardiovasc. disease - no.(%)	109 (34.7%)	103 (44.4%)	1.35 [1.05;1.76]	0.022	546
Respiratory disease - no.(%)	47 (15.0%)	36 (15.5%)	1.04 [0.73;1.49]	0.819	546
Diabetes - no.(%)	102 (32.5%)	97 (41.8%)	1.29 [1.00;1.68]	0.055	546
Cancer - no.(%)	37 (11.8%)	23 (9.91%)	0.86 [0.56;1.32]	0.489	546
Smoking - no.(%)	25 (8.80%)	15 (7.18%)	0.79 [0.46;1.33]	0.370	493
Admission characteristics					
Anosmia - no.(%)	40 (15.2%)	18 (9.68%)	0.66 [0.40;1.07]	0.090	449
Cough - no.(%)	196 (66.2%)	124 (55.9%)	0.74 [0.57;0.96]	0.025	518
Dyspnea - no.(%)	152 (48.4%)	145 (62.5%)	1.71 [1.31;2.23]	<0.001	546
Fever - no.(%)	216 (72.2%)	184 (83.3%)	1.61 [1.13;2.29]	0.009	520
Headache - no.(%)	57 (20.1%)	31 (14.6%)	0.73 [0.50;1.07]	0.109	497
Diarrhea - no.(%)	90 (29.4%)	66 (29.6%)	1.00 [0.75;1.33]	0.995	529
Time from diag. to admission [IQR] - d	6.00 [3.00-9.00]	6.00 [3.00-8.00]	0.99 [0.96;1.02]	0.473	511
Biological data					
CRP > 60 mg/l - no.(%)	119 (45.6%)	135 (71.1%)	2.54 [1.86;3.48]	<0.001	451
Median lymphocyte count [IQR] - G/l	0.85 [0.58-1.21]	0.75 [0.50-1.03]	0.74 [0.57;0.98]	0.035	450

	No event <i>N=314</i>	Event <i>N=232</i>	HR	p.ratio	N
Median platelet count [IQR] - G/l	189 [148-257]	197 [153-257]	1.00 [1.00;1.00]	0.548	453
Thrombopenia < 150 G/l - no.(%)	65 (25.3%)	45 (23.0%)	0.89 [0.64;1.24]	0.502	453
Median SCr [IQR] - μmol/l	102 [66.7-176]	127 [77.0-204]	1.00 [1.00;1.00]	0.373	495
SCr > 115 μmol/l	129 (46.7%)	118 (53.9%)	1.15 [0.88;1.50]	0.292	495

Patients were matched (1:1 ratio) for age, BMI, cardiovascular and respiratory diseases, cancer, and diabetes.

Abbreviations: IQR, interquartile range; BMI, body mass index; RAS, renin-angiotensin system; cardiovasc, cardiovascular; CRP, C-reactive protein; SCr, serum creatinine; diag, diagnosis; HR, hazard ratio.

Data are expressed as medians (IQRs) or counts (percentages), as appropriate.

Accepted Article

Table 3. Risk factors for death in univariable analysis in the matched transplant and non-transplant cohorts (n=546)

	No event N=454	Event N=92	HR	p.ratio	N
Baseline characteristics					
Renal Transplantation - no.(%)	218 (48.0%)	55 (59.8%)	1.55 [1.02;2.35]	0.039	546
Median age [IQR] - yr	60.0 [50.0-69.0]	71.0 [62.0-79.2]	1.05 [1.04;1.07]	<0.001	546
Age > 60 yr - no.(%)	232 (51.1%)	74 (80.4%)	3.61 [2.16;6.05]	<0.001	546
Male - no.(%)	295 (65.0%)	59 (64.1%)	0.97 [0.63;1.49]	0.891	546
BMI > 25 kg/m ² - no.(%)	298 (65.6%)	60 (65.2%)	1.00 [0.65;1.53]	0.994	546
Hypertension - no.(%)	298 (67.9%)	70 (79.5%)	1.76 [1.05;2.95]	0.033	527
RAS blockers - no.(%)	177 (40.6%)	38 (44.7%)	1.16 [0.76;1.78]	0.492	521
Cardiovasc disease - no.(%)	160 (35.2%)	52 (56.5%)	2.23 [1.47;3.36]	<0.001	546
Resp. disease - no.(%)	70 (15.4%)	13 (14.1%)	0.91 [0.51;1.64]	0.760	546
Diabetes - no.(%)	154 (33.9%)	45 (48.9%)	1.75 [1.16;2.64]	0.007	546
Cancer - no.(%)	51 (11.2%)	9 (9.78%)	0.86 [0.43;1.71]	0.668	546
Smoking - no.(%)	32 (7.75%)	8 (10.0%)	1.27 [0.61;2.64]	0.519	493
Admission characteristics					
Anosmia - no.(%)	55 (14.2%)	3 (4.84%)	0.33 [0.10;1.05]	0.060	449
Cough - no.(%)	274 (63.3%)	46 (54.1%)	0.71 [0.46;1.09]	0.118	518
Dyspnea - no.(%)	240 (52.9%)	57 (62.0%)	1.45 [0.95;2.20]	0.086	546
Fever - no.(%)	330 (75.9%)	70 (82.4%)	1.47 [0.84;2.56]	0.179	520
Headache - no.(%)	76 (18.2%)	12 (15.0%)	0.82 [0.44;1.51]	0.524	497
Diarrhea - no.(%)	135 (30.5%)	21 (24.4%)	0.76 [0.46;1.24]	0.275	529
Time from diag. to admission [IQR] - d	6.00 [3.00-9.00]	4.00 [2.00-7.00]	0.94 [0.89;0.99]	0.018	511
Biological data					
CRP > 60 mg/l - no.(%)	207 (54.0%)	47 (69.1%)	1.84 [1.10;3.08]	0.020	451
Median lymphocyte count [IQR] - G/l	0.80 [0.57-1.18]	0.71 [0.45-0.96]	0.43 [0.24;0.77]	0.005	450
Median platelet count [IQR] - G/l	190 [150-253]	201 [153-259]	1.00 [1.00;1.00]	0.526	453

	No event N=454	Event N=92	HR	p.ratio	N
Thrombopenia < 150 G/l - no.(%)	93 (24.7%)	17 (22.4%)	0.87 [0.51;1.49]	0.604	453
Median SCr [IQR] - $\mu\text{mol/l}$	102 [69.7-179]	151 [100-219]	1.00 [1.00;1.00]	0.029	495
SCr > 115 $\mu\text{mol/l}$	192 (46.4%)	55 (67.9%)	2.32 [1.45;3.70]	<0.001	495

Patients were matched (1:1 ratio) for age, BMI, cardiovascular and respiratory diseases, cancer, and diabetes.

Abbreviations: IQR, interquartile range; BMI, body mass index; RAS, renin-angiotensin system; cardiovasc, cardiovascular; CRP, C-reactive protein; SCr, serum creatinine; diag, diagnosis; HR, hazard ratio.

Data are expressed as medians (IQRs) or counts (percentages), as appropriate.

Table 4. Multivariable analysis of risk factors for severe disease^a and mortality in matched transplant and non-transplant cohorts (n=546)

A. Severe disease	HR	p	B. Mortality	HR	p
Cardiovasc disease	1.35 [1.03;1.76]	0.028	Cardiovasc disease	1.54 [0.96;2.46]	0.071
Cough	0.61 [0.46;0.80]	<0.001	Cough	0.58 [0.36;0.92]	0.022
Dyspnea	1.90 [1.43;2.53]	0.004	Dyspnea	1.74 [1.08;2.78]	0.022
Fever	1.70 [1.19;1.76]	0.004	Fever	1.81 [1.00;3.28]	0.050
			SCr > 115 $\mu\text{mol/L}$	2.40 [1.48;3.87]	<0.001
			Age > 60 years	3.47 [1.86; 6.47]	<0.001

Abbreviations: SCr, serum creatinine; HR, hazard ratio; cardiovasc, cardiovascular

Concordance for the severe disease model: 0.63; concordance for the mortality model: 0.73.

^a Severe disease defined as requirement for ICU admission or mechanical ventilation, or death

Patients were matched (1:1 ratio) for age, BMI, cardiovascular and respiratory diseases, cancer, and diabetes.

Missing data:

- Severe model: 34 observations

- Death model: 64 observations

Figure legend

Figure 1. Thirty-day cumulative incidence of the composite endpoint (ICU admission or death) and death only in the non-transplant patients (solid line) and kidney transplant recipients (dashed line) matched cohorts

A) Composite endpoint (non-transplant patients: 42.1%, kidney transplant recipients: 42.2%). B) Death only (non-transplant patients: 11.4%, kidney transplant recipients: 17.9%)

Figure 1

