



**HAL**  
open science

## Misdiagnosis of imported falciparum malaria from African areas due to an increased prevalence of pfhrp2/pfhrp3 gene deletion

Xavier Iriart, Sandie Menard, Pamela Chauvin, Hasna S Mohamed, Elena Charpentier, Mohamed A Mohamed, Antoine Berry, Mohamed H Aboubaker

► **To cite this version:**

Xavier Iriart, Sandie Menard, Pamela Chauvin, Hasna S Mohamed, Elena Charpentier, et al.. Misdiagnosis of imported falciparum malaria from African areas due to an increased prevalence of pfhrp2/pfhrp3 gene deletion: the Djibouti case. *Emerging microbes & infections*, 2020, 9 (1), pp.1984-1987. 10.1080/22221751.2020.1815590 . hal-03526514

**HAL Id: hal-03526514**

**<https://ut3-toulouseinp.hal.science/hal-03526514>**

Submitted on 14 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Misdiagnosis of imported *falciparum* malaria from African areas due to an increased prevalence of *pfhrp2/pfhrp3* gene deletion: the Djibouti case

Xavier Iriart<sup>a,b\*</sup>, Sandie Menard<sup>b\*</sup>, Pamela Chauvin<sup>b</sup>, Hasna S. Mohamed<sup>c</sup>, Elena Charpentier<sup>a,b</sup>, Mohamed A. Mohamed<sup>c</sup>, Antoine Berry<sup>a,b†</sup> and Mohamed H. Aboubaker<sup>d†</sup>

<sup>a</sup>Département de Parasitologie-Mycologie, Centre Hospitalier Universitaire de Toulouse, Toulouse, France; <sup>b</sup>Centre de Physiopathologie de Toulouse Purpan (CTPP), INSERM, CNRS, Université de Toulouse III, UPS, Toulouse, France; <sup>c</sup>Laboratoire de l'Hôpital Général Peltier, Djibouti, République de Djibouti; <sup>d</sup>Laboratoire d'analyse médicale Mer Rouge, Djibouti, République de Djibouti

### ABSTRACT

Following the diagnosis of a *falciparum* malaria case imported from Djibouti and not detected by a pfHRP2-based rapid diagnostic test (RDT), we investigated the prevalence of the *pfhrp2/pfhrp3*-deleted parasites in Djibouti using 378 blood samples collected between January and May 2019, from Djiboutian patients with suspected malaria. Malaria diagnosis by quantitative PCR confirmed the presence of *Plasmodium falciparum* for 20.9% (79/378) samples while RDTs did not detect HRP2 antigen in 83.5% (66/79) of these samples. Quantitative PCRs targeting the *pfhrp2/pfhrp3* genes confirmed the absence of both genes for 86.5% of *P. falciparum* strains. The very large number (86.5%) of *falciparum* parasites lacking the *pfhrp2/pfhrp3* genes observed in this study, now justifies the use of non-HRP2 alternative RDTs in Djibouti. In this area and in most countries where HRP2-based RDTs constitute the main arsenal for *falciparum* malaria diagnosis, it is important to implement a systematic surveillance and to inform biologists and clinicians about the risk of malaria misdiagnosis. Further investigations are needed to better understand the mechanism of selection and diffusion of the *pfhrp2/pfhrp3*-deleted parasites.

**ARTICLE HISTORY** Received 26 May 2020; Revised 3 August 2020; Accepted 23 August 2020

**KEYWORDS** *Plasmodium falciparum*; histidine-rich protein 2; deletion; Djibouti; rapid diagnostic test

A case of imported *falciparum* malaria acquired in Djibouti was diagnosed in 2019 in the Toulouse University Hospital (France) by microscopy and a positive *P. falciparum* specific real-time PCR (qPCR) [1]. An interesting feature of the *falciparum* strain from this patient was its inability to positivize the HRP2 band of the Rapid Diagnostic Test (RDT) (Palutop+4 Optima, Biosynex) despite a significant parasitaemia (1.3%). A *pfhrp2/pfhrp3* deletion was confirmed for this strain (data not shown).

In order to broadly evaluate the prevalence of *pfhrp2/pfhrp3*-deleted parasites in Djibouti, a study was carried out between January and May 2019. The flow chart for this study is described in Figure 1. Three hundred and seventy-eight blood samples were collected from malaria-suspected patients in 3 close sites located in Djibouti city (the Mer Rouge Medical Analysis Laboratory, the National Social Security Funds Laboratory and the Peltier Hospital Laboratory) and *Plasmodium* diagnosis was assessed with HRP2-based RDT (Palutop +4 Optima, Biosynex). Twenty-


two percent (83/378) of blood samples were tested pan-pLDH positive. After extraction of DNA from the 83 pan-pLDH(+) blood samples using a DNeasy Blood & Tissue Kit® (Qiagen®), qPCR, as previously used [1], confirmed the presence of *P. falciparum* alone for 90.4% (75/83) samples, both *P. falciparum* and *P. vivax* for 4.8% (4/83) samples and *P. vivax* only for 4.8% (4/83) samples. Among the 79 *P. falciparum* PCR(+) samples, the RDT did not detect the *P. falciparum* HRP2 antigen for 83.5% (66/79) of samples (Table 1). Before assessing the *pfhrp2/pfhrp3* gene deletion, we performed a qPCR targeting the single copy *P. falciparum*-specific  $\beta$ -tubulin gene (PF3D7\_1008700) [2]. This qPCR is less sensitive than the *pfhrp2/pfhrp3* ones with a limit of detection for  $\beta$ -tubulin qPCR of 15 parasites per  $\mu$ l compared to 5 and 1.67 parasites per  $\mu$ l for *pfhrp2* and *pfhrp3* qPCRs, respectively. The  $\beta$ -tubulin qPCR made it possible to rule out 5  $\beta$ -tubulin PCR(-) samples that would have been false negative in *pfhrp2/pfhrp3* qPCR due to a low quantity or quality of specific DNA. Two

**CONTACT** Xavier Iriart  iriart.x@chu-toulouse.fr and Antoine Berry  berry.a@chu-toulouse.fr,  Service de Parasitologie Mycologie, Centre Hospitalier Universitaire de Toulouse, Hôpital Purpan, Institut Fédératif de biologie (IFB), 330 avenue de Grande Bretagne, TSA 40031, 31059 Toulouse Cedex 9, Toulouse, France

\*Equal contribution as first author

†Equal contribution as last author

© 2020 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group, on behalf of Shanghai Shangyixun Cultural Communication Co., Ltd This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



**Figure 1.** Flow chart of samples throughout the study. \*Probable infection with *falciparum pfhrp2/pfhrp3* (-) strain with residual circulating HRP2 antigen from an old *falciparum pfhrp2/pfhrp3* (+) infection.

**Table 1.** Characteristics of *Plasmodium*-infected patients.

	N	%
<i>Plasmodium</i> identification by qPCR		
<i>Plasmodium</i> sp. Positive	83	
<i>P. falciparum</i> positive	79/83	95.2%
RDT results on <i>P. falciparum</i> strain		
pan-pLDH positive	79	
HRP2 negative	66/79	83.5%
<i>Pfhrp2/pfhrp3</i> genotyping on <i>P. falciparum</i> strain		
$\beta$ -tubulin positive	74/79	93.7%
<i>Pfhrp2/pfhrp3</i> negative	64/74	86.5%

modified qPCRs were carried out in order to amplify the exon 2 of *pfhrp2* and *pfhrp3* genes [3]. The amplification of *pfhrp2* and *pfhrp3* was performed in a final reaction volume of 10  $\mu$ L containing the DNA template, with 0.15  $\mu$ M of each primer (*pfhrp2-F1* and *pfhrp2-R1* for *pfhrp2* gene and *pfhrp3-F1* and *pfhrp3-R1* for *pfhrp3* gene), 5 mM or 4 mM MgCl<sub>2</sub>, respectively and LightCycler® FastStart DNA Master SYBR Green I (Roche Diagnostics GmbH). The qPCR conditions were identical for both genes with a

denaturation at 95°C for 15 min, followed by 45 cycles of 95°C for 15 s, 55°C for 20 s and 72°C for 60 s. The *pfhrp2/pfhrp3* qPCRs confirmed the absence of both genes for 86.5% (64/74) of the *P. falciparum* strains (Table 1). One hundred percent of the HRP2 RDT(-) samples were deleted for *pfhrp2/pfhrp3* genes.

## Discussion

For more than a decade, rapid diagnostic tests (RDTs) have become essential to malaria diagnosis. A total of 276 million RDTs were sold globally in 2017, and in sub-Saharan Africa, RDTs accounted for an estimated 75% of all the diagnosis tests carried out for suspected cases [4]. RDTs can detect malaria-specific antigens: Aldolase or lactate dehydrogenase enzymes (pLDH) common to all *Plasmodium* species, LDH specific to a particular one (i.e. pvLDH specific for *P. vivax*) and histidine-rich protein 2 (pfHRP2) only produced by *P. falciparum* [5]. Due to the dominance and the severity of *falciparum* infection, 90% of the RDTs used in the diagnosis of malaria were only based on pfHRP2 detection [4]. PfHRP2-based RDTs frequently cross-react with a second *P. falciparum* protein, the PfHRP3 antigen due to a high degree of similarity in the amino acid sequence [6].

However, since 2010, false negative results with pfHRP2-based RDTs have been reported in the Amazon region and in some parts of Africa and Asia [7–9], usually due to the deletion of *pfhrp2* and/or *pfhrp3*. These *pfhrp2/pfhrp3*-deleted *falciparum* strains raise a diagnosis issue as a large number of *falciparum*-infected patients will go undetected by the pfHRP2-based RDTs. The public health consequence can be an increased risk of malaria morbidity and mortality due to delayed treatments, and an increase in the transmission of malaria.

Our study identified a very high prevalence (86.5%) of parasites lacking *pfhrp2/pfhrp3* genes among the *falciparum* strains circulating in Djibouti. The lack of previous data makes it difficult to date this trend. However, the current warning of health professionals from Djibouti, who are concerned by a high level of negative RDTs despite clinical suspicions of malaria, suggests that this trend is recent.

Djibouti is a major transit point for migrants, predominantly Afar and Somalis and lesser numbers Ethiopians and Eritreans. These human migratory and commercial flows could be at the origin of the diffusion of parasites lacking *pfhrp2/pfhrp3* genes as these strains are already present in these areas at a high frequency (4.8% and 62% in Ethiopia (2017) and Eritrea (2016), respectively) [10,11]. This data would be in favour of the importation of these clones from neighbouring countries rather than a local emergence, but this question can solely be answered through genotyping.

The second main issue is to understand how these *pfhrp2/pfhrp3*-deleted parasites are selected. The consensual assumption is the misdiagnosis and the

selection of *pfhrp2/pfhrp3*-deleted strains by the exclusive use of HRP2-based tests over time, as previously observed in Eritrea [11] and predicted by mathematical modelling [12]. So, the delay or the absence of treatment against these *pfhrp2/pfhrp3*-deleted parasites due to misdiagnosis would have given them a selective advantage compared to *pfhrp2/pfhrp3* non-deleted strains. In Djibouti, malaria diagnosis is exclusively based on the use of RDT selective for the detection of *P. falciparum* and *P. vivax* but not for the *Plasmodium* species with pan-pLDH [13]. Furthermore, the low genetic diversity linked to the low transmission rate of malaria in Djibouti could have reinforced this trend [13]. Another possibility is that a *pfhrp2/pfhrp3*-deleted clone could have been indirectly selected due to the presence, on this same clone, of another mutation responsible for the resistance to an antimalarial used in this area. This scenario is unlikely in Djibouti since none of the 7 *pfhrp2/pfhrp3*-deleted strains tested in this study carried a mutation on the *PfKelch13* protein (data not shown) [14].

A systematic country-wide surveillance might ensure that the prevalence found in this study reflects the national one. However, the high prevalence of *pfhrp2/pfhrp3*-deleted parasites observed in Djibouti now justifies the use of non-HRP2 alternative RDTs like LDH- or aldolase-based RDTs. Indeed, the WHO recommends a policy switch to more effective WHO-approved non-HRP2 malaria tests when the prevalence of *pfhrp2*-deleted parasites meets or exceeds a 5% prevalence. Despite the costs and resources involved [15], these investments are required to control malaria morbidity and mortality in Djibouti.

In most countries, HRP2-based RDTs constitute the main diagnosis arsenal for *falciparum* imported malaria [16,17]. Consequently, a negative HRP2-based RDT should therefore not rule out a *falciparum* infection. The high frequency of mutant parasites in Djibouti raises the question on the risk of malaria misdiagnosis for locals but also for travellers or soldiers as Djibouti's strategic situation has made it a desirable location for foreign military bases.

## Acknowledgements

We would like to thank all the technical teams from the Mer Rouge Medical Analysis Laboratory, the National Social Security Funds Laboratory and the Peltier Hospital Laboratory for the sample collection.

We would like to thank all the molecular biology technicians as well as the biology students of the Parasitology-Mycology Department of the Toulouse University Hospital involved in malaria diagnosis.

## Disclosure statement

No potential conflict of interest was reported by the author(s).

## References

- [1] Fabre R, Berry A, Morassin B, et al. Comparative assessment of conventional PCR with multiplex real-time PCR using SYBR Green I detection for the molecular diagnosis of imported malaria. *Parasitology*. 2004;128:15–21.
- [2] Witkowski B, Duru V, Khim N, et al. A surrogate marker of piperazine-resistant *Plasmodium falciparum* malaria: a phenotype-genotype association study. *Lancet Infect Dis*. 2017;17:174–183.
- [3] Baker J, McCarthy J, Gatton M, et al. Genetic diversity of *Plasmodium falciparum* histidine-rich protein 2 (PfHRP2) and its effect on the performance of PfHRP2-based rapid diagnostic tests. *J Infect Dis*. 2005;192:870–877.
- [4] WHO | World malaria report 2018 [Internet]. WHO [cited 2020 Mar 26]. Available from: <http://www.who.int/malaria/publications/world-malaria-report-2018/report/en/>
- [5] WHO | Malaria diagnosis - new perspectives [Internet]. WHO [cited 2020 Mar 26]. Available from: <https://www.who.int/tdr/publications/tdr-research-publications/malaria-diagnosis/en/>
- [6] Baker J, Ho M-F, Pelecanos A, et al. Global sequence variation in the histidine-rich proteins 2 and 3 of *Plasmodium falciparum*: implications for the performance of malaria rapid diagnostic tests. *Malar J*. 2010;9:129.
- [7] Cheng Q, Gatton ML, Barnwell J, et al. *Plasmodium falciparum* parasites lacking histidine-rich protein 2 and 3: a review and recommendations for accurate reporting. *Malar J*. 2014;13:283.
- [8] Beshir KB, Sepúlveda N, Bharmal J, et al. *Plasmodium falciparum* parasites with histidine-rich protein 2 (*pfhrp2*) and *pfhrp3* gene deletions in two endemic regions of Kenya. *Sci Rep*. [Internet] 2017 [cited 2020 Jul 30];7. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5677122/>.
- [9] Grignard L, Nolder D, Sepúlveda N, et al. A novel multiplex qPCR assay for detection of *Plasmodium falciparum* with histidine-rich protein 2 and 3 (*pfhrp2* and *pfhrp3*) deletions in polyclonal infections. *EBioMedicine* [Internet] 2020 [cited 2020 Jul 23];55. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7218259/>.
- [10] Girma S, Cheaveau J, Mohon AN, et al. Prevalence and epidemiological characteristics of asymptomatic malaria based on ultrasensitive diagnostics: a cross-sectional study. *Clin Infect Dis Off Publ Infect Dis Soc Am*. 2019;69:1003–1010.
- [11] Berhane A, Anderson K, Mihreteab S, et al. Major threat to malaria control programs by *Plasmodium falciparum* lacking histidine-rich protein 2, Eritrea. *Emerg Infect Dis*. 2018;24:462–470.
- [12] Watson OJ, Verity R, Ghani AC, et al. Impact of seasonal variations in *Plasmodium falciparum* malaria transmission on the surveillance of *pfhrp2* gene deletions. *eLife*. 2019;8:e40339.
- [13] WHO | Country profiles [Internet]. WHO [cited 2020 Mar 26]. Available from: [http://www.who.int/malaria/publications/country-profiles/profile\\_dji\\_en.pdf?ua=1](http://www.who.int/malaria/publications/country-profiles/profile_dji_en.pdf?ua=1)
- [14] Ariey F, Witkowski B, Amaratunga C, et al. A molecular marker of artemisinin-resistant *Plasmodium falciparum* malaria. *Nature*. 2014;505:50–55.
- [15] WHO. Response plan to *pfhrp2* gene deletions. 2019 [cited 2020 Mar 26]. Available from: <https://apps.who.int/iris/handle/10665/325528>
- [16] Kamaliddin C, Le Bouar M, Berry A, et al. Assessment of diagnostic methods for imported malaria in mainland France. *Med Mal Infect*. 2020;50:141–160.
- [17] Lalloo DG, Shingadia D, Bell DJ, et al. UK malaria treatment guidelines 2016. *J Infect*. 2016;72:635–649.