

## Para -Substituted $\alpha$ -Phenyl- N - tert -butyl Nitrones

Anaïs Deletraz, Béatrice Tuccio, Julien Roussel, Maud Combes, Catherine Cohen-Solal, Paul-Louis Fabre, Patrick Trouillas, Michel Vignes, Noelle Callizot, Grégory Durand

### ► To cite this version:

Anaïs Deletraz, Béatrice Tuccio, Julien Roussel, Maud Combes, Catherine Cohen-Solal, et al.. Para - Substituted  $\alpha$ -Phenyl- N - tert -butyl Nitrones: Spin-Trapping, Redox and Neuroprotective Properties. ACS Omega, 2020, 5 (48), pp.30989-30999. 10.1021/acsomega.0c03907 . hal-03549080

## HAL Id: hal-03549080 https://ut3-toulouseinp.hal.science/hal-03549080

Submitted on 31 Jan2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





Article

# *Para*-Substituted $\alpha$ -Phenyl-*N*-tert-butyl Nitrones: Spin-Trapping, Redox and Neuroprotective Properties

Anaïs Deletraz, Béatrice Tuccio, Julien Roussel, Maud Combes, Catherine Cohen-Solal, Paul-Louis Fabre, Patrick Trouillas, Michel Vignes, Noelle Callizot, and Grégory Durand\*



was studied using two different in vitro models of cell death on primary cortical neurons injured by glutamate exposure or on glial cells exposed to <sup>t</sup>BuOOH. Trends between the protection afforded by the nitrones and their lipophilicity were observed. 4-CF<sub>3</sub>-PBN was the most potent agent against <sup>t</sup>BuOOH-induced oxidative stress on glial cells, while 4-Me<sub>2</sub>N-PBN showed potency in both models.

#### ■ INTRODUCTION

Oxidative stress results from an exaggerated production of reactive oxygen species (ROS). These include free radicals such as the superoxide ion  $O_2^{\bullet-}$  or the hydroxyl radical HO<sup>•</sup> to name two. However, reactive nitrogen species (RNS) and reactive sulfur species (RSS) must also be considered. Some of these reactive species are beneficial to living organisms as modulators of cellular function, signaling, and immune response; however, when they are present in high levels, they lead to cellular injury.<sup>1,2</sup> Oxidative stress is often associated with several pathologies such as cancer and cardiovascular and neurodegenerative diseases to name a few.<sup>1,3</sup> To prevent oxidative damage, synthetic antioxidants have been developed and nitrones are promising agents with considerable potential as therapeutics.<sup>4–6</sup> Nitrones have also been widely used as spin traps for the detection and characterization of free radicals by electron paramagnetic resonance (EPR) spectroscopy. In the spin-trapping technique, free radicals react with the nitronyl function to form stable and EPR-identifiable aminoxyl radicals.<sup>7,8</sup> Two families of nitrones have been developed: the cyclic ones derived from 5,5-dimethyl-1-pyrroline N-oxide (DMPO) and the linear ones derived from  $\alpha$ -phenyl-N-tertbutyl nitrone (PBN).

densities. Finally, the neuroprotective effect of these derivatives

Over the past several decades, several analogues of DMPO and PBN have been designed to improve their spin-trapping

properties and their biological activities.<sup>6,9</sup> When used as spin traps, it is important to design nitrones with a high rate constant of free radical trapping as well as with great stability of the corresponding aminoxyl spin adduct to ensure efficient detection. In general, cyclic nitrones lead to more persistent adducts than linear ones, and the analogues of DMPO with high potency have been designed. One can cite the phosphorylated analogue 5-diethoxyphosphoryl-5-methyl-1pyrrolidine N-oxide (DEPMPO),<sup>10</sup> the ester derivative 5ethoxycarbonyl-5-methyl-1-pyrrolidine N-oxide (EMPO),<sup>11</sup> and the amido analogue 5-carbamoyl-5-methyl-1-pyrrolidine N-oxide (AMPO).<sup>12,13</sup> AMPO was reported to have the highest rate constant of superoxide trapping, followed by EMPO and then DEPMPO. Linear nitrones have also been employed successfully in spin-trapping experiments but are generally less efficient than cyclic ones.<sup>14,15</sup> However, due to their better distribution within tissues and cells, linear nitrones have been widely used for radical detection in ex vivo and in

Received: August 13, 2020 Accepted: October 15, 2020 Published: November 20, 2020



vivo studies.<sup>16–18</sup> Linear nitrones are usually easier to synthesize and purify as they are often solid at room temperature and can be recrystallized to afford paramagnetic impurity-free samples. Moreover, linear PBN-type nitrones allow the possibility of rather easy functionalization both on the aromatic ring and on the *N-tert*-butyl group and therefore provide more chemical versatility. Several studies showed that PBN-type nitrones combine anti-inflammatory, antioxidant, and neuroprotective properties and are able to cross the blood-brain barrier.<sup>19,20</sup> These nitrones could therefore be used for the treatment of stroke,<sup>21</sup> visual loss,<sup>22,23</sup> neuronal damage,<sup>24</sup> and other age-related diseases.<sup>25</sup> The PBN derivative called 2,4-disulfophenyl-*N-tert*-butyl nitrone (NXY-059) was the first neuroprotective agent to reach phase III clinical trials in the United States.<sup>26,27</sup>

Nitrones have a very broad activity that depends on the nature and the position of the substituents on the nitronyl function. Therefore, the choice of substituents is very important and depends on the properties to be improved such as water solubility,<sup>28,29</sup> lipophilicity,<sup>29,30</sup> rate constant of radical trapping,<sup>31–33</sup> adduct stability,<sup>34</sup> bioactivity,<sup>23,35</sup> and the possibility of cellular or tissue targeting thanks to the ligation to specific molecular targets.<sup>36-41</sup> Over the past several years, the reactivity of the para-substituted derivatives of PBN has been explored to identify the most promising substituent for improved and optimal reactivity toward free radicals. It has been shown that the electronic nature of the substituent influences the rate of radical trapping on the nitronyl function. The presence of an electron-withdrawing group on the para position of the phenyl ring increased the reactivity toward nucleophilic addition reactions.<sup>31–33,42</sup> Contrarily, nitrones with an electron-donating group exhibited high reactivity toward electrophilic radicals.<sup>43,44</sup> The polar effect of the substituents has also been correlated with the electrochemical properties of the nitronyl function, with the derivatives bearing an electron-withdrawing group easier to reduce and harder to oxidize than those bearing an electron-donating group.<sup>33</sup>

In this work, we studied the spin-trapping properties of *para*substituted nitrones (4-iPr-PBN (3), 4-Ph-PBN (5), 4-MeS-PBN (6), 4-MeCONH-PBN (7), 4-F-PBN (8), 4-CF<sub>3</sub>O-PBN (9), and 4-CF<sub>3</sub>-PBN (12)) (Figure 1) as well as their electrochemical properties. The electronic effects of the substituents were computationally rationalized by density functional theory (DFT), and correlations between the experimental and theoretical data were established. Finally,



Figure 1. Chemical structures of DMPO, PBN, NXY-059, and X-PBNs studied in this work. Molecular weight, SMILES string, and PubChem IDs of all of the compounds are given in the Supporting Information (Table S1).

the in vitro protection of the nitrones including previously synthesized 4-X-PBN derivatives, namely,  $4-Me_2N-PBN$  (1), 4-MeO-PBN (2),  $4-AcNHCH_2-PBN$  (4), 4-MeNHCO-PBN (10), 4-HOOC-PBN (11), and 4-NC-PBN (13)<sup>33,42,45</sup> was investigated in two paradigms of cell death on neurons and glial cells.

#### RESULTS AND DISCUSSION

**Synthesis.** The derivatives 4-iPr-PBN (3), 4-Ph-PBN (5), 4-MeS-PBN (6), 4-F-PBN (8), 4-CF<sub>3</sub>O-PBN (9), and 4-CF<sub>3</sub>-PBN (12) were obtained by the "one-pot" reduction/ condensation of 2-methyl-2-nitropropane onto the appropriate benzaldehyde in the presence of zinc powder and AcOH, (method A in Scheme 1).<sup>46,47</sup> 4-MeCONH-PBN (7) was obtained by direct condensation of 4-acetamidobenzaldehyde with *N*-(*tert*-butyl)hydroxylamine acetate using pyrrolidine as a catalyst according to the methodology developed by Morales et al.<sup>48</sup> with a few modifications (method B in Scheme 1). The final compounds were purified by flash chromatography and two successive crystallizations, to ensure high purity. For 4-Me<sub>2</sub>N-PBN (1), 4-MeO-PBN (2), 4-AcNHCH<sub>2</sub>-PBN (4), 4-MeNHCO-PBN (10), 4-HOOC-PBN (11), and 4-NC-PBN (13), the samples already synthesized by our team were used.<sup>42</sup>

**EPR Study of Hydroxymethyl Radical Trapping.** To evaluate their ability to scavenge carbon-centered radicals, the relative rate constant of trapping of the hydroxymethyl radical (°CH<sub>2</sub>OH) by 4-iPr-PBN, 4-Ph-PBN, 4-MeS-PBN, 4-Me-CONH-PBN, 4-F-PBN, 4-CF<sub>3</sub>O-PBN, and 4-CF<sub>3</sub>-PBN was measured. Scheme 2 depicts the general mechanism of spin trapping by nitrones.

 $\alpha$ -Hydroxycarbon-centered radicals can be produced during oxidative stress by an attack of HO<sup>•</sup> on alcohols and are therefore of interest when evaluating nitrone antioxidant activities.<sup>49</sup> The Fenton reaction was used to produce in situ •CH<sub>2</sub>OH radicals in the presence of methanol. Under these conditions, all of the generated spin adducts of the °CH<sub>2</sub>OH radical (noted N-CH<sub>2</sub>OH) gave rise to a triplet of doublets EPR spectrum characteristic of such a nitroxide adduct, as shown in Figure 2.50 The EPR hyperfine splitting constants (hfsc's)  $a_N$  and  $a_H$  of the simulated  $^{\circ}CH_2OH$  adducts are reported in Table 1. All of the derivatives present similar nitrogen hfsc (~15.3 G) except 4-Ph-PBN (14.0 G), probably due to the higher resonance of the phenyl substituent. We next used a kinetic competition method described elsewhere<sup>45</sup> to determine the trapping efficiency of the compounds 4-iPr-PBN, 4-Ph-PBN, 4-MeS-PBN, 4-MeCONH-PBN, 4-F-PBN, 4-CF<sub>3</sub>O-PBN, and 4-CF<sub>3</sub>-PBN. The nitrone of interest, denoted as N, was tested in competition with 1,3,5-tri[(N-(1diethylphosphono)-1-methylethyl) N-oxy-aldimine] TN for •CH<sub>2</sub>OH trapping. A series of experiments were performed in the presence of various amounts of N and TN. The analysis of the experimental EPR spectra led to the relative trapping rate of the nitrones noted  $k_{\rm N}/k_{\rm TN}$ , as depicted in Figure 2C. To compare the new nitrones to PBN, the same experiment was performed using PBN versus TN, and from this, the  $k_{\rm N}/k_{\rm PBN}$ ratio was calculated for the whole series with values ranging from 0.25 to 3.18, as reported in Table 1. The details are given in the Experimental Section.

4-iPr-PBN, 4-MeS-PBN, 4-MeCONH-PBN, and 4-Ph-PBN trapped  $^{\circ}$ CH<sub>2</sub>OH slower than PBN, while 4-F-PBN and 4-CF<sub>3</sub>O-PBN trapped  $^{\circ}$ CH<sub>2</sub>OH 1.7 and 1.8 times faster than PBN, respectively. With a spin-trapping rate 3.2 times higher

Scheme 1. Methods Used for the Synthesis of 4-X-PBN Nitrones



Scheme 2. General Spin-Trapping Mechanism by Nitrones



than that observed for PBN, 4-CF<sub>3</sub>-PBN, which bears a strong electron-withdrawing substituent, was the most potent of the series even including the *para*-PBN derivatives previously studied.<sup>33,45</sup> The two very potent PBN-type nitrones that have been previously used as spin traps PPN and EPPN exhibited  $k_{\rm N}/k_{\rm PBN}$  values of 6.4 and 1.5, respectively.<sup>45</sup> This indicates the high efficiency of the fluorinated derivatives 4-F-PBN, 4-CF<sub>3</sub>O-PBN, and 4-CF<sub>3</sub>-PBN.

The plot of the rate constants by the different nitrones versus the Hammett values of their substituents showed a very good correlation (Figure 3A,  $R^2 = 0.86$ ). A positive value of the slope  $\rho$ , as observed, is indicative of the nucleophilic nature of the addition of the radical on the nitronyl function, which agrees with the work of De Vleeschouwer et al., who classified hydroxymethyl radicals as strong nucleophiles.<sup>52</sup> Moreover, it has been already shown that an electron-withdrawing substituent in PBN-type derivatives increases the reactivity of the nitronyl function for nucleophilic radical addition,<sup>13,53</sup> further supporting our observation.

The atomic partial charges of the nitronyl atoms (H, C, N, and O) and the atomic total charge of the nitronyl moiety were calculated for all of the derivatives using natural population analysis (NPA) within the natural bond orbital (NBO) framework and are reported in Table S2.<sup>54,55</sup> A positive



	EPR									
			•CH <sub>2</sub> OH							
nitrones	$\sigma_{\rm p}{}^a$	$a_{\rm N}$ (G)	$a_{\rm H}$ (G)	$k_{\rm N}/k_{\rm PBN}^{\ b}$ (± 0.05)						
4-iPr-PBN (3)	-0.15	15.4	3.8	0.25						
4-Ph-PBN (5)	-0.01	14.0	3.0	0.78						
4-MeS-PBN (6)	0	15.4	3.5	0.33						
PBN	0	15.3	3.8	1.00						
4-MeCONH- PBN (7)	0	15.3	3.6	0.58						
4-F-PBN (8)	0.06	15.4	3.4	1.72						
4-CF <sub>3</sub> O-PBN (9)	0.35	15.3	3.3	1.84						
4-CF <sub>3</sub> -PBN (12)	0.54	15.2	3.3	3.18						

<sup>*a*</sup>Data from Hansch et al.<sup>51</sup> <sup>*b*</sup>Ratio of the second-order rate constants for the hydroxymethyl radical trapping by various nitrones ( $k_{\rm N}$ ) and by PBN ( $k_{\rm PBN}$ ) in methanol, calculated with the ratio  $k_{\rm PBN}/k_{\rm TN} = 0.057$ .

correlation between the rate constants of  ${}^{\circ}\text{CH}_2\text{OH}$  addition to nitrones with the atomic total charge of the nitronyl moiety was observed (Figure 3B) as well as a good correlation between the atomic total charge of the nitronyl atoms of all of the series and the Hammett constants  $(\sigma_p)^{51}$  of the substituents (Figure S1). This confirms again that  ${}^{\circ}\text{CH}_2\text{OH}$  free radical scavenging occurs through nucleophilic radical addition requiring electron-withdrawing groups in the nitrones, which directly impact on the partial charges of the nitronyl moiety, where the nucleophilic addition is expected to occur.<sup>42</sup>



**Figure 2.** EPR signals of **TN** and **N** hydroxymethyl radical adducts, respectively. (A) **N** = PBN and (B) **N** = 4-CF<sub>3</sub>-PBN. The hydroxymethyl radical was generated by a Fenton system and the concentration ratio [N]/[TN] = 4. The peaks topped by a cross (X) correspond to the hydroxymethyl radical adduct of **N**. (C) Determination of the relative rate constants  $k_N/k_{TN}$  of •CH<sub>2</sub>OH trapping by PBN, 4-CF<sub>3</sub>-PBN, and 4-iPr-PBN.



**Figure 3.** Correlation of relative rate constants ( $k_{\rm N}/k_{\rm PBN}$ ) of •CH<sub>2</sub>OH addition to nitrones with (A) Hammett constants ( $\sigma_{\rm p}$ ) ( $R^2 = 0.86$ ) and with (B) atomic total charge of the nitronyl moiety ( $R^2 = 0.83$ , excluding the outlier 4-Ph-PBN marked as O).

Table 2. Electrochemical<sup>a</sup> Properties and Calculated Ionization Potentials<sup>b</sup> of 4-X-PBN Derivatives

	in $CH_3CN^c$ compound (1 mM)									
	$E_{\rm p}({\rm c}) ~({\rm V})^d$			$E_{\rm p}({\rm a})~({\rm V})$		stability domain (V) <sup>d</sup>	IP (eV)			
nitrones	first peak	second peak	third peak	first peak	second peak					
4-iPr-PBN (3)	-2.21	-2.30		1.52		3.73	5.9			
4-Ph-PBN (5)	-1.96	-2.56 (r)		1.52		3.48	5.9			
4-MeS-PBN (6)	-2.06			1.32	1.62	3.38	7.0			
PBN	-2.12	-2.27		1.60		3.72	6.0			
4-MeCONH-PBN (7)	-1.93	-2.35	-2.56	1.41	1.80	3.34	5.7			
4-F-PBN (8)	-2.10	-2.19		1.57	1.78	3.67	6.0			
4-CF <sub>3</sub> O-PBN (9)	-2.02	-2.13		1.64	1.86	3.66	6.1			
4-CF <sub>3</sub> -PBN (12)	-1.83	-2.13	-2.27	1.89		3.72	6.2			

<sup>*a*</sup>The peak potentials are given versus a silver wire electrode for a potential scan rate of 0.1 V/s; in general, the electron transfers appeared irreversible (no backward peak observed) except for nitrone 4-Ph-PBN noted (r). <sup>*b*</sup>The IPs were calculated at the (CPCM)/M06-2X/6-31+g(d,p) level of theory. <sup>*c*</sup>Containing 0.1 M TBAP with reduction  $E_p(c)$  and oxidation  $E_p(a)$  at a glassy carbon (GC) electrode. <sup>*d*</sup>The stability domain is  $E_p(a) - E_p(c)$ .

Cyclic Voltammetry. The electrochemical properties of 4iPr-PBN, 4-Ph-PBN, 4-MeS-PBN, 4-MeCONH-PBN, 4-F-PBN, 4-CF<sub>3</sub>O-PBN, and 4-CF<sub>3</sub>-PBN were investigated using cyclic voltammetry in acetonitrile containing tetra-butylammonium perchlorate (TBAP) as an electrolyte and the redox potentials are reported in Table 2. Reduction and oxidation potentials of all of the nitrones have been observed in the electroactivity domain of the solvent (Figures 4, S2, and S3). As expected, the peak currents are linearly related to the square root of the potential scan rate (Figures S4–S6), demonstrating that the process is diffusion-controlled.<sup>56,57</sup> By calibrating the current versus ferrocene, the number of electrons involved in the reduction or oxidation processes can be deduced.<sup>58</sup> PBN, 4-iPr-PBN, 4-Ph-PBN, 4-F-PBN, and 4-CF<sub>3</sub>O-PBN were reduced through two successive two-electron transfers in agreement with previous observations on other PBN-type nitrones.<sup>33,59,60</sup> For the derivatives 4-MeCONH-PBN and 4-CF<sub>3</sub>-PBN, an additional electron transfer was also observed, as shown in Figure 4. This is likely due to the reduction of the trifluoromethyl substituent of 4-CF<sub>3</sub>-PBN, while this corresponds to the reduction of the carbonyl group for 4-MeCONH-PBN. 4-MeS-PBN, which possesses a thiomethyl substituent, underwent a one-step two-electron reduction. For all of the derivatives except 4-Ph-PBN, the reduction appeared irreversible, as no associated backward peaks were observed. The reduction of the nitronyl group into a radical nitroxide anion corresponds to the first peak obtained, with values ranging from -2.21 to -1.83 V.

Considering the anodic behavior, PBN and the derivatives 4iPr-PBN, 4-Ph-PBN, and 4-CF<sub>3</sub>-PBN were oxidized through an irreversible one-electron transfer, whereas 4-MeS-PBN, 4-MeCONH-PBN, 4-F-PBN, and 4-CF<sub>3</sub>O-PBN possess two oxidation peaks. The first peak observed corresponds to the oxidation of the nitronyl function with values ranging from 1.41 to 1.89 V, except for 4-MeS-PBN where the first peak would correspond to the oxidation of the thiomethyl group.

4-CF<sub>3</sub>-PBN bearing an electron-withdrawing group is the easiest to reduce, with the lowest cathodic peak potential in the series and the hardest to oxidize. On the contrary, the derivative with the highest reduction potential was 4-iPr-PBN, which has an electron-donating group and very good correlations between the Hammett constants ( $\sigma_p$ ),<sup>S1</sup> and the redox potentials were observed (Figure S7). The ionization potentials (IPs) were also computationally determined (Table 2), and a good correlation between IP and the oxidation potential ( $R^2 = 0.71$ , Figure 4) was also observed.

In connection with the spin-trapping applications, the potential window where the spin-trapping technique can be safely applied with electrochemically generated radicals, refer to as the stability domain, was calculated and is reported in Table 2. It represents the potential window where there is no risk of an inverted spin-trapping process.<sup>61,62</sup> 4-iPr-PBN, 4-F-PBN, 4-CF<sub>3</sub>O-PBN, and 4-CF<sub>3</sub>-PBN have a similar stability domain to PBN (~3.70 V), whereas 4-Ph-PBN, 4-MeS-PBN, and 4-MeCONH-PBN present a slightly slower stability domain (~3.40 V). Therefore, all of the spin traps exhibit

Article



**Figure 4.** Cyclic voltammograms of PBN, 4-iPr-PBN, 4-MeCONH-PBN, and 4-CF<sub>3</sub>-PBN in acetonitrile containing 0.1 M TBAP at a GC electrode and potential scan rate  $\nu = 0.1$  V/s: (A) reduction and (B) oxidation. Inset: correlation of ionization potential with the oxidation potential of nitrones ( $R^2 = 0.71$ ), excluding the outlier 4-MeS-PBN for which a more complex oxidation process was observed. The correlation includes the values for 4-AcNHCH<sub>2</sub>-PBN, 4-MeNHCO-PBN, 4-HOOC-PBN, and 4-NC-PBN from Rosselin et al<sup>33</sup> and for 4-MeO-PBN from Deletraz et al.<sup>45</sup>

potential windows broad enough to be used in electrochemical investigations.<sup>60,63</sup>

**Cell Culture and Viability Studies.** Excitotoxicity and oxidative stress trigger different mechanisms of cell death but there is strong evidence that both mechanisms may cooperate in inducing neuron cell death. Therefore, the in vitro cytoprotective effect of the derivatives was next evaluated on two types of assays. The first one relies on <sup>t</sup>BuOOH-induced

oxidative stress and the second one relies on glutamateinduced excitotoxicity. We added to the evaluation of other 4-X-PBN nitrones previously synthesized in our laboratory. This includes 4-Me<sub>2</sub>N-PBN, 4-MeO-PBN, 4-AcNHCH<sub>2</sub>-PBN, 4-MeNHCO-PBN, 4-HOOC-PBN, and 4-NC-PBN (Figure 1).  $^{33,42,45}$  In the first test, nitrones at 10  $\mu$ M were preincubated with glial cells for 24 h before the <sup>t</sup>BuOOH exposure (300  $\mu$ M), and the cell survival was evaluated by a 3-(4,5dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) assay (Figure 5). We checked the effect of 4-X-PBN derivatives at 10  $\mu$ M after 24 h of incubation (Figure S8). None of the nitrones tested showed any toxicity at this concentration with a survival rate being comprised between 98.4  $\pm$  4.2 and 103.4  $\pm$  10.9%. Exposure of 300  $\mu$ M <sup>t</sup>BuOOH resulted in a significant drop ( $\sim 2.5$  times) of the cell viability indicating high toxicity of <sup>t</sup>BuOOH toward glial cells. Preincubation with nitrones led to different levels of protection depending on the nature of the substituents. 4-MeO-PBN, 4-NC-PBN, 4-MeNHCO, and 4-Ph-PBN did not afford any protection with cellular viability very similar to that obtained with <sup>t</sup>BuOOH alone. PBN led to slight protection with a cellular viability of 45.3  $\pm$  1.4% while the control was 41.3  $\pm$ 2.0%. 4-AcNHCH2-PBN, 4-HOOC-PBN, 4-F-PBN, 4-MeS-PBN, 4-MeCONH-PBN, and 4-iPr-PBN led to moderate protection, which was however not statistically significant when compared to the <sup>t</sup>BuOOH control condition. In contrast, 4-CF<sub>3</sub>O-PBN, 4-Me<sub>2</sub>N-PBN, and 4-CF<sub>3</sub>-PBN exhibited statistically significant protection with viability of  $55.6 \pm 4.1$ , 56.9  $\pm$  4.5, and 57.8  $\pm$  4.5%, respectively. No correlation between the neuroprotection afforded by the nitrones and their spin trapping (Figure S9) or redox properties was observed; however, it has to be underlined that the most efficient derivative in this model, 4-CF<sub>3</sub>-PBN, exhibited the highest spin-trapping rate as well as the lowest reduction and the highest oxidation potentials. This indicates the peculiar properties of this derivative. Previous work showed that 4-CF<sub>3</sub>-PBN efficiently protected rat retina from light damage when administered intraperitoneal.<sup>23</sup> Our data therefore confirm the potency of  $4-CF_3$ -PBN to reduce oxidative stress and cell death.

In the second assay, the nitrones were preincubated with primary cortical neurons (consisting of 80% neurons and 20% glial cells; mainly astrocytes) for 1 h at different concen-



Figure 5. Antioxidant effect of 4-X-PBN derivatives at 10  $\mu$ M on glial cells challenged by <sup>t</sup>BuOOH (300  $\mu$ M, 24 h), after 24 h of incubation with nitrones. Cell survival evaluation: MTT assay. Significance was accepted with \*p < 0.05 versus <sup>t</sup>BuOOH condition by the one-way analysis of variance (ANOVA) test.



**Figure 6.** Neuroprotective effect of PBN derivatives at 0.1, 1, and 10  $\mu$ M on primary cortical neurons injured by glutamate (20  $\mu$ M, 20 min, evaluation performed 48 h after the glutamate washout) after 1 h of incubation with nitrones. Cell survival evaluation: MTT assay. Significance was accepted with \**p* < 0.05 versus glutamate condition by one-way ANOVA, followed by PLSD Fisher's test.

trations, that is, 0.1, 1, and 10  $\mu$ M before 20 min of excitotoxic glutamate exposure (20  $\mu$ mol/L), followed by a washout. Glutamate concentrations in the extracellular space are low (nanomolar concentrations) and tightly controlled by a large number of mechanisms operating at the synapse. Disturbances in this regulatory system can have deleterious effects such as the excessive release of glutamate (micromolar concentration), which can induce hyperexcitability of post-synaptic neurons to the point of excitotoxicity, followed by cell death (cytotoxicity).<sup>64,65</sup> Glutamate excitotoxicity is one of the well-accepted pathophysiological mechanisms behind many neurodegenerative diseases like amyotrophic lateral sclerosis (ALS), Alzheimer's, and Huntington's disease.<sup>66</sup> Glutamate, the major excitatory neurotransmitter of the central nervous system, acts as a neurotoxin at high concentrations. High glutamate concentration causes an overstimulation of glutamate receptors and an increase in the intracellular calcium level, which can further activate various enzymes such as proteases, endonucleases, phospholipases, and nitric oxide synthase (NOS). This results in enhanced structural degradation, mitochondrial damage, ROS/RNS production, DNA damage, and increased expression of inflammatory mediators, which eventually lead to neuronal death.

Forty-eight hours after the glutamate washout, cell survival was evaluated by an MTT assay (Figure 6). In the absence of nitrone, the survival was dropped by about 30%, demonstrating the toxicity of glutamate exposure. Brain-derived neurotrophic factor (BDNF), a well-known neurotrophin acting via its TrkB receptor, was used as a positive control. The stimulation of the signaling cascades includes the phosphatidylinositol-3-kinase (PI3K), PLC $\gamma$ , and MAPK pathways. All these pathways promote cell survival and aid in cell growth and differentiation.<sup>67</sup> 4-MeO-PBN, 4-iPr-PBN, 4-MeS-PBN, 4-Me-CONH-PBN, and the three fluorinated derivatives 4-F-PBN, 4-CF<sub>3</sub>-PBN, and 4-CF<sub>3</sub>O-PBN failed to show any protection against glutamate, and the cellular viability was in certain conditions even lower than the control glutamate, suggesting slight toxicity. In contrast, a dose-dependence effect was noted for PBN, 4-MeNHCO-PBN, 4-Me2N-PBN, 4-AcNHCH2-PBN, 4-Ph-PBN, and 4-NC-PBN, but only 4-Me<sub>2</sub>N-PBN, 4-MeNHCO-PBN, and 4-NC-PBN led to statistically significant protection at 10  $\mu$ M with cellular viability of 78.8  $\pm$  2.4, 79.9  $\pm$ 3.0, and 81.7  $\pm$  3.7%, respectively. Irrespective of the

concentration (0.1, 1, or 10  $\mu$ M), treatment with 4-HOOC-PBN led to a significant protection higher than 80%, indicating a plateau effect.

Previous studies showed that, in vitro, PBN-protected rat cerebellar neurons against excitotoxic glutamate exposure (100  $\mu$ M) with a half-maximal effective concentration (EC<sub>50</sub>) of 2.7 mM.<sup>68</sup> The ability of PBN to alleviate glutamate neurotoxicity in vivo was also demonstrated but was attributed to the free radical scavenging mechanism of PBN.<sup>69</sup> The absence of correlation between the neuroprotection and the spin trapping (Figure S9) or redox properties was also noted in this assay.

Taken together, these data show that 4-Me<sub>2</sub>N-PBN is able to protect neurons and glial cells from a glutamate injury and is also able to reduce glial cell death induced by massive oxidative stress. Although 4-Me<sub>2</sub>N-PBN was found to be a poor spin trap,<sup>33</sup> it exhibits here the lowest ionization potential of the series. Moreover, its strong antioxidant potency was noted on the oxygen radical absorbance capacity (ORAC) test, the activity being superior to curcumin, while none of the other nitrones presented here showed any activity (data not shown). In contrast, 4-CF<sub>3</sub>-PBN, which was the most potent spin trap, was also the most potent agent against the <sup>t</sup>BuOOH astrocyte toxicity but failed to show any significant protection of neurons. Altogether, these results indicate that the cell protection afforded by nitrones depends on the type of toxicity (oxidative stress or excitotoxic event) as well as on the type of cells (neurons or glial cells). However, the mechanisms leading to cell death triggered in neurons and glial cells are rather similar in these cell types, as they both involve oxidative stress. Some evidence suggest that antioxidants may act directly by scavenging free radicals or indirectly by increasing endogenous cellular antioxidant defenses such as activation of the nuclear factor erythroid 2 (Nrf2). The potential for Nrf2mediated transcription to protect from neurodegeneration resulting from oxidative stress mechanisms is a well-known neuroprotective mechanism.<sup>70</sup> The Nrf2 signaling pathway is closely associated with the protective effect on neurons exposed to oxidative stress insults. In addition, Nrf2 has been reported to inhibit apoptosis based on its effect on mitochondria-related apoptotic proteins such as Bcl-2 and Bax.<sup>71</sup> The activation of this pathway is also protective against glial cell death.<sup>72</sup> In addition, reductions in nitric oxide (NO) production modulate the expression of proteins, such as iNOS,



**Figure 7.** Correlation of octanol–water partition coefficient ( $c \log P$ ) of the nitrones with their protective effect at 10  $\mu$ M on (A) glial cells challenged by 'BuOOH ( $R^2 = 0.41$ ) and (B) primary cortical neurons injured by glutamate ( $R^2 = 0.42$ ), excluding the outlier 4-Ph-PBN marked as O.

or regulators of the genes encoding proinflammatory mediators, such as nuclear factor- $\kappa$ B. Particularly in cell survival, NF- $\kappa$ B is well known to promote the transcription of bcl-2 and inhibition of apoptosis proteins.<sup>73</sup> The difference in the protection by nitrones on the two models may originate from their intrinsic properties highlighted in a cell-specific manner. The lipophilicity of the nitrones could be a key factor in their protective action in either model. Indeed, as shown in Figure 7, a positive trend is noted when plotting the protection of nitrones against <sup>t</sup>BuOOH toxicity in glial cells and their lipophilicity. Conversely, the lipophilicity of nitrones could be detrimental to the protection of neurons challenged with glutamate.

#### CONCLUSIONS

A series of para-substituted PBN nitrones were studied to investigate the influence of the nature of the substituent on the spin-trapping, redox, and cytoprotective activities of the nitronyl function. The spin-trapping ability toward hydroxymethyl radicals showed that the presence of an electronwithdrawing group at the para position significantly increased the trapping rate. The derivative 4-CF<sub>2</sub>-PBN gave the best spin-trapping activity with a trapping rate 3.2 times higher than PBN, and it is therefore a potential candidate for spin-trapping experiments. The spin-trapping rate was positively correlated with the atomic total charge of the nitronyl function; the higher the charge, the faster the reaction. The electrochemical properties of the derivatives were studied and the polar effect of para substituents was confirmed. With a strong electronwithdrawing CF<sub>3</sub> group, 4-CF<sub>3</sub>-PBN was the easiest to reduce and the hardest to oxidize. 4-CF3-PBN also showed good protection against <sup>t</sup>BuOOH toxicity on glial cells but no protection of neurons challenged by glutamate exposure was observed. In contrast, 4-Me<sub>2</sub>N-PBN showed good protection on both models, which indicates broad neuroprotection of this agent. Finally, while no correlation between the cytoprotection and the spin-trapping and redox properties was observed, we demonstrated that the cytoprotection was linked to the lipophilicity of the nitrones. Increasing lipophilicity may improve the protection against <sup>t</sup>BuOOH-induced toxicity in glial cells where it could be detrimental to neurons challenged by glutamate. This work therefore provides insight into the design of nitrone derivatives that could be further used as spin traps and therapeutics.

#### EXPERIMENTAL SECTION

The general methods and procedures for the synthesis of the derivatives,  $c \log P$  determination, EPR measurements, hydroxymethyl spin-trapping kinetics, computational calculations, cyclic voltammetric measurements, and *in vitro* neuroprotective measurements are described in the Supporting Information. They were described by Deletraz et al.<sup>74</sup> except for the glial cell model.

General Procedure for the Synthesis of 4-X-PBNs (Except 4-MeCONH-PBN). Under an argon atmosphere and under stirring, the corresponding benzaldehyde (1.0 equiv), 2-methyl-2-nitropropane (2.0 equiv), and AcOH (6.0 equiv) were dissolved in dry EtOH. The mixture was cooled down to 0 °C, and then zinc powder (4.0 equiv) was slowly added to keep the temperature below 15 °C. The mixture was stirred at room temperature for 30 min and then heated overnight at 60 °C in the dark in the presence of molecular sieves (4 Å). The reaction mixture was filtered through a pad of Celite, and the solvent was removed under vacuum. The crude mixture was purified by flash chromatography (EtOAc/cyclohexane), followed by two successive crystallizations from EtOAc/*n*-hexane.

*N*-tert-Butyl- $\alpha$ -(4-isopropyl)phenylnitrone (3). Following the general procedure, the reaction of 4-isopropylbenzaldehyde (319 mg, 2.16 mmol), 2-methyl-2-nitropropane (444 mg, 4.31 mmol), AcOH (0.74 mL, 12.96 mmol), and Zn (562 mg, 8.64 mmol) in dry EtOH for 16 h gave, after flash chromatography (EtOAc/cyclohexane, 2:8 v/v), nitrone 3 (284 mg, 60%) as a white powder.  $R_f$  (EtOAc/cyclohexane, 2:8 v/v) = 0.18. Elemental analysis calculated for C<sub>14</sub>H<sub>21</sub>NO: C, 76.67; H, 9.65; N, 6.39; found: C, 77.17; H, 9.49; N, 5.88%. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.22 (2H, d, J = 8.4 Hz), 7.51 (1H, s), 7.27 (2H, d, J = 8.4 Hz), 2.98-2.89 (1H, m), 1.61 (9H, s), 1.25 (6H, d, J = 6.8 Hz); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$ 151.5, 129.9, 129.1, 128.9, 126.6, 70.6, 34.3, 28.5, 23.9. HPLC (XTerra, Waters, RP18; 280 nm) t<sub>R</sub>: 11.6 min, 98.7% purity. HR-MS (ESI/Q-TOF) m/z:  $[M + H]^+$  calcd for  $C_{14}H_{22}NO_{14}$ 220.1701; found, 220.1704. The spectral data were in agreement with the literature.<sup>75</sup>

*N-tert-Butyl-* $\alpha$ -(4-phenyl)phenylnitrone (5). Following the general procedure, the reaction of 4-biphenylcarboxaldehyde (910 mg, 5.00 mmol), 2-methyl-2-nitropropane (1.03 g, 10.00 mmol), AcOH (1.72 mL, 30.00 mmol), and Zn (1.30 g, 20.00 mmol) in dry EtOH for 20 h gave, after flash chromatography (EtOAc/cyclohexane, 2:8 v/v), nitrone 5 (718 mg, 57%) as a white powder.  $R_f$  (EtOAc/cyclohexane, 3:7 v/v) = 0.32.

Elemental analysis calculated for C<sub>17</sub>H<sub>19</sub>NO: C, 80.60; H, 7.56; N, 5.53; found: C, 80.82; H, 7.34; N, 5.40%. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.37 (2H, d, *J* = 8.4 Hz), 7.68-7.63 (4H, m), 7.59 (1H, s), 7.47–7.43 (2H, m), 7.38–7.36 (1H, m), 1.63 (9H, s); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  142.7, 140.4, 130.2, 129.7, 129.4, 129.0, 127.9, 127.2, 127.1, 70.9, 28.5. HPLC (XTerra, Waters, RP18; 280 nm) *t*<sub>R</sub>: 12.6 min, 92.1% purity. HR-MS (ESI/Q-TOF) *m*/*z*: [M + H]<sup>+</sup> calcd for C<sub>17</sub>H<sub>20</sub>NO, 254.1545; found, 254.1546. The spectral data were in agreement with the literature.<sup>75</sup>

*N*-tert-Butyl- $\alpha$ -(4-methylthio)phenylnitrone (6). Following the general procedure, the reaction of 4-(methylthio)benzaldehyde (455 mg, 2.99 mmol), 2-methyl-2-nitropropane (616 mg, 5.98 mmol), AcOH (1.03 mL, 17.94 mmol), and Zn (777 mg, 11.96 mmol) in dry EtOH for 18 h gave, after flash chromatography (EtOAc/cyclohexane, 3:7 v/v), nitrone 6 (259 mg, 39%) as a white powder.  $R_f$  (EtOAc/cyclohexane, 3:7 v/v) = 0.20. Elemental analysis calculated for C<sub>12</sub>H<sub>17</sub>NOS: C, 64.54; H, 7.67; N, 6.27; found: C, 64.05; H, 7.79; N, 5.57%. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.22 (2H, d, J = 8.4 Hz), 7.49 (1H, s), 7.25 (2H, d, I = 8.4 Hz), 2.51 (3H, s), 1.61 (9H, s);<sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  141.5, 129.5, 129.2, 127.8, 125.6, 70.7, 28.5, 15.3. HPLC (XTerra, Waters, RP18; 280 nm)  $t_{\rm R}$ : 9.5 min, 99.3% purity. HR-MS (ESI/Q-TOF) m/  $z: [M + H]^+$  calcd for  $C_{12}H_{18}NOS$ , 224.1109; found, 224.1112. The spectral data were in agreement with the literature.<sup>76</sup>

*N*-tert-Butyl- $\alpha$ -(4-fluoro)phenylnitrone (8). Following the general procedure, the reaction of 4-fluorobenzaldehyde (410 mg, 3.31 mmol), 2-methyl-2-nitropropane (682 mg, 6.62 mmol), AcOH (1.14 mL, 19.86 mmol), and Zn (861 mg, 13.24 mmol) in dry EtOH for 20 h gave, after flash chromatography (EtOAc/cyclohexane, 3:7 v/v), nitrone 8 (357 mg, 55%) as a white powder.  $R_f$  (EtOAc/cyclohexane, 3:7 v/v = 0.25. Elemental analysis calculated for C<sub>11</sub>H<sub>14</sub>FNO: C, 67.67; H, 7.23; N, 7.17; found: C, 68.02; H, 7.01; N, 7.32%. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>)  $\delta$  8.35–8.32 (2H, m), 7.53 (1H, s), 7.10 (2H, t, J = 8.8 Hz), 1.61 (9H, s); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$  163.4 (d, J = 251 Hz), 131.1 (d, J = 8 Hz), 128.9, 127.6 (d, J = 3 Hz), 115.6 (d, J = 21 Hz), 70.9, 28.5. HPLC (XTerra, Waters, RP18; 280 nm) t<sub>R</sub>: 6.9 min, 98.1% purity. HR-MS (ESI/Q-TOF) m/z:  $[M + H]^+$  calcd for C11H15FNO, 196.1132; found, 196.1146. The spectral data were in agreement with the literature.<sup>7</sup>

N-tert-Butyl- $\alpha$ -(4-trifluoromethoxy)phenylnitrone (9). Following the general procedure, the reaction of 4-(trifluoromethoxy)benzaldehyde (531 mg, 2.79 mmol), 2methyl-2-nitropropane (575 mg, 5.58 mmol), AcOH (0.96 mL, 16.74 mmol), and Zn (725 mg, 11.16 mmol) in dry EtOH for 16 h gave, after flash chromatography (EtOAc/cyclohexane, 2:8 v/v), nitrone 9 (395 mg, 54%) as a white powder.  $R_f$  (EtOAc/cyclohexane, 2:8 v/v) = 0.19. Elemental analysis calculated for C<sub>12</sub>H<sub>14</sub>F<sub>3</sub>NO<sub>2</sub>: C, 55.17; H, 5.40; N, 5.36; found: C, 55.52; H, 5.25; N, 5.08%. <sup>1</sup>H NMR (400 MHz,  $CDCl_3$ )  $\delta$  8.36 (2H, d, J = 8.8 Hz), 7.56 (1H, s), 7.25 (2H, d, J = 8.8 Hz), 1.62 (9H, s);  ${}^{13}C{}^{1}H$  NMR (100 MHz, CDCl<sub>3</sub>)  $\delta$ 149.8, 130.4, 129.8, 128.5, 120.7, 120.5 (d, J = 256 Hz), 71.3, 28.5. HPLC (XTerra, Waters, RP18; 280 nm) t<sub>R</sub>: 11.7 min, 99.9% purity. HR-MS (ESI/Q-TOF) m/z: [M + H]<sup>+</sup> calcd for C12H15F3NO2, 262.1055; found, 262.1060. The spectral data were in agreement with the literature.

*N-tert-Butyl-* $\alpha$ -(4-trifluoromethyl)phenylnitrone (12). Following the general procedure, the reaction of 4-

(trifluoromethyl)benzaldehyde (433 mg, 2.49 mmol), 2methyl-2-nitropropane (513 mg, 4.98 mmol), AcOH (0.85 mL, 14.94 mmol) and Zn (647 mg, 9.96 mmol) in dry EtOH for 21 h gave, after flash chromatography (EtOAc/cyclohexane, 2:8 v/v), nitrone **12** (274 mg, 60%) as a white powder.  $R_f$  (EtOAc/cyclohexane, 1:9 v/v) = 0.13. Elemental analysis calculated for C<sub>12</sub>H<sub>14</sub>F<sub>3</sub>NO: C, 58.77; H, 5.75; N, 5.71; found: C, 59.14; H, 5.53; N, 5.57%. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.39 (2H, d, J = 8.4 Hz), 7.65 (2H, d, J = 8.4 Hz), 7.62 (1H, s), 1.63 (9H, s); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz, CDCl<sub>3</sub>) δ 134.3, 131.3 (d, J = 32 Hz), 128.8, 128.6, 125.5–125.3 (m), 124.0 (d, J = 269 Hz), 71.8, 28.5. HPLC (XTerra, Waters, RP18; 280 nm)  $t_{\rm R}$ : 11.6 min, 99.8% purity. HR-MS (ESI/Q-TOF) *m/z*: [M + H]<sup>+</sup> calcd for C<sub>12</sub>H<sub>15</sub>F<sub>3</sub>NO, 246.1106; found, 246.1111. The spectral data were in agreement with the literature.<sup>75</sup>

*N*-tert-Butyl- $\alpha$ -(4-acetamido)phenylnitrone (7). Under an argon atmosphere and under stirring, 4-acetamidobenzaldehyde (656 mg, 4.02 mmol, 1.0 equiv) and N-(tert-butyl)hydroxylamine acetate (599 mg, 4.02 mmol, 1.0 equiv) were dissolved in dry DCM. Pyrrolidine (0.40 mL, 4.82 mmol, 1.2 equiv) was added and the mixture was stirred at room temperature for 18 h. The solvent was removed under vacuum and the crude mixture was purified by flash chromatography (EtOAc/cyclohexane, 9:1 v/v), followed by two successive crystallizations from MeOH/Et<sub>2</sub>O to afford 500 mg of nitrone 7 (2.14 mmol, 53%) as a white powder.  $R_f$  (EtOAc/ cyclohexane, 8:2 v/v = 0.10. Elemental analysis calculated for C<sub>13</sub>H<sub>18</sub>N<sub>2</sub>O<sub>2</sub>: C, 66.64; H, 7.74; N, 11.96; found: C, 64.91; H, 7.63; N, 11.12%. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.24 (2H, d, J = 8.8 Hz), 7.97 (1H, bs), 7.58 (2H, d, J = 8.8 Hz), 7.50 (1H, s), 2.16 (3H, s), 1.59 (9H, s); <sup>13</sup>C{<sup>1</sup>H} NMR (100 MHz,  $CDCl_3$ )  $\delta$  168.7, 139.8, 130.0, 129.8, 126.9, 119.2, 70.6, 28.4, 24.8. HPLC (XTerra, Waters, RP18; 280 nm) t<sub>R</sub>: 3.6 min, 97.6% purity. HR-MS (ESI/Q-TOF) m/z: [M + H]<sup>+</sup> calcd for C13H19N2O2, 235.1447; found, 235.1449. The spectral data were in agreement with the literature.

#### ASSOCIATED CONTENT

#### **1** Supporting Information

The Supporting Information is available free of charge at https://pubs.acs.org/doi/10.1021/acsomega.0c03907.

Molecular weight, SMILES string, and PubChem IDs of the 4-X-PBN derivatives as well as PBN, DMPO, and NXY-059. Octanol–water partition coefficients ( $c \log P$ ) and calculated atomic partial charges of the nitronyl atoms and ionization potentials of 4-X-PBN derivatives; correlation of the  $\sigma_{\rm p}$  values with atomic total charge of nitrones; cyclic voltammograms of 4-F-PBN, 4-CF<sub>3</sub>O-PBN, 4-MeS-PBN, and 4-Ph-PBN; normalized cyclic voltammograms  $(I/\nu^{1/2} = f(E))$  of PBN and 4-MeS-PBN as a function of the potential scan rate  $\nu$ ; linear regressions corresponding to potential scan rate variations on PBN, 4-CF<sub>3</sub>-PBN, 4-iPr-PBN, and 4-MeCONH-PBN; correlation of the Hammett constants  $(\sigma_{\rm p})$  with the reduction potential and with the oxidation potential of the nitronyl function; effect of 4-X-PBN derivatives at 10  $\mu$ M on glial cells after 24 h of incubation with nitrones; plots of relative rate constants of <sup>•</sup>CH<sub>2</sub>OH addition to nitrones with the protective effect of nitrones at 10  $\mu$ M on glial cells and primary cortical neurons; general methods and procedures for the synthesis of the derivatives,  $c \log P$  determination,

EPR measurements, hydroxymethyl spin-trapping kinetics, computational calculations, cyclic voltammetric measurements, and in vitro neuroprotective measurements; and <sup>1</sup>H and <sup>13</sup>C NMR spectra, reverse-phase high-performance liquid chromatography (HPLC) chromatograms, and mass spectrometry spectra of 4-iPr-PBN, 4-Ph-PBN, 4-MeS-PBN, 4-MeCONH-PBN, 4-F-PBN, 4-CF<sub>3</sub>O-PBN, and 4-CF<sub>3</sub>-PBN (PDF)

#### AUTHOR INFORMATION

#### **Corresponding Author**

Grégory Durand – Institut des Biomolécules Max Mousseron, UMR 5247 CNRS-Université Montpellier-ENSCM & Avignon Université, Equipe Chimie Bioorganique et Systèmes Amphiphiles, Avignon 84916, France; orcid.org/0000-0001-6680-2821; Phone: +33 (0)4 9014 4445; Email: gregory.durand@univ-avignon.fr

#### Authors

- Anaïs Deletraz Institut des Biomolécules Max Mousseron, UMR 5247 CNRS-Université Montpellier-ENSCM & Avignon Université, Equipe Chimie Bioorganique et Systèmes Amphiphiles, Avignon 84916, France
- **Béatrice Tuccio** Aix-Marseille Université, CNRS, ICR UMR 7273, Avenue Escadrille Normandie Niemen, 13397 Marseille, France
- Julien Roussel Institut des Biomolécules Max Mousseron, UMR 5247 CNRS-Université Montpellier-ENSCM-Site faculté des Sciences, 34095 Montpellier, France
- Maud Combes Neuro-Sys, 13120 Gardanne, France
- Catherine Cohen-Solal Institut des Biomolécules Max Mousseron, UMR 5247 CNRS-Université Montpellier-ENSCM-Site faculté des Sciences, 34095 Montpellier, France
- Paul-Louis Fabre Pharma-Dev, UMR152, Université de Toulouse, 31400 Toulouse, France
- Patrick Trouillas INSERM U1248 IPPRITT, Univ. Limoges, Faculté de Médecine et Pharmacie, 87000 Limoges, France; Regional Centre of Advanced Technologies and Materials, Faculty of Science, Palacký University, 771 46 Olomouc, Czech Republic
- Michel Vignes Institut des Biomolécules Max Mousseron, UMR 5247 CNRS-Université Montpellier-ENSCM-Site faculté des Sciences, 34095 Montpellier, France; orcid.org/ 0000-0003-0969-2966
- Noelle Callizot Neuro-Sys, 13120 Gardanne, France

Complete contact information is available at: https://pubs.acs.org/10.1021/acsomega.0c03907

#### Notes

The authors declare no competing financial interest.

#### ACKNOWLEDGMENTS

A.D. was the recipient of a fellowship from the "Région Provence Alpes Côte d'Azur". We thank Fédération RENARD IR-RPE CNRS 3443 and "Structure Fédérative de Recherche" TERSYS for providing funds for the EPR studies; CALI for computational facilities and the National Program of Sustainability I from the Ministry of Youth, Education and Sports of the Czech Republic (LO1305); and Jamila Chetouani for her excellent technical assistance in acquiring the computational data. This study was conducted with the financial support of the European Regional Development Fund, the French Government, the "Région Provence Alpes Côte d'Azur", the Departmental Council of Vaucluse, and the Urban Community of Avignon. We thank Dr Micheál Moloney for carefully proofreading the article.

#### REFERENCES

(1) Halliwell, B.; Gutteridge, J. M. C. Free Radicals in Biology and Medicine; Oxford University Press, 2015.

(2) Finkel, T.; Holbrook, N. J. Oxidants, Oxidative Stress and the Biology of Ageing. *Nature* 2000, 408, 239–247.

(3) Villamena, F. A. Reactive Species Detection in Biology: From Fluorescence to Electron Paramagnetic Resonance Spectroscopy; Elsevier, 2017.

(4) Floyd, R. A.; Kopke, R. D.; Choi, C.-H.; Foster, S. B.; Doblas, S.; Towner, R. A. Nitrones as Therapeutics. *Free Radical Biol. Med.* **2008**, 45, 1361–1374.

(5) Villamena, F. A.; Das, A.; Nash, K. M. Potential Implication of the Chemical Properties and Bioactivity of Nitrone Spin Traps for Therapeutics. *Future Med. Chem.* **2012**, *4*, 1171–1207.

(6) Rosselin, M.; Poeggeler, B.; Durand, G. Nitrone Derivatives as Therapeutics: From Chemical Modification to Specific-Targeting. *Curr. Top. Med. Chem.* **2017**, *17*, 2006–2022.

(7) Villamena, F. A.; Zweier, J. L. Detection of Reactive Oxygen and Nitrogen Species by EPR Spin Trapping. *Antioxid. Redox Signaling* **2004**, *6*, 619–629.

(8) Hawkins, C. L.; Davies, M. J. Detection and Characterisation of Radicals in Biological Materials Using EPR Methodology. *Biochim. Biophys. Acta, Gen. Subj.* **2014**, *1840*, 708–721.

(9) Goldstein, S.; Lestage, P. Chemical and Pharmacological Aspects of Heteroaryl-Nitrones. *Curr. Med. Chem.* **2000**, *7*, 1255–1267.

(10) Frejaville, C.; Karoui, H.; Tuccio, B.; Le Moigne, F.; Culcasi, M.; Pietri, S.; Lauricella, R.; Tordo, P. 5-(Diethoxyphosphoryl)-5-Methyl-1-Pyrroline N-Oxide: A New Efficient Phosphorylated Nitrone for the in Vitro and in Vivo Spin Trapping of Oxygen-Centered Radicals. J. Med. Chem. **1995**, 38, 258–265.

(11) Olive, G.; Mercier, A.; Le Moigne, F.; Rockenbauer, A.; Tordo, P. 2-Ethoxycarbonyl-2-Methyl-3,4-Dihydro-2H-Pyrrole-1-Oxide: Evaluation of the Spin Trapping Properties. *Free Radical Biol. Med.* **2000**, *28*, 403–408.

(12) Villamena, F. A.; Rockenbauer, A.; Gallucci, J.; Velayutham, M.; Hadad, C. M.; Zweier, J. L. Spin Trapping by 5-Carbamoyl-5-Methyl-1-Pyrroline *N* -Oxide (AMPO): Theoretical and Experimental Studies. *J. Org. Chem.* **2004**, *69*, 7994–8004.

(13) Villamena, F. A.; Xia, S.; Merle, J. K.; Lauricella, R.; Tuccio, B.; Hadad, C. M.; Zweier, J. L. Reactivity of Superoxide Radical Anion with Cyclic Nitrones: Role of Intramolecular H-Bond and Electrostatic Effects. J. Am. Chem. Soc. 2007, 129, 8177–8191.

(14) Allouch, A.; Roubaud, V.; Lauricella, R.; Bouteiller, J.-C.; Tuccio, B. Preparation and Use as Spin Trapping Agents of New Ester-Nitrones. *Org. Biomol. Chem.* **2003**, *1*, 593–598.

(15) Roubaud, V.; Dozol, H.; Rizzi, C.; Lauricella, R.; Bouteiller, J.-C.; Tuccio, B. Poly( $\beta$ -Phosphorylated Nitrones): Preparation and Characterisation of a New Class of Spin Trap. *J. Chem. Soc., Perkin Trans.* 2 2002, 5, 958–964.

(16) Hughes, H. M.; George, I. M.; Evans, J. C.; Rowlands, C. C.; Powell, G. M.; Curtis, C. G. The Role of the Liver in the Production of Free Radicals during Halothane Anaesthesia in the Rat. *Biochem. J.* **1991**, 277, 795–800.

(17) Kadiiska, M. B.; Hanna, P. M.; Jordan, S. J.; Mason, R. P. Electron Spin Resonance Evidence for Free Radical Generation in Copper-Treated Vitamin E- and Selenium-Deficient Rats: In Vivo Spin-Trapping Investigation. *Mol. Pharmacol.* **1993**, *44*, 222–227.

(18) Connor, H. D.; Gao, W.; Nukina, S.; Lemasters, J. J.; Mason, R. P.; Thurman, R. G. Evidence That Free Radicals Are Involved in Graft Failure Following Orthotopic Liver Transplantation in the Rat - An Electron Paramagnetic Resonance Spin Trapping Study. *Transplantation* **1992**, *54*, 199–204.

(19) Chioua, M.; Sucunza, D.; Soriano, E.; Hadjipavlou-Litina, D.; Alcázar, A.; Ayuso, I.; Oset-Gasque, M. J.; González, M. P.; Monjas, L.; Rodríguez-Franco, M. I.; Marco-Contelles, J.; Samadi, A.  $\alpha$ -Aryl- N-Alkyl Nitrones, as Potential Agents for Stroke Treatment: Synthesis, Theoretical Calculations, Antioxidant, Anti-Inflammatory, Neuroprotective, and Brain–Blood Barrier Permeability Properties. *J. Med. Chem.* **2012**, *55*, 153–168.

(20) Chioua, M.; Martínez-Alonso, E.; Gonzalo-Gobernado, R.; Ayuso, M. I.; Escobar-Peso, A.; Infantes, L.; Hadjipavlou-Litina, D.; Montoya, J. J.; Montaner, J.; Alcázar, A.; Marco-Contelles, J. New Quinolylnitrones for Stroke Therapy: Antioxidant and Neuroprotective (Z)- N - Tert -Butyl-1-(2-Chloro-6-Methoxyquinolin-3-Yl)Methanimine Oxide as a New Lead-Compound for Ischemic Stroke Treatment. J. Med. Chem. **2019**, 62, 2184–2201.

(21) Maples, K. R.; Green, A. R.; Floyd, R. A. Nitrone-Related Therapeutics: Potential of NXY-059 for the Treatment of Acute Ischaemic Stroke. *CNS Drugs* **2004**, *18*, 1071–1084.

(22) Mandal, M. N. A.; Moiseyev, G. P.; Elliott, M. H.; Kasus-Jacobi, A.; Li, X.; Chen, H.; Zheng, L.; Nikolaeva, O.; Floyd, R. A.; Ma, J.; Anderson, R. E.  $\alpha$ -Phenyl- *N-Tert* -Butylnitrone (PBN) Prevents Light-Induced Degeneration of the Retina by Inhibiting RPE65 Protein Isomerohydrolase Activity. *J. Biol. Chem.* **2011**, 286, 32491–32501.

(23) Stiles, M.; Moiseyev, G. P.; Budda, M. L.; Linens, A.; Brush, R. S.; Qi, H.; White, G. L.; Wolf, R. F.; Ma, J.; Floyd, R.; et al. others. PBN (Phenyl-N-Tert-Butylnitrone)-Derivatives Are Effective in Slowing the Visual Cycle and Rhodopsin Regeneration and in Protecting the Retina from Light-Induced Damage. *PLoS One* **2015**, *10*, No. e0145305.

(24) Willis, C. L.; Ray, D. E. Antioxidants Attenuate MK-801-Induced Cortical Neurotoxicity in the Rat. *NeuroToxicology* **200**7, *28*, 161–167.

(25) Floyd, R. A. Nitrones as Therapeutics in Age-Related Diseases. *Aging Cell* **2006**, *5*, 51–57.

(26) Shuaib, A.; Lees, K. R.; Lyden, P.; Grotta, J.; Davalos, A.; Davis, S. M.; Diener, H.-C.; Ashwood, T.; Wasiewski, W. W.; Emeribe, U. NXY-059 for the Treatment of Acute Ischemic Stroke. *N. Engl. J. Med.* **2007**, 357, 562–571.

(27) Diener, H.-C.; Lees, K. R.; Lyden, P.; Grotta, J.; Davalos, A.; Davis, S. M.; Shuaib, A.; Ashwood, T.; Wasiewski, W.; Alderfer, V.; Hardemark, H.-G.; Rodichok, L. NXY-059 for the Treatment of Acute Stroke: Pooled Analysis of the SAINT I and II Trials. *Stroke* **2008**, *39*, 1751–1758.

(28) Janzen, E. G.; Zawalski, R. C. Synthesis of Nitronyl Alcohols and Their Benzoate Esters. J. Org. Chem. 1978, 43, 1900-1903.

(29) Janzen, E. G.; West, M. S.; Kotake, Y.; DuBose, C. M. Biological Spin Trapping Methodology. III. Octanol-Water Partition Coefficients of Spin-Trapping Compounds. J. Biochem. Biophys. Methods **1996**, 32, 183–190.

(30) Balogh, G. T.; Szántó, Z.; Forrai, E.; Győrffy, W.; Lopata, A. Use of Reversed-Phase Liquid Chromatography for Determining the Lipophilicity of  $\alpha$ -Aryl-N-Cyclopropylnitrones. *J. Pharm. Biomed.* Anal. **2005**, *39*, 1057–1062.

(31) Greenstock, C. L.; Wiebe, R. H. Substituent Effects in the Kinetic Analysis of Free Radical Reactions with Nitrone Spin Traps. *Can. J. Chem.* **1982**, *60*, 1560–1564.

(32) Sueishi, Y.; Yoshioka, C.; Olea-Azar, C.; Reinke, L. A.; Kotake, Y. Substituent Effect on the Rate of the Hydroxyl and Phenyl Radical Spin Trapping with Nitrones. *Bull. Chem. Soc. Jpn.* **2002**, *75*, 2043–2047.

(33) Rosselin, M.; Tuccio, B.; Pério, P.; Villamena, F. A.; Fabre, P.-L.; Durand, G. Electrochemical and Spin-Trapping Properties of Para -Substituted  $\alpha$ -Phenyl- N - Tert -Butyl Nitrones. *Electrochim. Acta* **2016**, 193, 231–239.

(34) Stolze, K.; Udilova, N.; Rosenau, T.; Hofinger, A.; Nohl, H. Spin Adducts of Several N-2-(2-Alkoxycarbonyl-Propyl)- $\alpha$ -Pyridylnitrone Derivatives with Superoxide, Alkyl and Lipid-Derived Radicals. *Biochem. Pharmacol.* **2004**, *68*, 185–194. (35) Kim, S.; de A. Vilela, G. V. M.; Bouajila, J.; Dias, A. G.; Cyrino, F. Z. G. A.; Bouskela, E.; Costa, P. R. R.; Nepveu, F.  $\alpha$ -Phenyl-N-Tert-Butyl Nitrone (PBN) Derivatives: Synthesis and Protective Action against Microvascular Damages Induced by Ischemia/Reperfusion. *Bioorg. Med. Chem.* **2007**, *15*, 3572–3578.

(36) Quin, C.; Trnka, J.; Hay, A.; Murphy, M. P.; Hartley, R. C. Synthesis of a Mitochondria-Targeted Spin Trap Using a Novel Parham-Type Cyclization. *Tetrahedron* **2009**, *65*, 8154–8160.

(37) Robertson, L.; Hartley, R. C. Synthesis of N-Arylpyridinium Salts Bearing a Nitrone Spin Trap as Potential Mitochondria-Targeted Antioxidants. *Tetrahedron* **2009**, *65*, 5284–5292.

(38) Han, Y.; Liu, Y.; Rockenbauer, A.; Zweier, J. L.; Durand, G.; Villamena, F. A. Lipophilic  $\beta$ -Cyclodextrin Cyclic–Nitrone Conjugate: Synthesis and Spin Trapping Studies. *J. Org. Chem.* **2009**, *74*, 5369–5380.

(39) Abbas, K.; Hardy, M.; Poulhès, F.; Karoui, H.; Tordo, P.; Ouari, O.; Peyrot, F. Detection of Superoxide Production in Stimulated and Unstimulated Living Cells Using New Cyclic Nitrone Spin Traps. *Free Radical Biol. Med.* **2014**, *71*, 281–290.

(40) Hardy, M.; Poulhés, F.; Rizzato, E.; Rockenbauer, A.; Banaszak, K.; Karoui, H.; Lopez, M.; Zielonka, J.; Vasquez-Vivar, J.; Sethumadhavan, S.; Kalyanaraman, B.; Tordo, P.; Ouari, O. Mitochondria-Targeted Spin Traps: Synthesis, Superoxide Spin Trapping, and Mitochondrial Uptake. *Chem. Res. Toxicol.* **2014**, *27*, 1155–1165.

(41) Socrier, L.; Rosselin, M.; Choteau, F.; Durand, G.; Morandat, S. Cholesterol-Nitrone Conjugates as Protective Agents against Lipid Oxidation: A Model Membrane Study. *Biochim. Biophys. Acta, Biomembr.* **201**7, 1859, 2495–2504.

(42) Durand, G.; Choteau, F.; Pucci, B.; Villamena, F. A. Reactivity of Superoxide Radical Anion and Hydroperoxyl Radical with  $\alpha$ -Phenyl- N - Tert -Butylnitrone (PBN) Derivatives. J. Phys. Chem. A **2008**, 112, 12498–12509.

(43) Murofushi, K.; Abe, K.; Hirota, M. Substituent Effect on the Spin-Trapping Reactions of Substituted N-Benzylidene-t-Butylamine N-Oxides. *J. Chem. Soc., Perkin Trans.* 2 **1987**, *12*, 1829–1833.

(44) Abe, Y.; Seno, S.; Sakakibara, K.; Hirota, M. Spin Trapping of Oxygen-Centred Radicals by Substituted N-Benzylidene-Tert-Butylamine N-Oxides. J. Chem. Soc., Perkin Trans. 2 1991, 6, 897–903.

(45) Deletraz, A.; Zéamari, K.; Di Meo, F.; Fabre, P.-L.; Reybier, K.; Trouillas, P.; Tuccio, B.; Durand, G. Reactivities of MeO-Substituted PBN-Type Nitrones. *New J. Chem.* **2019**, *43*, 15754–15762.

(46) Huie, R.; Cherry, W. R. Facile One-Step Synthesis of Phenyl-Tert-Butylnitrone (PBN) and Its Derivatives. J. Org. Chem. **1985**, 50, 1531–1532.

(47) Choteau, F.; Tuccio, B.; Villamena, F. A.; Charles, L.; Pucci, B.; Durand, G. Synthesis of Tris-Hydroxymethyl-Based Nitrone Derivatives with Highly Reactive Nitronyl Carbon. *J. Org. Chem.* **2012**, *77*, 938–948.

(48) Morales, S.; Guijarro, F. G.; Alonso, I.; García Ruano, J. L.; Cid, M. B. Dual Role of Pyrrolidine and Cooperative Pyrrolidine/ Pyrrolidinium Effect in Nitrone Formation. *ACS Catal.* **2016**, *6*, 84–91.

(49) Phaniendra, A.; Jestadi, D. B.; Periyasamy, L. Free Radicals: Properties, Sources, Targets, and Their Implication in Various Diseases. *Indian J. Clin. Biochem.* **2015**, *30*, 11–26.

(50) Buettner, G. R. Spin Trapping: ESR Parameters of Spin Adducts 1474 1528V. Free Radical Biol. Med. **198**7, *3*, 259–303.

(51) Hansch, C.; Leo, A.; Taft, R. W. A Survey of Hammett Substituent Constants and Resonance and Field Parameters. *Chem. Rev.* **1991**, *91*, 165–195.

(52) De Vleeschouwer, F.; Van Speybroeck, V.; Waroquier, M.; Geerlings, P.; De Proft, F. Electrophilicity and Nucleophilicity Index for Radicals. *Org. Lett.* **2007**, *9*, 2721–2724.

(53) Sueishi, Y.; Yoshioka, D.; Yoshioka, C.; Yamamoto, S.; Kotake, Y. High Static Pressure Alters Spin Trapping Rates in Solution. Dependence on the Structure of Nitrone Spin Traps. *Org. Biomol. Chem.* **2006**, *4*, 896–901.

(54) Foster, J. P.; Weinhold, F. Natural Hybrid Orbitals. J. Am. Chem. Soc. 1980, 102, 7211-7218.

(55) Reed, A. E.; Weinhold, F. Natural Localized Molecular Orbitals. J. Chem. Phys. **1985**, 83, 1736–1740.

(56) Suryanarayanan, V.; Yoshihara, S.; Shirakashi, T. Electrochemical Behavior of Carbon Electrodes in Organic Liquid Electrolytes Containing Tetrafluoroborate and Hexafluorophosphate Anionic Species in Different Non-Aqueous Solvent Systems. *Electrochim. Acta* **2005**, *51*, 991–999.

(57) Nakahara, K.; Iwasa, S.; Iriyama, J.; Morioka, Y.; Suguro, M.; Satoh, M.; Cairns, E. J. Electrochemical and Spectroscopic Measurements for Stable Nitroxyl Radicals. *Electrochim. Acta* **2006**, *52*, 921– 927.

(58) Amatore, C.; Azzabi, M.; Calas, P.; Jutand, A.; Lefrou, C.; Rollin, Y. Absolute Determination of Electron Consumption in Transient or Steady State Electrochemical Techniques. *J. Electroanal. Chem. Interfacial Electrochem.* **1990**, 288, 45–63.

(59) McIntire, G. L.; Blount, H. N.; Stronks, H. J.; Shetty, R. V.; Janzen, E. G. Spin Trapping in Electrochemistry. 2. Aqueous and Nonaqueous Electrochemical Characterizations of Spin Traps. *J. Phys. Chem. A.* **1980**, *84*, 916–921.

(60) Acken, B. J.; Gallis, D. E.; Warshaw, J. A.; Crist, D. R. Electrochemical Behavior of C-Methoxy, Amino, Cyano, and Mercapto Nitrones. *Can. J. Chem.* **1992**, *70*, 2076–2080.

(61) Eberson, L. 'Inverted Spin Trapping'. Reactions between the Radical Cation of  $\alpha$ -Phenyl-N-Tert-Butylnitrone and Ionic and Neutral Nucleophiles. *J. Chem. Soc., Perkin Trans.* 2 **1992**, 1807–1813.

(62) Eberson, L. Inverted Spin Trapping. Part III. Further Studies on the Chemical and Photochemical Oxidation of Spin Traps in the Presence of Nucleophiles. *J. Chem. Soc., Perkin Trans.* 2 1994, 2, 171–176.

(63) Walter, T. H.; Bancroft, E. E.; McIntire, G. L.; Davis, E. R.; Gierasch, L. M.; Blount, H. N. Spin Trapping in Heterogeneous Electron Transfer Processes. *Can. J. Chem.* **1982**, *60*, 1621–1636.

(64) Choi, D. W. Glutamate Receptors and the Induction of Excitotoxic Neuronal Death. In *Progress in Brain Research*; Elsevier, 1994 Chapter 6, Vol. 100, pp 47–51.

(65) Doble, A. The Role of Excitotoxicity in Neurodegenerative Disease: Implications for Therapy. *Pharmacol. Ther.* **1999**, *81*, 163–221.

(66) Tanović, A.; Alfaro, V. Glutamate-related excitotoxicity neuroprotection with memantine, an uncompetitive antagonist of NMDA-glutamate receptor, in Alzheimer's disease and vascular dementia. *Rev. Neurol.* **2006**, *42*, 607–616.

(67) Sun, D.; Wurzelmann, M.; Romeika, J. Therapeutic Potential of Brain-Derived Neurotrophic Factor (BDNF) and a Small Molecular Mimics of BDNF for Traumatic Brain Injury. *Neural Regener. Res.* **2017**, *12*, 7–12.

(68) Yue, T.-L.; Gu, J.-L.; Lysko, P. G.; Cheng, H.-Y.; Barone, F. C.; Feuerstein, G. Neuroprotective Effects of Phenyl-t-Butyl-Nitrone in Gerbil Global Brain Ischemia and in Cultured Rat Cerebellar Neurons. *Brain Res.* **1992**, *574*, 193–197.

(69) Lancelot, E.; Revaud, M.-L.; Boulu, R. G.; Plotkine, M.; Callebert, J. Alpha-Phenyl-N-Tert-Butylnitrone Attenuates Excitotoxicity in Rat Striatum by Preventing Hydroxyl Radical Accumulation. *Free Radical Biol. Med.* **1997**, *23*, 1031–1034.

(70) Zhang, M.; An, C.; Gao, Y.; Leak, R. K.; Chen, J.; Zhang, F. Emerging Roles of Nrf2 and Phase II Antioxidant Enzymes in Neuroprotection. *Prog. Neurobiol.* **2013**, *100*, 30–47.

(71) Li, L.; Wu, J.; Luo, M.; Sun, Y.; Wang, G. The Effect of Heat Stress on Gene Expression, Synthesis of Steroids, and Apoptosis in Bovine Granulosa Cells. *Cell Stress Chaperones* **2016**, *21*, 467–475.

(72) Assis, L. C.; Straliotto, M. R.; Engel, D.; Hort, M. A.; Dutra, R. C.; de Bem, A. F.  $\beta$ -Caryophyllene Protects the C6 Glioma Cells against Glutamate-Induced Excitotoxicity through the Nrf2 Pathway. *Neuroscience* **2014**, *279*, 220–231.

(73) Zhu, M.; Liu, M.; Guo, Q.; Zhu, C.; Guo, J. Prolonged DADLE Exposure Epigenetically Promotes Bcl-2 Expression and Elicits Neuroprotection in Primary Rat Cortical Neurons via the PI3K/ Akt/NF-KB Pathway. Acta Pharmacol. Sin. 2018, 39, 1582–1589.

(74) Deletraz, A.; Zéamari, K.; Hua, K.; Combes, M.; Villamena, F. A.; Tuccio, B.; Callizot, N.; Durand, G. Substituted  $\alpha$ -Phenyl and  $\alpha$ -Naphthlyl-N-Tert-Butyl Nitrones: Synthesis, Spin-Trapping, and Neuroprotection Evaluation. *J. Org. Chem.* **2020**, *85*, 6073–6085.

(75) Song, X.; Qian, Y.; Ben, R.; Lu, X.; Zhu, H.-L.; Chao, H.; Zhao, J. Activation of C–H Bonds in Nitrones Leads to Iridium Hydrides with Antitumor Activity. *J. Med. Chem.* **2013**, *56*, 6531–6535.

(76) Kelly, M.; Kincaid, J.; Janagani, S. 2-Substituted and 4-Substituted Aryl Nitrone Compounds. U.S.Patent US2,556,270A12005.

(77) Ikeda, K.; Tatsuno, T.; Ogo, H.; Masumoto, S.; Fujibayashi, T.; Nagata, R. Nitrones Derivatives. U.S. Patent US6,194,461B12001.

(78) Hinton, R. D.; Janzen, E. G. Synthesis and Characterization of Phenyl-Substituted C-Phenyl-N-Tert-Butylnitrones and Some of Their Radical Adducts. J. Org. Chem. 1992, 57, 2646-2651.