



## Pythagoras' theorem for areas

Jean-Baptiste Hiriart-Urruty, Jean Pierre Quadrat, Jean Bernard Lasserre

### ► To cite this version:

Jean-Baptiste Hiriart-Urruty, Jean Pierre Quadrat, Jean Bernard Lasserre. Pythagoras' theorem for areas. The American Mathematical Monthly, 2001, 108 (6), pp.549-551. hal-04118478

**HAL Id: hal-04118478**

**<https://ut3-toulouseinp.hal.science/hal-04118478>**

Submitted on 6 Jun 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# NOTES

Edited by Jimmie D. Lawson and William Adkins

## Pythagoras' Theorem for Areas

Jean-P. Quadrat, Jean B. Lasserre, and Jean-B. Hiriart-Urruty

The theorem of Pythagoras relating the squares of the lengths of the sides of a right triangle is one of the key results in elementary geometry. At a more advanced level, one learns that the Pythagorean theorem extends to prehilbert spaces, but it still expresses a relation among *lengths* of vectors.

We incidentally came across a result relating the areas of the faces of a right tetrahedron whose analogy with the Pythagorean theorem is striking. Here it is.

**Theorem 1.** *Let  $OABC$  be a tetrahedron with three perpendicular triangular faces  $OAB$ ,  $OAC$ ,  $OBC$ , and “hypotenuse-face”  $ABC$ ; see Figure 1. Let  $S_1$ ,  $S_2$ ,  $S_3$  denote the areas of the perpendicular faces and let  $S$  denote the area of the hypotenuse-face. Then*

$$S^2 = S_1^2 + S_2^2 + S_3^2. \quad (1)$$




Figure 1. The right tetrahedron  $OABC$

Theorem 1 is very easy to prove; we derive a generalized version in the  $n$ -dimensional context.

When we came across the result of Theorem 1, which we propose to call *the Pythagorean theorem for areas*, we found it noteworthy and asked colleagues: is it original (of course we thought no ...)? is it well-known (maybe ...)? does it bear a name (who knows ...)? After several unsuccessful attempts, we finally found mention of this result; here is a brief historical account.

The result of Theorem 1 indeed appears in some old books of geometry, in chapters devoted to “Applications of vector calculus to analytical geometry in 3-dimensional space”; the areas are there calculated via vector products. A sample reference is [8, pp. 121–123]. According to [4, p. 98], it was very likely known to R. Descartes. Theorem 1 also appears in [7] (a “bible” for results in elementary geometry) where (p. 911–912) it is attributed to J.-P. Gua de Malves (published in his memoirs of 1783). However, the analogy with the Pythagorean theorem for lengths is nowhere pointed

out. Later on, in his famous work [3], L.N.M. Carnot states the result of Theorem 1 and refers to it as a “known result” [3, section 263, p. 311]; he also provides an extension to any convex polyhedron ([3, section 268, p. 313]), which is a generalization of the cosine theorem for triangles.

We now propose a generalization of Theorem 1 to  $n$ -dimensional spaces. Let the standard Euclidean affine space  $\mathbb{R}^n$  be marked off with  $(O; \vec{e}_1, \vec{e}_2, \dots, \vec{e}_n)$ , where  $\{\vec{e}_1, \vec{e}_2, \dots, \vec{e}_n\}$  is an orthonormal basis of the vector space  $\mathbb{R}^n$ . We consider there the compact convex polyhedron (or polytope, or  $n$ -simplex)  $\Omega_n$  described as follows:

$$\Omega_n := \left\{ (x_1, \dots, x_n) \in \mathbb{R}^n : \sum_{i=1}^n \frac{x_i}{a_i} \leq 1, x_i \geq 0, \text{ for all } i = 1, \dots, n \right\}, \quad (2)$$

where  $a_i > 0$  for all  $i = 1, \dots, n$ .

The “multi-orthogonal”  $\Omega_n$  is just a generalized version of the right tetrahedron of Theorem 1, and its structure from the convexity viewpoint is well-known ([1, p. 84], [2, p. 77–79], [6, p. 53]). Indeed,  $\Omega_n$  has

- $n + 1$  vertices: the origin  $O$  and the  $n$  points  $A_i$  defined as  $\vec{OA}_i = a_i \vec{e}_i$  for all  $i = 1, 2, \dots, n$ .
- $n + 1$   $(n - 1)$ -dimensional faces (also called *facets*):  $n$  of these facets contain the origin (obtained as convex hulls of  $O$  and  $n - 1$  points among the  $A_i$ ’s); we call them *facets issuing from the origin*. One facet does not contain the origin (obtained as the convex hull of the  $A_i$ ’s); we call it *hypotenuse facet*.

The  $(n - 1)$ -dimensional content of the facets of  $\Omega_n$  is called *area*, in analogy with the usual case with  $n = 3$ . The  $n$ -dimensional content of  $\Omega_n$ , called the “volume”  $V$  of  $\Omega_n$ , can be computed from the areas of facets by the following formula [1, p. 87]:

$$V = \frac{1}{n} (\text{area of a facet}) \times (\text{height from the vertex outside the considered facet}). \quad (3)$$

### Theorem 2. (an $n$ -Dimensional Version of the Pythagorean Theorem for Areas)

For the compact convex polyhedron  $\Omega_n$  in (2), the square of the area of the hypotenuse facet is equal to the sum of the squares of the areas of the  $n$  facets issuing from the origin.

For example, with  $n = 4$ , Theorem 2 gives a relation among volumes (in the usual sense of the word) of the four 3-dimensional faces:  $V^2 = V_1^2 + V_2^2 + V_3^2 + V_4^2$ .

*Proof.* To follow the proof, the reader is invited to take  $n = 3$  and keep in mind Figure 1. According to (3), if  $S_i$  denotes the area of the facet issuing from the origin “opposed” to the vertex  $A_i$ ,

$$V = \frac{1}{n} S_i \times \|\vec{OA}_i\| = \frac{1}{n} S_i a_i. \quad (4)$$

The height from the origin  $O$  to the hypotenuse facet is the distance from  $O$  to the affine hyperplane whose equation is  $\sum_{i=1}^n x_i/a_i = 1$  (the one containing all the vertices  $A_1, A_2, \dots, A_n$ ); it equals  $(\sum_{i=1}^n a_i^{-2})^{-1/2}$ . Hence, if  $S$  denotes the area of the

hypotenuse facet of  $\Omega_n$ , we infer from (3) that

$$V = \frac{1}{n} S \times \left( \sum_{i=1}^n \frac{1}{a_i^2} \right)^{-1/2}. \quad (5)$$

We infer from (4)

$$S_i^2 = n^2 V^2 \frac{1}{a_i^2} \quad \text{for all } i = 1, \dots, n$$

and from (5)

$$S^2 = n^2 V^2 \left( \sum_{i=1}^n \frac{1}{a_i^2} \right). \quad \blacksquare$$

**ACKNOWLEDGMENTS.** We thank our colleagues Carral, Guillemot, Wind (IREM Toulouse), and Kahane (Paris), who helped us find historical references, the referee who pointed out to us the masterpiece [3], and H.S.M. Coxeter (Toronto), who provided us with the reference [5] where Theorem 2 is given.

## REFERENCES

1. M. Berger, *Géométrie*, vol. 2, Nathan, Paris, 1990.
2. A. Brøndsted, *An Introduction to Convex Polytopes*, Springer-Verlag, New York, 1983.
3. L.N.M. Carnot, *Geométrie de Position*, Imprimerie Crapelet, Paris, 1803.
4. P. Costabel, *Descartes. Exercices pour la Géométrie des Solides (De Solidorum Elementis)*, Presses Universitaires de France, Paris, 1987.
5. P.S. Donchian and H.S.M. Coxeter, An  $n$ -dimensional extension of Pythagoras' Theorem, *Math. Gazette* **19** (1935) 206.
6. B. Grünbaum, *Convex Polytopes*, J. Wiley & Sons, London, 1967.
7. F. G-M, *Exercices de Géométrie, comprenant l'exposé des méthodes géométriques et 2000 questions résolues*, 6ième édition, J. Gabay, Paris, 1920, réédité par les Editions Jacques Gabay en 1991.
8. P. Nillus, *Leçons de calcul vectoriel. Tome I: Règles de calcul vectoriel - Applications à la géométrie élémentaire, à la trigonométrie et à la géométrie analytique*, Editions Eyrolles, Paris, 1931.

INRIA, Domaine de Voluceau Rocquencourt, B.P. 105 78153 Le Chesnay Cédex, France.  
 jean-pierre.quadrat@inria.fr

LAAS-CNRS, 7 Av. du Colonel Roche, 31077 Toulouse Cédex 4, France.  
 lasserre@laas.fr

Université Paul Sabatier, 118 route de Narbonne, 31062 Toulouse Cedex 4, France.  
 jbhu@cict.fr