



Erratum to 'Germline-focussed analysis of tumour-only sequencing: recommendations from the ESMO Precision Medicine Working Group'

D. Mandelker, M. Donoghue, S. Talukdar, C. Bandlamudi, P. Srinivasan, M. Vivek, S. Jezdic, H. Hanson, K. Snape, A. Kulkarni, et al.

► To cite this version:

D. Mandelker, M. Donoghue, S. Talukdar, C. Bandlamudi, P. Srinivasan, et al.. Erratum to 'Germline-focussed analysis of tumour-only sequencing: recommendations from the ESMO Precision Medicine Working Group'. *Annals of Oncology*, 2021, 32 (8), pp.1069-1071. 10.1016/j.annonc.2021.05.798 . hal-04324470

HAL Id: hal-04324470

<https://ut3-toulouseinp.hal.science/hal-04324470v1>

Submitted on 5 Dec 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

ERRATUM



Erratum to ‘Germline-focussed analysis of tumour-only sequencing: recommendations from the ESMO Precision Medicine Working Group’

[Annals of Oncology 30 (2019) 1221–1231]

D. Mandelker^{1*}, M. Donoghue², S. Talukdar³, C. Bandlamudi², P. Srinivasan², M. Vivek^{4,5}, S. Jezdic⁶, H. Hanson³, K. Snape³, A. Kulkarni⁷, L. Hawkes⁸, J.-Y. Douillard⁶, S. E. Wallace⁹, E. Rial-Sebbag¹⁰, F. Meric-Bernstam¹¹, A. George^{12,13}, D. Chubb¹³, C. Loveday¹³, M. Ladanyi^{1,4}, M. F. Berger^{1,2}, B. S. Taylor^{2,3,5} & C. Turnbull^{7,13,14,15*}

¹Department of Pathology; ²Marie-Josée and Henry R. Kravis Center for Molecular Oncology, Memorial Sloan Kettering Cancer Center, New York; ³Department of Clinical Genetics, St George’s University of London, London; ⁴Human Oncology and Pathogenesis Program; ⁵Department of Epidemiology and Biostatistics, Memorial Sloan Kettering Cancer Center, New York, USA; ⁶European Society for Medical Oncology (ESMO) Head Office, Lugano, Switzerland; ⁷Department of Clinical Genetics, Guy and St Thomas’ NHS Foundation Trust, London; ⁸Department of Clinical Genetics, Oxford University Hospitals NHS Foundation Trust, Oxford; ⁹Department of Health Sciences, University of Leicester, Leicester, UK; ¹⁰University of Toulouse, Toulouse, France; ¹¹Department of Investigational Cancer Therapeutics, The University of Texas MD Anderson Cancer Center, Houston, USA; ¹²Cancer Genetics Unit, The Royal Marsden NHS Foundation Trust, London; ¹³Division of Genetics and Epidemiology, Institute of Cancer Research, London; ¹⁴William Harvey Research Institute, Queen Mary University of London, London; ¹⁵Public Health England, London, UK

The publisher regrets that in the original publication figures 1, 2 and 3 were incorrectly presented. The correct figures 1, 2 and 3 are as follows.




Figure 1. Distribution of variant allele frequency observed in the tumour for variants of true germline origin which were (i) small insertion/deletions (ii) SNVs.

DOI of original article: <https://doi.org/10.1093/annonc/mdz136>

*Correspondence to: Dr Diana Mandelker, Department of Pathology, Memorial Sloan Kettering Cancer Center, 1275 York Avenue, New York, NY 10065, USA. Tel:+1-212-639-4820

E-mail: mandelkd@mskcc.org (D. Mandelker).

*Prof. Clare Turnbull, Division of Genetics and Epidemiology, Institute of Cancer Research, 123 Old Brompton Road, Kensington, London SW7 3RP, UK. Tel:+44-208-722-4175

E-mail: clare.turnbull@icr.ac.uk (C. Turnbull).

0923-7534/© 2021 The Author(s). Published by Elsevier Ltd on behalf of European Society for Medical Oncology. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Figure 2. Distribution of germline and somatic pathogenic variants detected upon tumour analysis. Only variants classified pathogenic/likely pathogenic AND above VAF threshold are included (blue, germline origin; red, somatic origin; numbers, total number of pathogenic variants observed in tumour).




Figure 3. Distribution of germline and somatic pathogenic variants detected upon tumour analysis for 30 high-actionability CSGs. (A) Offtumour and (B) on-tumour.

The publisher would like to apologise for any inconvenience caused.