



HAL
open science

Blood Biomarkers from Research Use to Clinical Practice

D. Angioni, J. Delrieu, O. Hansson, H. Fillit, P. Aisen, J. Cummings, J.R. Sims, J.B. Braunstein, M. Sabbagh, T. Bittner, et al.

► **To cite this version:**

D. Angioni, J. Delrieu, O. Hansson, H. Fillit, P. Aisen, et al.. Blood Biomarkers from Research Use to Clinical Practice: What Must Be Done? A Report from the EU/US CTAD Task Force. The Journal of prevention of Alzheimer's disease, 2022, 9 (4), pp.569-579. 10.14283/jpad.2022.85 . hal-04419100

HAL Id: hal-04419100

<https://ut3-toulouseinp.hal.science/hal-04419100>

Submitted on 26 Jan 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution 4.0 International License



Published in final edited form as:

J Prev Alzheimers Dis. 2022 ; 9(4): 569–579. doi:10.14283/jpad.2022.85.

Blood Biomarkers from Research Use to Clinical Practice: What Must Be Done? A Report from the EU/US CTAD Task Force

D. Angioni^{1,*}, J. Delrieu^{1,*}, O. Hansson², H. Fillit³, P. Aisen⁴, J. Cummings⁵, J.R. Sims⁶, J.B. Braunstein⁷, M. Sabbagh⁸, T. Bittner⁹, M. Pontecorvo¹⁰, S. Bozeat¹¹, J.L. Dage¹², E. Largent¹³, S. Mattke¹⁴, O. Correa¹⁵, L.M. Gutierrez Robledo¹⁶, V. Baldivieso¹⁷, D.R. Willis¹⁸, A. Atri^{19,20}, R.J. Bateman²¹, P-J. Ousset¹, B. Vellas^{1,22}, M. Weiner²³,

EU/US/CTAD Task Force

¹Gerontopole of Toulouse, Alzheimer's Disease Research and Clinical Center, Toulouse University Hospital, Toulouse, France

²Memory Clinic, Skåne University Hospital, Malmö, Sweden

³Alzheimer's Drug Discovery Foundation, New York, NY, USA

⁴USC Alzheimer's Therapeutic Research Institute, San Diego, CA, USA

⁵Chambers-Grundy Center for Transformative Neuroscience, Department of Brain Health, School of Integrated Health Science, University of Nevada Las Vegas (UNV), Las Vegas, Nevada, USA

⁶Lilly, Indianapolis, USA

⁷C(2)N Diagnostics, Saint Louis, MO, USA

⁸Barrow Neurological Institute, Phoenix Arizona, AZ, USA

⁹Genentech, A Member of the Roche Group, Basel, Switzerland

Open Access: This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits use, duplication, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license and indicate if changes were made.

Corresponding Author: Davide Angioni, Gerontopole of Toulouse, Alzheimer's Disease Research and Clinical Center, Toulouse University Hospital, Toulouse, France, davideangioni2@gmail.com.

*These authors have contributed equally to this work and share first authorship

Task force members: Sandrine Andrieu (Toulouse, France), Phyllis Barkman Ferrell (Indianapolis, USA), Russell Barton (Baltimore, USA), Monika Baudler (Basel, Switzerland), Navia Bradford (Cambridge, USA), Brady Burgess (San Francisco, USA), Brian Campbell (San Francisco, USA), Chris Dague (Malvern, USA), Francesca De Simone (Malvern, USA), Robert Dean (Indianapolis, USA), Jean-Cosme Dodart (Dallas, USA), Rachele Doody (Basel, Switzerland), Kristen Eisterhold (Indianapolis, USA), Stephane Epelbaum (Indianapolis, USA), Sven Eriksson (Stockholm, Sweden), Rianne Esquvel (Malvern, USA), Michela Gallagher (Baltimore, USA), Hideki Garren (San Francisco, USA), Sarah Giardina (New York, USA), Ishtar Govia (Kingston, Jamaica), Virginie Gregoire (Indianapolis, USA), Harald Hampel (Nutley, USA), Charles Harrington (Aberdeen, United Kingdom), James Hendrix (Indianapolis, USA), Dave Henley (Titusville, USA), Joseph Herring (North Wales, USA), Helen Hu (Nutley, USA), Farhad Imam (Kirkland, USA), Michael Irizarry (Nutley, USA), Peter Johannsen (Bagsvaerd, Denmark), Paul Kinnon (Solana Beach, USA), Hartmuth Kolb (Titusville, USA), Hisatomo Kowa (Kobe, Japan), Akihiko Koyama (Nutley, USA), Lynn Kramer (Nutley, USA), Lars Lannfelt (Uppsala, Sweden), John Lawson (Malvern, USA), Ferenc Martenyi (San Francisco, USA), Sarah McManus (Titusville, USA), David Miller (Blue Bell, USA), Sonya Miller (Aberdeen, United Kingdom), Richard Mohs (Baltimore, USA), Laura Nisenbaum (New York, USA), Gunilla Osswald (Stockholm, Sweden), Suzanne Ostrowitzki (San Francisco, USA), Lewis Penny (Aberdeen, United Kingdom), Elizabeth Pittampalli (San Francisco, USA), Craig Ritchie (Edinburgh, United Kingdom), Andrew Roche (Dublin, Ireland), Yves Rolland (Toulouse, France), Sharon Rosenzweig-Lipson (Baltimore, USA), Peter Schuler (Langen, Germany), Maria Eugenia Soto Martin (Toulouse), Reisa Sperling (Boston, USA), Erik Stoops (Gent, Belgium), Aishwarya Sukumar (Kirkland, USA), Charlotte Thim Hansen (Bagsvaerd, Denmark), Jacques Touchon (Montpellier, France), Martin Traber (Basel, Switzerland), Gallen Triana-Baltzer (Titusville, USA), Eugene Vanmechelen (Gent, Belgium), Huijing Yu (Dallas, USA), Wagner Zago (San Francisco, USA), Nancy Zhang (Titusville, USA).

10. Eli Lilly and Company, Indianapolis, IN, USA
11. Hoffmann-La Roche, Basel, Switzerland
12. Indiana University School of Medicine, Department of Neurology, Indianapolis, IN, USA
13. Department of Medical Ethics and Health Policy, University of Pennsylvania Perelman School of Medicine, Philadelphia, USA
14. Center for Economic & Social Research, University of Southern California, Los Angeles, USA
15. GRIS - Gestão Responsável Integrada de Saúde, Rio de Janeiro, Brazil
16. National Institute of Geriatrics, Mexico City, Mexico
17. AdventHealth Memory Care Clinic, Winter Park, USA
18. Department of Family Medicine, Indiana University School of Medicine, Indianapolis, USA
19. Banner Sun Health Research Institute, Banner Health, Sun City, USA
20. Department of Neurology, Brigham and Women's Hospital and Harvard Medical School, Boston, Massachusetts, USA
21. Washington University School of Medicine, St. Louis, MO, USA
22. UNM Division of General Internal Medicine and Geriatric Medicine, Albuquerque NM, USA
23. University of California, San Francisco, CA, USA

Abstract

Timely and accurate diagnosis of Alzheimer's disease (AD) in clinical practice remains challenging. PET and CSF biomarkers are the most widely used biomarkers to aid diagnosis in clinical research but present limitations for clinical practice (i.e., cost, accessibility). Emerging blood-based markers have the potential to be accurate, cost-effective, and easily accessible for widespread clinical use, and could facilitate timely diagnosis. The EU/US CTAD Task Force met in May 2022 in a virtual meeting to discuss pathways to implementation of blood-based markers in clinical practice. Specifically, the CTAD Task Force assessed: the state-of-art for blood-based markers, the current use of blood-based markers in clinical trials, the potential use of blood-based markers in clinical practice, the current challenges with blood-based markers, and the next steps needed for broader adoption in clinical practice.

Keywords

Alzheimer's disease; blood biomarkers; diagnostic; clinical trials; amyloid; p-tau; neurofilament light

Introduction

Timely and accurate diagnosis of Alzheimer's disease (AD) in clinical practice is a great challenge. Currently, the clinical diagnosis of AD is poor; about 20-25% of the patients are misdiagnosed if cerebrospinal fluid (CSF) or positron emission tomography (PET)

biomarkers are not used (1, 2). This misdiagnosis is particularly important during early disease stages with subtle or mild symptoms (subjective cognitive decline [SCD] and mild cognitive impairment [MCI]) and in primary care, where more than 50% of patients with cognitive impairment are not recognized or correctly diagnosed (2). This misdiagnosis results in suboptimal treatment and care, delayed or incorrect therapies, and inaccurate information about disease and prognosis (3).

In 2021, on the basis of the dramatic effect on cerebral amyloid-plaque load, aducanumab was granted accelerated approval for the treatment of AD in the United States (US) and phase III trials with other anti-amyloid monoclonal antibodies are ongoing (gantenerumab, donanemab, lecanemab) (4). If these potentially disease-modifying anti-amyloid therapies become approved and widely available for use in clinical practice, it will become even more important to establish an accurate early diagnosis. To date, PET and CSF biomarkers are the most widely used biomarkers in clinical practice but easily accessible, cost-effective, and accurate biomarkers such as blood-based markers (BBM) have the potential for widespread clinical use, including in primary care (3).

However, several issues remain to be overcome before BBM can be expected to represent a routine component of clinical care. A consensus on progress, barriers and next steps is needed to make efficient advances in the field.

The EU/US CTAD Task Force met in May 2022 in a virtual meeting to discuss the pathways to implementation of BBM use in clinical practice. In this objective, the CTAD Task Force assessed:

1. The state-of-scientific development of BBM.
2. The current use of BBM in clinical trials.
3. Limitations of the currently available data and gaps to be resolved
4. The potential use of BBM in clinical practice.
5. The next steps for broader adoption of BBM in clinical practice.

State-of-scientific development of blood-based markers

Amyloid and p-tau blood-based markers in AD

Plasma A β 42/40 ratio, p-tau, serum neurofilament light chain (NfL) and glial fibrillary acidic protein (GFAP) are the most advanced BBM developed for the diagnosis and prognosis of AD and to potentially monitor the effects of disease-modifying therapies (DMTs).

There are many assays to measure A β in blood and to detect positive amyloid status in patients with early AD. Plasma A β 42/A β 40 ratio, measured by high-precision assay, can detect abnormal amyloid status in cognitively impaired (5, 6), and cognitively unimpaired subjects even before amyloid PET reach its threshold of positivity (7). Due to their high-precision and robust implementation, mass spectrometry-based (MS) methods and fully automated immunoassay methods are considered to have the best opportunity for clinical

implementation in routine care; MS methods have been implemented as lab developed tests (LDTs) in the United States (8).

Similarly to A β isoforms, several forms of p-tau such as p-tau181, p-tau217 and p-tau231 can be detected in plasma with high accuracy (9–11). Plasma p-tau217 correlates with tau tangles in AD, but not in non-AD pathologies (12). P-tau217 levels appear to be elevated in the presence of amyloid plaques but not in the absence of amyloid plaques as occurs in primary age related tauopathy (PART) or other non-AD tauopathies (e.g., progressive supranuclear palsy or corticobasal degeneration) (13). Plasma p-tau217 levels are increased by 300-700% in symptomatic AD and can differentiate AD from non-AD diseases with an accuracy similar to CSF p-tau and Tau PET (12). In cognitively unimpaired subjects, plasma p-tau217 with or without A β 42/40 ratio can predict cerebral amyloid pathology assessed by amyloid PET (14). Plasma p-tau217 can also detect AD pathology in cognitively unimpaired subjects. Its level is increased by 80-350% in cognitively unimpaired subjects with positive amyloid status and changes up to 20 years before onset of cognitive decline (12, 15). In non-demented subjects with cognitive complaints, individual probability of conversion to AD has been assessed by a model combining plasma p-tau, apolipoprotein E (APOE) ϵ 4 genotype, and simple executive (animal fluency and trail making test part B) and memory (Alzheimer's Disease Assessment Scale–Cognitive Subscale memory subtests) tests. This model predicted dementia better than a clinical assessment performed by specialists in memory disorders and performed similarly to CSF biomarker-based algorithms (16).

Robustness of blood-based markers in clinical routine

There is a theoretical concern that plasma A β 42/40 ratio may have a lower diagnostic accuracy than the A β 42/40 ratio in CSF, representing a potential problem for clinical robustness. Indeed, A β 42/40 ratio is reduced by only 8-15% in blood while A β 42/p-tau ratio is reduced by more than 40-60% in CSF, making it challenging to apply a cutoff at the individual level. From the Alzheimer's Disease Neuroimaging Initiative (ADNI, n = 118 from cognitive unimpaired to dementia subjects) cohort, the performance of BBM has been evaluated by a theoretical simulation evaluating an increase in the inter-assay coefficient of variability (17). The robustness of plasma A β 42/40 ratio is quickly reduced with an increase of inter-assay variability. An increase of only 5% in coefficient of variability impacted its accuracy for discriminating amyloid positive PET versus negative cases. Cullen et al. measured the test-retest variability (random effect) of blood A β 42/A β 40 ratio, NfL, GFAP, and p-tau217 from the Biomarkers For Identifying Neurodegenerative Disorders Early and Reliably (BioFINDER) -1 study (n=399) (18). In this study, the test-retest variability reduced the performance of biomarkers for predicting abnormal amyloid accumulation and progression to AD dementia. Compared to the other analytes, p-tau217 was less affected by test-retest variability. Direct head-to-head studies have demonstrated that high precision and accurate plasma A β 42/A β 40 ratios perform as well or better than immunoassay measures (19) in identifying AD pathology. For example, a study of five different assays compared in the same cohort, the IP-MS plasma A β 42/A β 40 assay outperformed all immunoassays compared to amyloid PET or CSF A β as the reference standard (8). Multiple international studies, some with longitudinal collection and conducted over several years, have consistently shown performance characteristics of receiver operator curve (ROC) area

under the curve (AUC) of ~ 0.85 (5–7, 20–22). These performances are similar to several blood plasma p-tau immunoassays, and with rigorous stability that provides a benefit to aid in the diagnosis of AD. Further, results from Cullen et al. (18) shown that combining BBM reduced the impact of assay variability on biomarker performance. Recently, it was shown that in cognitive impairment at the earliest stages of pathological amyloid accumulation, the combination of p-tau217 with A β 42/40 may be more sensitive than p-tau217 alone for the detection of AD (23). Combined BBM could be used in clinical practice and can assist in avoiding misdiagnoses.

Confounding factors

Performances of p-tau181 and p-tau217 have been studied in heterogeneous community-based populations from the Mayo Clinic Study of Aging (n = 1329). This analysis showed the effect of comorbidities on plasma p-tau181 and p-tau217 levels. Increase in p-tau181 and p-tau217 were associated with age especially in subjects with a positive amyloid status. Several comorbidities including chronic kidney disease, hypertension, stroke, and myocardial infarction were associated with higher plasma p-tau181 and p-tau217 levels, even in an age- and sex-adjusted model (24). In BioFINDER-1 and -2 studies, plasma NfL, GFAP and p-tau levels were affected by kidney function and body mass index, but these potential confounding factors seem to have only small effects on the clinical performance of the biomarkers. Based on these data, a significant research priority is to better understand the effect of comorbidities on BBM for their proper interpretation in clinical practice.

A secondary analysis from the ADNI study shown that sex may impact the clinical interpretation of plasma p-tau181 concentrations (25). In women, high plasma p-tau levels were associated with worse phenotypic markers (i.e., greater cortical A β deposition, higher CSF p-tau181 levels, lower brain glucose metabolism) compared with men. Moreover, higher baseline p-tau levels in women were associated with greater risk of progression to dementia and faster cognitive decline, compared with men.

Race and ethnicity are also pertinent demographic variables that may impact the interpretation of BBM data in the clinical setting. A recent evaluation studied the performance of various BBM in a cohort of 76 matched pairs (91% cognitively normal) of African-American (AA) and Non-Hispanic White individuals (NHW) (26). The results suggested that a high-performance plasma A β 42/40 assay may provide a consistent and reliable measure of brain amyloidosis across AA and NHW groups, but models based on plasma p-tau181, p-tau231, and NfL, may perform inconsistently and could result in a disproportionate misdiagnosis of AA individuals. Special attention should be given to the use of algorithms including BBM and APOE genotype in populations without a predominant European ancestry. Indeed the association between APOE alleles and AD risk varies across different ethnic groups (27, 28). Additional studies are necessary to understand how to best interpret BBM results across racially and ethnically diverse populations for the benefit of clinical care of these patients.

Limitations

Despite the encouraging data available thus far, it should be emphasized that most of the data reviewed above come from previously collected, retrospective studies, where the plasma samples were analyzed in large batches, applying optimal cut points derived from batch data, and evaluating AUCs derived from these cut points. There are few prospective data available where plasma samples were obtained longitudinally over a long period of time and clinical performances calculated from a predetermined cut point. Such data are being developed through clinical studies and are recommended before implementation of any particular BBM or combination of BBM in clinical practice.

Current use of plasma biomarkers in clinical trials

The introduction of new BBM for AD will improve the design and conduct of clinical trials evaluating DMTs. For the screening of these trials, BBM can help identify individuals with a high probability of not having AD pathology. This approach is of particular interest in preclinical trials in AD and could significantly reduce the number of negative CSF and PET scans commonly seen in the screening phases of trials for putative DMTs (29).

Within the framework of clinical trials, three potential uses of BBM can be identified:

1. As pre-screening tools to eliminate individuals who are likely to be amyloid negative with CSF or PET to reduce the number of required confirmatory CSF and PET testing
2. As an inclusion criterion (diagnosis tool) without confirmatory CSF or PET testing
3. As pharmacodynamic markers to detect direct target engagement and monitor disease-modifying effects

Blood-based markers as screening tools

This use of BBM as a pre-screening tool is currently the most common. The obvious goal is to increase the prevalence of amyloid positive individuals before they undergo CSF or PET testing, which are needed to confirm the diagnosis of AD for final inclusion decisions for trials. This approach makes it possible to avoid unnecessary examinations and therefore reduce the cost and burden associated with these invasive and costly examinations. Biomarkers currently used in this pre-screening perspective include the A β 42/A β 40 ratio (30), p-tau181, p-tau217 (31), or algorithms considering, in addition to the biomarker, data such as age and APOE status (32, 33). Currently available data support the utility of plasma prescreening: for example, the preliminary analysis of the AHEAD 3-45 study demonstrated an improvement of positive predictive value from 28.9% PET eligible to 61.5% PET confirmed with a plasma prescreen (30). However, the value of BBM as screening tools prospectively needs to be confirmed.

Blood-based markers as an inclusion criterion

Use as the main inclusion criterion for trials, indicating Alzheimer's pathology, is conceivable only if the BBM can achieve sufficiently high diagnostic performance. This

is particularly advantageous for trials where CSF or PET is not required to demonstrate drug target engagement or pharmacodynamic response in each participant (34). Currently, only one trial, TRAILBLAZER-ALZ3, evaluating the efficacy of donanemab in cognitively normal people with elevated plasma p-tau 217, uses this recruitment strategy. This approach is based on the good discriminating performance established by the p-tau217 assay (12). This strategy seems particularly attractive for this type of trial including participants at the preclinical stage of AD, where participants are less likely to accept invasive diagnostic tests. However, at the current time there are limited data concerning the value of BBM used prospectively as inclusion criteria.

Blood-based markers as pharmacodynamic markers

BBM could also be used as pharmacodynamic markers to detect direct target engagement and disease modifying effects. Both effects are critical for identifying an appropriate dose and increasing confidence that the treatment leads to disease-relevant outcomes.

BBM could have utility as surrogate endpoints, resulting in shorter clinical trials or smaller sample sizes. In the early stages of drug development, they could help make go/no go decisions for later larger trials of drug candidates, before effects are demonstrated on measures of cognition or function (34). Plasma A β 42/A β 40 ratio has reflected amyloid removal, corroborated with amyloid PET, in several drug trials of amyloid-removing antibodies including gantenerumab and lecanemab (35, 36). There are promising data from epidemiological studies showing that plasma p-tau has properties that may make it suitable as a pharmacodynamic marker, including longitudinal increases in preclinical and prodromal stages, and longitudinal correlations with brain structure and cognitive measures (37). In the treatment arms of EMERGE and ENGAGE trials of aducanumab (4) a reduction of p-tau181 levels was observed over time in concert with amyloid plaque lowering. Conversely, p-tau181 levels increased in the placebo groups. A greater reduction in plasma p-tau181 levels was associated with less clinical decline. Moreover, reduction in plasma p-tau181 levels was correlated with reductions in amyloid PET standard uptake value ratio (SUVR). Additionally, in exploratory analysis of TRAILBLAZER-ALZ trial, donanemab showed a reduction of plasma p-tau217 levels over time. P-tau217 reduction was correlated with changes in amyloid and tau load as assessed by PET (press release July 2021). Recently, the Alzheimer's Association recommended use of BBM for the prescreening or screening of AD subjects in DMT trials before confirmation by PET and CSF testing (38). Table 1 summarizes the current use of plasma biomarkers in ongoing disease-modifying trials.

Blood biomarkers in clinical practice

From lab to clinical practice

Advancing BBM to clinical application may improve disease detection and patient care in the coming years. The scalability of a biomarker, from research to clinical practice, requires the application of 3R's in its development: AccuRacy, Robustness and Regulation. AccuRacy: clinical performance is demonstrated in the population in which the test will be used in clinical practice. Robustness: ability to apply a predefined cut-off prospectively while maintaining clinical performance despite differences in sample handling, operators,

and laboratories. Regulation: demonstration of the analytical and clinical performance of a test before it is brought to the market. As discussed above, the ability of BBM to be used in clinical practice will critically depend on the results of prospective trials demonstrating assay stability and the robustness of predetermined cut points.

Which marker in clinical practice?

Plasma p-tau markers vary in their capability to predict AD according to the specific protein species considered (i.e., p-tau217, p-tau181 and p-tau231), the assay, and thresholds used (10, 12, 24). A recent comparison study showed a relatively high correlation between three p-tau181 assays (Eli Lilly, ADx, Quanterix), one p-tau217 assay (Eli Lilly), and one p-tau231 assay (Gothenburg) (10). Recently it was shown that the p-tau217 assay developed by Janssen exhibits similar performance to the one developed by Eli Lilly (39).

The PrecivityAD™ test is a tandem MS assay used to quantify A β 42 and A β 40 isoforms and determine APOE genotype (32). The PrecivityAD test is intended for use in subjects older than 60 years with MCI or dementia. The PrecivityAD test output is the Amyloid Probability Score (APS), an algorithm derived from the A β 42/40 ratio, the APOE genotype and the patient's age. The APS reflects the likelihood that a patient, on a scale of 0-100, will be amyloid positive on an amyloid PET scan. In two studies, the PARIS study ([ClinicalTrials.gov Identifier: NCT02420756](https://clinicaltrials.gov/ct2/show/study/NCT02420756)) and the Mission AD study ([ClinicalTrials.gov Identifier: NCT02956486](https://clinicaltrials.gov/ct2/show/study/NCT02956486)), involving 686 participants, the APS showed high concordance with amyloid PET status with an area under the curve of 0.88 and an overall accuracy of 81% (6). The cutoff values were established to balance the tradeoff between low frequency of intermediate results and high accuracy (6). Three APS categories were defined: low APS score (0-35) consistent with a low likelihood of amyloid plaques; intermediate APS score (36-57) unable to distinguish between the presence or absence of amyloid plaques; high APS score (58-100), consistent with a high likelihood of amyloid plaques. Importantly, 86% of patients fell into the actionable low or high APS score categories. The PrecivityAD assay is now being used prospectively with predetermined cut points. Interim results from a study underway ([NCT05477056](https://clinicaltrials.gov/ct2/show/study/NCT05477056)) are encouraging in supporting an association of the APS with improved AD diagnostic confidence and clinical decision making. However, at the current time, there are few published data available demonstrating the robustness of the value of the PrecivityAD test in a prospective setting.

The Quest AD-Detect™ plasma biomarker test provides an A β 42/40 ratio and uses a cut off score of 0.160. Ratios \geq 0.160 suggest a lower-than-normal risk of AD and the need to investigate for non-AD causes of cognitive impairment in symptomatic individuals. Ratios $<$ 0.160 are consistent with AD and indicate the need for further evaluation to verify the AD diagnosis. This assay and the threshold score require prospective validation.

NfL is a neuron-specific filament protein that has recently emerged as a biomarker of neuronal injury. The neurofilament proteins levels increase in CSF and in blood, following neuroaxonal damage and regardless of causal pathways. NfL levels increase with aging (40). They are associated with cognitive status and can potentially predict cognitive decline in multiple neurodegenerative conditions such as cerebral small-vessel disease, dementia with Lewy bodies, dementia due to Parkinson disease, frontotemporal dementia, and

multiple sclerosis (41). Although Immunoblot and enzyme-linked immunosorbent assay provided limited sensitivity, third (electrochemiluminescence) and fourth-generation (single-molecule array) assays permit highly sensitive detection in blood samples (41). Recently, the Elecsys Amyloid Plasma Panel (EAPP), developed by Roche, which measures p-tau181 and APOE4 in an individual's blood sample earned breakthrough label from Food and Drug Administration. This is an important step toward more widespread availability of these biomarkers.

What objective?

The different roles of BBM are summarized in Table 2. In clinical practice, BBM may assist in diagnosis and prognosis. In the future, in individuals diagnosed with AD and treated with new DMTs, BBM would be useful to monitor the changes related to the treatment. In this context, BBM might also be essential for the safe and effective use of the treatment. Moreover, BBM would help to identify patients who are most likely to benefit from the treatment as well as individuals at increased risk of serious adverse reactions because of treatment. The same biomarkers can play different roles, especially if different thresholds are used (e.g., lower thresholds to screen patients for additional evaluation as PET or CSF, and higher thresholds to confirm the presence of the disease).

Where?

If it is assumed that BBM are demonstrated to be stable and robust in a prospective setting, BBM could be used in primary care. A possibility is to utilize BBM in primary care to identify the patients with cognitive impairment that need to be referred for further cognitive evaluation by specialists in memory disorders. For this reason, a biomarker of neurodegeneration or one not specific to AD, such as the NfL, may represent a good candidate to be implemented in primary care. However, the robustness of this strategy in primary care needs to be determined. Another possibility is to recommend that trained primary care physicians can use AD specific BBM, including A β 42/A β 40 ratio and p-tau measures, for diagnosis and management without referral to a memory disorder specialist. The following arguments support this option:

- Timely detection of cognitive impairment, and accurate etiologic diagnosis are notoriously poor in primary care clinics due to several challenges including the lack of simple, accurate and accessible tests and the scarcity of clinicians trained in cognitive disorders. BBM are reaching performance levels similar to PET or CSF and are now available with improved versions coming soon.
- In many areas, most AD care is still provided by non-memory specialists, even more so in regions with low access to specialty care.
- If AD-specific treatments (e.g., anti-amyloid) are approved and become widely accessible, the need for detection and accurate diagnosis will be amplified.

In the last decade, digital technologies (e.g., data collection application, sensors, wearable devices) that allow remote assessment and collection of real-world data on cognitive outcomes have attracted great attention and interest in the field of cognitive diseases (42, 43). The World Health Organization (WHO) ICOPE step 1 is a simple tool, which can

be used to detect an impairment in one or more domain(s) of the intrinsic capacity (i.e., cognition, mood, nutrition, mobility, sensory). In France, a mobile app and a conversational robot, that enable self-assessment, have been developed to promote ICOPE step 1 use (44). These tools are now extended internationally. In ICOPE step 1, memory is assessed by a subjective question, an orientation task, and a free recall of 3 words. In a population of about 18000 individuals with a mean age of 76 years old, it was shown that 60% presented either a subjective or an objective complaint (45). In 50 % of cases these alterations were persistent after 6 months (not published data). In primary care, digital biomarkers can be used to preselect individuals at risk of cognitive decline that could benefit from BBM assessment. In Mexico, the ICOPE step 1 is followed by the Davos Alzheimer's Collaborative program to explore the addition of a BBM in a real-world setting, but under a research protocol, to assess facilitators and barriers to such an approach from a health system viewpoint.

In the specialty care setting, the goal of the BBM is the etiologic diagnosis of dementia and MCI. For this purpose, A β 42/40 ratio, p-tau217 and multiplex assays would be performed as part of (not replacement for) a amyloid, tau, neurodegeneration (AT(N)) classification system (46). A great benefit of a BBM in specialty care would be to reduce unnecessary confirmatory procedures as PET and CSF biomarkers. In this way, the specialty care pathway would be more streamlined and made more efficient.

When?

In the first step, BBM would be collected only in subjects with an objective cognitive impairment (MCI and dementia). BBM should be ordered as part of the initial diagnostic workup, unless other (non-AD) obvious and untreated causes of cognitive decline are present. Since the first signs of cognitive decline and in case of suspicion of AD, BBM can be conducted in parallel with history, collection of risk factors and medications, cognitive and behavioral status exam, medical/neurologic exam and brain structural imaging. BBM will also have great utility in patients who have contraindications to lumbar puncture or in those reluctant to lumbar puncture.

How?

For diagnosis, BBM could be useful when high confidence is required for AD or non-AD pathologies. For the diagnosis of typical amnesic AD, BBM, especially p-tau217, could be sufficient or could be part of a multiplex assessment with A β 42/40 and APOE genotype. For atypical presentations (non-amnesic, early onset, rapidly progressive, logopenic aphasia, posterior cortical atrophy), BBM for AD are likely sufficient if positive; but if negative, and AD remains in the differential diagnosis with at least moderate likelihood, suggest the need for CSF biomarkers or FDG-PET. In case of discordance of amyloid and tau BBM or intermediate results, CSF may resolve the uncertainty (or PET if CSF is contraindicated, Table 3). The results of BBM should be integrated with all other clinical features in a patient-centric manner. Even in the case of AD blood biomarkers positivity, clinician should not overlook other potentially contributing factors and conditions (e.g., depression, obstructive sleep apnea, iatrogenic etiologies).

Advantages of blood-based markers in health care systems

Previous work has suggested that health systems are ill-prepared to cope with the expected workload of new treatments: the prevalent caseload is large and the number of dementia specialists is not sufficient. Thus, the wait time for comprehensive cognitive evaluations is likely to increase (47). One strategy to facilitate the diagnosis of AD and potentially to facilitate access to new treatments would therefore be to optimize triage in primary care. A recent study estimated impact of BBM on wait times and cost with a simulation model in several scenarios (Mini Mental State Examination MMSE only, BBM only, combination of both) (48). The combination of MMSE and BBM could strongly decrease the demand for specialist visits, the average wait time to complete the diagnosis process and the average annual cost by \$400 to \$700 million for the U.S. health care system. Carefully and appropriately implementing of a similar strategy, combining MMSE (or a similar brief validated cognitive test e.g., MoCA) and BBM in primary care settings, could also potentially improve the diagnosis accuracy; while utilizing a BBM only strategy would lead to many inappropriate referrals of cognitively unimpaired individuals to specialists in memory disorders.

The BBM have the potential for being more accessible and cheaper (less invasive, time-intensive, costly, infrastructure dependent, and resource requiring – can be done in the community setting) than CSF and PET biomarkers. They could improve detection, diagnostic accuracy, patient-centered autonomy and empowerment, and better overall care. The use of BBM in clinical practice would be a great opportunity for referral to research (e.g., use of legacy samples and screening for clinical trials). Programs, such as the Davos Alzheimer's Collaborative are exploring this workflow in a real-world setting, but under a research protocol, to assess facilitators and barriers to such an approach from a health system viewpoint.

Potential risks of blood-based markers use in clinical practice

Among the risks, discussed above, there are lack of stability of BBM assay results when used prospectively, lack of robustness of cut points determining normal/abnormal results and the limited assessment of these cut points/ranges in real-world populations with ethnocultural diversity, advanced age (oldest old), and comorbidities. Attention to strict quality control measures over time is essential to mitigate part of this risk. These issues must also be resolved prior to implementation in clinical practice. In the absence of consensus guidelines regulating the use of BBM, there is an important risk of misuse and misinterpretation of the results. A common example of misuse is the use of BBM in cognitive unimpaired individuals, especially without associated cognitive testing, when the BBM has not been validated for this application. An important risk is the underdiagnosis of non-AD pathologies and other conditions potentially contributing to cognitive disorder in patients with negative BBM. Indeed, unexperienced users could interpret negative values as supporting a lack of need for further assessment (for other causes and contributors) and to refer patients to specialists in memory disorders. On the contrary, positive values could lead unexperienced BBM users to make a diagnosis of AD while disregarding other likely causes (e.g., co-existing vascular cognitive impairment or Lewy Body Disease) and potentially contributing conditions.

Ethical considerations

Being minimally invasive, cost-effective, and globally scalable, BBM represent an opportunity to improve equitable access to medical care and research and to limit social disparities in healthcare. Patients throughout the United States and other regions of the world do not have easy and uniform access to specialized care centers, which are often the locations for PET imaging or specialists that can perform lumbar punctures for CSF analysis. These strategies are largely concentrated in major metropolitan areas, which present substantial geographic access barriers for many patients. Research has demonstrated that PET scan usage is associated with challenges in reaching healthcare equity and inclusion (49). Studies indicate that learning biomarker results is safe, and that patients understand their results correctly (50, 51). However, these data came from motivated populations of research participants (52, 53) and the disclosure of results was conducted by experienced providers; thus, they may not be generalizable to other patient populations or settings. To ensure a safe use of BBM in clinical practice, a process of informed consent, including the provision of adequate information before testing, is appropriate. The clinician should ensure that patients and care partners understand what BBM results can and cannot inform and that the individual desires to be aware of this information. The disclosure of BBM results is likely to have a major impact for some patients and care partners. For this reason, patients and care partners may need emotional support after the disclosure of BBM results. Finally, to be diagnosed with AD remains a source of stigma and discrimination, and this can lead some individuals to delay or forego care. Stigma must be addressed to avoid its effects on patients and care partners and to increase the likelihood that we can realize the potential of BBM.

Key points about blood-based markers use to aid in diagnosis of AD in clinical practice

1. Trained primary care physicians can use BBM in AD diagnosis without referral to a specialist in memory disorders.
2. BBM such as p-tau measures and A β 42/40 ratio can be used alone or in combination to aid in AD diagnosis.
3. BBM should be used in combination with clinical and cognitive evaluation. The interpretation of BBM needs to be integrated into the standard process for AD diagnosis (history, cognitive evaluation, structural brain imaging).
4. In primary care, soon (when prospective studies are completed) BBM could be used after a digital cognitive evaluation to assess subjects with objective and persistent cognitive impairment. Prospective studies are needed to validate this use.

What are the next steps?

Over the past decade substantial progress in AD biomarkers research has been made, and several actions are underway to promote BBM implementation in clinical practice. With few exceptions, almost all current data on BBM performance comes from retrospective studies on batches of samples collected in research settings and in populations with limited ethnocultural diversity and comorbidities. Therefore, it will be critical to demonstrate the

effectiveness of BBM when used prospectively and longitudinally with preset cut points in more diverse real-world clinical populations. A close cooperation among pharmaceutical and diagnostic industry partners, academic institutions, regulatory bodies, and patient advocacy associations is needed. It will be useful to continue the Global Biomarker Standardization Consortium, to explore the development of a regulatory focus group with international participation, and to increase efforts to understand medical specialty and primary care needs and constraints.

The advent of BBM with reasonable sensitivity and specificity would lead to substantial changes in health care delivery. We expect to see an important surge in the number of patients seeking potential assessments. There is an urgent need for improved training and proficiency for primary care providers and specialists in memory disorders, as well as a need for development of clinical algorithms to better triage patients from primary care to specialists in memory disorders. Appropriate use criteria in the different settings should be drafted, clearly by defining for each biomarker the clinical purpose, the target population, the potential benefits and risks, and guidance regarding results interpretation and disclosure. There is a crucial need for standardization of operating procedures, to minimize and control pre-analytical, analytical, and post-analytical sources of variability. Work to understand what is needed for reimbursement of BBM is necessary also.

The development of AD BBM is expected to advance into other proteinopathies (TDP-43 pathologies, non-AD tauopathies, alpha-synucleinopathies). This could offer to the clinicians in the future high performance diagnostic tools in the differential diagnosis of cognitive impairment. Further studies with fixed biomarker cut-off set before the study starts need to demonstrate the impact of BBM use on clinical outcomes (e.g., time to diagnosis, diagnosis of AD at an early stage, inclusion in clinical trials, cost/effectiveness, etc).

Large epidemiological prospective studies must be implemented to better understand lifetime development of BBM as well as their interaction with age and comorbidities. Given the high incidence of cognitive decline in people older than 80 years old, more data are needed to interpret such biomarkers in the oldest old. Prospective studies that enroll ethnically and racially diverse individuals are needed. Large prospective studies, in clinical practice must be implemented now to be ready when new therapies are widely accessible. Clear evidence of clinical utility will be required to justify reimbursement.

This is a pivotal moment in the implementation of BBM in clinical trials. Their widespread use should quickly bring dramatic improvements to the conduct of trials, facilitating recruitment, reducing time and costs, and increasing the prospect of effective treatments for the disease. The time needed to implement BBM in clinical routine will inevitably depend on the results of the ongoing clinical trials with DMTs. This process will be accelerated by positivity of one or more clinical trial.

Conflict of interest:

The Task Force was partially funded by registration fees from industrial participants. These corporations placed no restrictions on this work. D. Angioni is an investigator in clinical trials sponsored by Roche, Alektor, Jansen, Alzheon, Otsuka, Novonordisk, UCB Pharma, Medesis Pharma and Eisai. No direct personal benefit is to be declared. J. Delrieu has received payment/honoraria from Biogen (presentation for Biogen in 2021); has

participated on a Data Safety Monitoring Board or Advisory Board for French board for Roche in 2020-2022. H. Fillit is an unpaid consultant to Lilly and Biogen. He have received consulting fees from Alektor, Samus Therapeutics, TauRx, Pinteon, Otsuka/Lundbeck, and Genentech during the past 3 years. P. Aisen reports grants from Janssen, NIA, FNIH, Alzheimer's Association, Eisai and consultant fees from Merck, Biogen, Roche, ImmunoBrain Checkpoint, Abbvie, Rainbow Medical, Shionogi, Vigil Neuroscience, Inc, Hengrui USA, outside the submitted work. J. Cummings reports grants from: NIGMS grant P20GM109025; NINDS grant U01NS093334; NIA grant R01AG053798; NIA grant P20AG068053; NIA grant P30AG072959; NIA grant R35AG71476; Alzheimer's Disease Drug Discovery Foundation (ADDF); Ted and Maria Quirk Endowment for the Pam Quirk Brain Health and Biomarker Laboratory; and the Joy Chambers-Grundy Endowment during the conduct of the study. J. Cummings receives consultant fees from AB Science, Acadia, Alkahest, AlphaCognition, ALZPathFinder, Annovis, AriBio, Artery, Avanir, Biogen, Biosplice, Cassava, Cerevel, Clinilabs, Cortexyme, Diadem, EIP Pharma, Eisai, GatehouseBio, GemVax, Genentech, Green Valley, Grifols, Janssen, Karuna, Lexeo, Lilly, Lundbeck, LSP, Merck, NervGen, Novo Nordisk, Oligomerix, Ono, Otsuka, PharmacotrophiX, PRODEO, Prothena, ReMYND, Renew, Resverlogix, Roche, Signant Health, Suven, Unlearn AI, Vaxxinity, VigilNeuro pharmaceutical, assessment, and investment companies, outside the submitted work. J. Cummings is the Chief Scientific Officer for CNS Innovations, LLC. J.R. Sims is an employee of Eli Lilly and Company: salary and minor stockholder. J. Braunstein is Employee of C2N Diagnostics. M. Sabbagh. Ownership interest (stock or stock options): NeuroTau, Optimal Cognitive Health Company, uMethod Health, Versanum, Athira, TransDermix, Seq BioMarque, NeuroReserve, Cortexyme/Quince Therapeutics; Consulting: Alzheon, Roche-Genentech, Eisai, KeifeRx, Lilly, Synaptogenix, NeuroTherapia, T3D, Signant Health, Novo Nordisk; Royalties: Humanix; Board of Director: Cortexyme/Quince Therapeutics, EIP Pharma. T. Bittner is a full time employee of F.Hoffmann-LaRoche AG and Genentech. M. Pontecorvo is an employee of Avid Radiopharmaceuticals, a wholly owned affiliate of Eli Lilly and Corporation and is a minor stockholder in Lilly. S. Bozeat is a full-time employee of Hoffman-La Roche and has stock options in the company. J.L. Dage is an inventor on patents or patent applications of Eli Lilly and Company relating to the assays, methods, reagents and/or compositions of matter related to measurement of P-tau217. Dr. Dage has served as a consultant for Genotix Biotechnologies Inc, Gates Ventures, Karuna Therapeutics, AlzPath Inc, Cognito, and received research support from ADx Neurosciences, Roche Diagnostics and Eli Lilly and Company in the past two years. E. Largent reports grants from National Institute on Aging, grants from Greenwall Foundation, outside the submitted work. S. Mattke reports personal fees from Biogen, Eisai, Roche/Genentech, Novartis, C2N, outside the submitted work. D.R. Willis is the principal investigator on a grant funded by the Devos Alzheimer's Collaborative. This grant provides travel to the DAC in-person meetings. D.R. Willis did not participate or consume any food provided by industry partners. The DAC includes in-kind contributions from industry partners for patients to complete digital cognitive assessments (no charge for software or hardware) and blood-based biomarkers testing (sample processing). Contractual relationships between her university and each industry partner offering those in-kind resources (Linus Health and C2N Diagnostics respectively) prevent any undue influence over any scholarly work or dissemination related to the project. At the time of submission of this paper, they did not have sent any specimen to C2N for in-kind processing. To date no financial benefit has been received from C2N. Alireza Atri has received honoraria or support for consulting; participating in independent data safety monitoring boards; providing educational lectures, programs, and materials; or serving on advisory boards for AbbVie, Acadia, Allergan, the Alzheimer's Association, Axovant, AZ Therapies, Biogen, Eisai, Grifols, Harvard Medical School Graduate Continuing Education, JOMDD, Lundbeck, Merck, Roche/Genentech, Novo Nordisk, Qynapse, Sunovion, Suven, and Synexus. Dr. Atri receives book royalties from Oxford University Press for a medical book on dementia. Dr. Atri currently receives institutional research grant/contract funding from NIA/NIH 1P30AG072980, AZ DHS CTR040636, Washington University St Louis, and Gates Ventures. Dr. Atri's institution receives/received funding for clinical trial grants, contracts and projects from government, consortia, foundations and companies for which he serves/served as contracted site-PI. Washington University and R.J. Bateman have equity ownership interest in C2N Diagnostics and receive income based on technology (blood plasma assay) licensed by Washington University to Diagnostics. R.J. Bateman receives income from C2N Diagnostics for serving on the scientific advisory board. Washington University, with R.J. Bateman as co-inventor, has submitted the US nonprovisional patent application "Plasma Based Methods for Determining A-Beta Amyloidosis." R.J. Bateman has received honoraria as a speaker/consultant/advisory board member from Amgen, Eisai, Hoffman-LaRoche, and Janssen; and reimbursement of travel expenses from, Hoffman-La Roche and Janssen. As an investigator, P.J. Ousset has received research funding from the following companies: Acadia Pharmaceuticals, Alektor Inc., Alzheon Inc., Araclon Biotech S.L., AstraZeneca, Biogen, Avanir Pharmaceuticals, Cortexyme Inc., Eisai Inc., Eli Lilly, Genentech, Inc., Hoffmann-La Roche, Janssen Research & Development, Merck Sharp & Dohme, Novartis Pharmaceuticals, Novo Nordisk, Shanghai Green Valley Pharmaceutical Co., TauRx Therapeutics Ltd, UCB Biopharma. No direct personal benefit is to be declared. B. Vellas is an investigator in clinical trials sponsored by Biogen, Lilly, Roche, Eisai Pharmaceuticals and the Toulouse University Hospital. He has served as SAB member for Biogen, Alzheon, Green Valley, Norvo Nordisk, Longeveron, but received no personal compensation. He has served as consultant and/or SAB member for Roche, Lilly, Eisai, TauX with personal compensation. He is member of the Editorial Board of JPAD with no personal compensation; and did not have a role in the editorial process/review for this manuscript. M. Weiner reports serves on Editorial Boards for Alzheimer's & Dementia, MRI and TMRI. He has served on Advisory Boards for Acumen Pharmaceutical, ADNI, Alzheon, Inc., Biogen, Brain Health Registry, Cerecin, Dolby Family Ventures, Eli Lilly, Merck Sharp & Dohme Corp., National Institute on Aging (NIA), Nestle/Nestec, PCORI/PPRN, Roche, University of Southern California (USC), NervGen. He has provided consulting to Baird Equity Capital, BioClinica, Cerecin, Inc., Cytex, Dolby

Family Ventures, Duke University, Eisai, FUJIFILM-Toyama Chemical (Japan), Garfield Weston, Genentech, Guidepoint Global, Indiana University, Japanese Organization for Medical Device Development, Inc. (JOMDD), Medscape, Nestle/Nestec, NIH, Peerview Internal Medicine, Roche, T3D Therapeutics, University of Southern California (USC), WebMD, and Vida Ventures. He has acted as a speaker/lecturer to The Buck Institute for Research on Aging; China Association for Alzheimer's Disease (CAAD); Japan Society for Dementia Research; and Korean Dementia Society. He holds stock options with Alzheon, Inc., Alzeca, and Anven. The following entities have provided funding for academic travel; University of Southern California (USC), NervGen, ASFNR, and CTAD Congress. Mr. Weiner receives support for his research from the following funding sources: National Institutes of Health (NIH), Department of Defense (DOD), Patient-Centered Outcomes Research Institute (PCORI), California Department of Public Health (CDPH), University of Michigan, Siemens, Biogen, Hillblom Foundation, Alzheimer's Association, The State of California, Johnson & Johnson, Kevin and Connie Shanahan, GE, VUmc, Australian Catholic University (HBI-BHR), The Stroke Foundation, and the Veterans Administration.

References

1. Beach TG, Monsell SE, Phillips LE, Kukull W. Accuracy of the clinical diagnosis of Alzheimer disease at National Institute on Aging Alzheimer Disease Centers, 2005-2010. *J Neuropathol Exp Neurol.* 2012 Apr;71(4):266–73. [PubMed: 22437338]
2. Bradford A, Kunik ME, Schulz P, Williams SP, Singh H. Missed and delayed diagnosis of dementia in primary care: prevalence and contributing factors. *Alzheimer Dis Assoc Disord.* 2009 Dec;23(4):306–14. [PubMed: 19568149]
3. Hansson O. Biomarkers for neurodegenerative diseases. *Nat Med.* 2021 Jun;27(6):954–63. [PubMed: 34083813]
4. Budd Haeberlein S, Aisen PS, Barkhof F, Chalkias S, Chen T, Cohen S, et al. Two Randomized Phase 3 Studies of Aducanumab in Early Alzheimer's Disease. *J Prev Alzheimers Dis.* 2022;9(2):197–210. [PubMed: 35542991]
5. West T, Kirmess KM, Meyer MR, Holubasch MS, Knapik SS, Hu Y, et al. A blood-based diagnostic test incorporating plasma A β 42/40 ratio, ApoE proteotype, and age accurately identifies brain amyloid status: findings from a multi cohort validity analysis. *Mol Neurodegener.* 2021 May 1;16(1):30. [PubMed: 33933117]
6. Hu Y, Kirmess KM, Meyer MR, Rabinovici GD, Gatsonis C, Siegel BA, et al. Assessment of a Plasma Amyloid Probability Score to Estimate Amyloid Positron Emission Tomography Findings Among Adults With Cognitive Impairment. *JAMA Netw Open.* 2022 Apr 1;5(4):e228392. [PubMed: 35446396]
7. Schindler SE, Bollinger JG, Ovod V, Mawuenyega KG, Li Y, Gordon BA, et al. High-precision plasma β -amyloid 42/40 predicts current and future brain amyloidosis. *Neurology.* 2019 22;93(17):e1647–59. [PubMed: 31371569]
8. Janelidze S, Teunissen CE, Zetterberg H, Allué JA, Sarasa L, Eichenlaub U, et al. Head-to-Head Comparison of 8 Plasma Amyloid- β 42/40 Assays in Alzheimer Disease. *JAMA Neurol.* 2021 Nov 1;78(11):1375–82. [PubMed: 34542571]
9. Mielke MM, Frank RD, Dage JL, Jeromin A, Ashton NJ, Blennow K, et al. Comparison of Plasma Phosphorylated Tau Species With Amyloid and Tau Positron Emission Tomography Neurodegeneration, Vascular Pathology, and Cognitive Outcomes. *JAMA Neurol.* 2021 Sep 1;78(9):1108–17. [PubMed: 34309632]
10. Bayoumy S, Verberk IMW, den Dulk B, Hussainali Z, Zwan M, van der Flier WM, et al. Clinical and analytical comparison of six Simoa assays for plasma P-tau isoforms P-tau181, P-tau217, and P-tau231. *Alzheimers Res Ther.* 2021 Dec 4;13(1):198. [PubMed: 34863295]
11. Barthélemy NR, Horie K, Sato C, Bateman RJ. Blood plasma phosphorylatedtau isoforms track CNS change in Alzheimer's disease. *J Exp Med.* 2020 Nov 2;217(11):e20200861. [PubMed: 32725127]
12. Palmqvist S, Janelidze S, Quiroz YT, Zetterberg H, Lopera F, Stomrud E, et al. Discriminative Accuracy of Plasma Phospho-tau217 for Alzheimer Disease vs Other Neurodegenerative Disorders. *JAMA.* 2020 Aug 25;324(8):772–81. [PubMed: 32722745]
13. Wennström M, Janelidze S, Nilsson KPR, Netherlands Brain Bank, Serrano GE, Beach TG, et al. Cellular localization of p-tau217 in brain and its association with p-tau217 plasma levels. *Acta Neuropathol Commun.* 2022 Jan 6;10(1):3. [PubMed: 34991721]

14. Janelidze S, Palmqvist S, Leuzy A, Stomrud E, Verberk IMW, Zetterberg H, et al. Detecting amyloid positivity in early Alzheimer's disease using combinations of plasma A β 42/A β 40 and p-tau. *Alzheimers Dement*. 2022 Feb;18(2):283–93. [PubMed: 34151519]
15. Barthélemy NR, Li Y, Joseph-Mathurin N, Gordon BA, Hassenstab J, Benzinger TLS, et al. A soluble phosphorylated tau signature links tau, amyloid and the evolution of stages of dominantly inherited Alzheimer's disease. *Nat Med*. 2020 Mar;26(3):398–407. [PubMed: 32161412]
16. Palmqvist S, Tideman P, Cullen N, Zetterberg H, Blennow K, Alzheimer's Disease Neuroimaging Initiative, et al. Prediction of future Alzheimer's disease dementia using plasma phospho-tau combined with other accessible measures. *Nat Med*. 2021 Jun;27(6):1034–42. [PubMed: 34031605]
17. Benedet AL, Brum WS, Hansson O, Alzheimer's Disease Neuroimaging Initiative, Karikari TK, Zimmer ER, et al. The accuracy and robustness of plasma biomarker models for amyloid PET positivity. *Alzheimers Res Ther*. 2022 Feb 7;14(1):26. [PubMed: 35130933]
18. Cullen NC, Janelidze S, Mattsson-Carlgen N, Palmqvist S, Bittner T, Suridjan I, et al. Test-retest variability of plasma biomarkers in Alzheimer's disease and its effects on clinical prediction models. *Alzheimers Dement*. 2022 Jun 14;
19. Tosun D, Veitch D, Aisen P, Jack CR, Jagust WJ, Petersen RC, et al. Detection of β -amyloid positivity in Alzheimer's Disease Neuroimaging Initiative participants with demographics, cognition, MRI and plasma biomarkers. *Brain Commun*. 2021;3(2):fcab008. [PubMed: 33842885]
20. Nakamura A, Kaneko N, Villemagne VL, Kato T, Doecke J, Doré V, et al. High performance plasma amyloid- β biomarkers for Alzheimer's disease. *Nature*. 2018 08;554(7691):249–54. [PubMed: 29420472]
21. Li Y, Schindler SE, Bollinger JG, Ovod V, Mawuenyega KG, Weiner MW, et al. Validation of Plasma Amyloid- β 42/40 for Detecting Alzheimer Disease Amyloid Plaques. *Neurology*. 2022 Feb 15;98(7):e688–99. [PubMed: 34906975]
22. Zicha S, Bateman RJ, Shaw LM, Zetterberg H, Bannon AW, Horton WA, et al. Comparative analytical performance of multiple plasma A β 42 and A β 40 assays and their ability to predict positron emission tomography amyloid positivity. *Alzheimers Dement*. 2022 Jul 12;
23. Meyer MR, Kirmess KM, Wentz-Roth TL, Irvin F, Holubasch MS, Verghese PB, et al. Plasma ratio of phospho- to non-phospho-tau at threonine 217 identifies brain amyloid burden as measured by quantitative amyloid PET in the PARIS sub-study of IDEAS. AAIC Scientific Session, July 31, 2022.
24. Mielke MM, Dage JL, Frank RD, Algeciras-Schimmich A, Knopman DS, Lowe VJ, et al. Performance of plasma phosphorylated tau 181 and 217 in the community. *Nat Med*. 2022 May 26;
25. Tsiknia AA, Edland SD, Sundermann EE, Reas ET, Brewer JB, Galasko D, et al. Sex differences in plasma p-tau181 associations with Alzheimer's disease biomarkers, cognitive decline, and clinical progression. *Mol Psychiatry*. 2022 Jun 29;
26. Schindler SE, Karikari TK, Ashton NJ, Henson RL, Yarasheski KE, West T, et al. Effect of Race on Prediction of Brain Amyloidosis by Plasma A β 42/A β 40, Phosphorylated Tau, and Neurofilament Light. *Neurology*. 2022 Jul 19;99(3):e245–57. [PubMed: 35450967]
27. Granot-HersHKovitz E, Tarraf W, Kurniansyah N, Daviglus M, Isasi CR, Kaplan R, et al. APOE alleles' association with cognitive function differs across Hispanic/Latino groups and genetic ancestry in the study of Latinos-investigation of neurocognitive aging (HCHS/SOL). *Alzheimers Dement*. 2021 Mar;17(3):466–74. [PubMed: 33155766]
28. Crean S, Ward A, Mercaldi CJ, Collins JM, Cook MN, Baker NL, et al. Apolipoprotein E ϵ 4 prevalence in Alzheimer's disease patients varies across global populations: a systematic literature review and meta-analysis. *Dement Geriatr Cogn Disord*. 2011;31(1):20–30. [PubMed: 21124030]
29. Cummings J. The Role of Biomarkers in Alzheimer's Disease Drug Development. *Adv Exp Med Biol*. 2019;1118:29–61. [PubMed: 30747416]
30. Sperling RA, Johnson KA, Zhou J, Irizarry MC, Dhadda S, Kramer LD, Li D, Li P, Swanson CJ, West T, Yarasheski KE, Kirmess KM, Meyer MR, Braunstein JB, Rissman RA, Rafii M, Grill J, Raman R, Sethuraman G, Donohue MC, Langford O, Aisen PS. on behalf of the AHEAD 3 45

- Study Team. Introduction of plasma biomarker screening for the AHEAD 3 45 Study. Clinical Trials on Alzheimer's Disease Taskforce meeting, 31 May 2022.
31. Sims JR. Plasma Biomarkers in Screening: Lilly Trials. Clinical Trials on Alzheimer's Disease Taskforce meeting, 31 May 2022.
 32. Kirmess KM, Meyer MR, Holubasch MS, Knapik SS, Hu Y, Jackson EN, et al. The PrecivityADTM test: Accurate and reliable LC-MS/MS assays for quantifying plasma amyloid beta 40 and 42 and apolipoprotein E proteotype for the assessment of brain amyloidosis. *Clin Chim Acta*. 2021 Aug;519:267–75. [PubMed: 34015303]
 33. Bittner T. Blood-based prescreening for SKYLINE. Clinical Trials on Alzheimer's Disease Taskforce meeting, 31 May 2022.
 34. Leuzy A, Mattsson-Carlgrén N, Palmqvist S, Janelidze S, Dage JL, Hansson O. Blood-based biomarkers for Alzheimer's disease. *EMBO Mol Med*. 2022 Jan 11;14(1):e14408. [PubMed: 34859598]
 35. Swanson CJ, Zhang Y, Dhadda S, Wang J, Kaplow J, Lai RYK, et al. A randomized, double-blind, phase 2b proof-of-concept clinical trial in early Alzheimer's disease with lecanemab, an anti-A β protofibril antibody. *Alzheimers Res Ther*. 2021 Apr 17;13(1):80. [PubMed: 33865446]
 36. Salloway S, Farlow M, McDade E, Clifford DB, Wang G, Llibre-Guerra JJ, et al. A trial of gantenerumab or solanezumab in dominantly inherited Alzheimer's disease. *Nat Med*. 2021 Jul;27(7):1187–96. [PubMed: 34155411]
 37. Mattsson-Carlgrén N, Janelidze S, Palmqvist S, Cullen N, Svenningsson AL, Strandberg O, et al. Longitudinal plasma p-tau217 is increased in early stages of Alzheimer's disease. *Brain*. 2020 Dec 5;143(11):3234–41. [PubMed: 33068398]
 38. Hansson O, Edelmayer RM, Boxer AL, Carrillo MC, Mielke MM, Rabinovici GD, et al. The Alzheimer's Association appropriate use recommendations for blood biomarkers in Alzheimer's disease. *Alzheimers Dement*. 2022 Jul 31;
 39. Groot C, Cicognola C, Bali D, Triana-Baltzer G, Dage JL, Pontecorvo MJ, et al. Correction: Diagnostic and prognostic performance to detect Alzheimer's disease and clinical progression of a novel assay for plasma p-tau217. *Alzheimers Res Ther*. 2022 Jun 13;14(1):82. [PubMed: 35698175]
 40. Khalil M, Pirpamer L, Hofer E, Voortman MM, Barro C, Leppert D, et al. Serum neurofilament light levels in normal aging and their association with morphologic brain changes. *Nat Commun*. 2020 Feb 10;11(1):812. [PubMed: 32041951]
 41. Khalil M, Teunissen CE, Otto M, Piehl F, Sormani MP, Gatteringer T, et al. Neurofilaments as biomarkers in neurological disorders. *Nat Rev Neurol*. 2018 Oct;14(10):577–89. [PubMed: 30171200]
 42. Kaye J, Aisen P, Amariglio R, Au R, Ballard C, Carrillo M, et al. Using Digital Tools to Advance Alzheimer's Drug Trials During a Pandemic: The EU/US CTAD Task Force. *J Prev Alzheimers Dis*. 2021;8(4):513–9. [PubMed: 34585227]
 43. Gold M, Amatniek J, Carrillo MC, Cedarbaum JM, Hendrix JA, Miller BB, et al. Digital technologies as biomarkers, clinical outcomes assessment, and recruitment tools in Alzheimer's disease clinical trials. *Alzheimers Dement (N Y)*. 2018;4:234–42. [PubMed: 29955666]
 44. Piau A, Steinmeyer Z, Cesari M, Kornfeld J, Beattie Z, Kaye J, et al. Intrinsic Capacity Monitoring by Digital Biomarkers in Integrated Care for Older People (ICOPE). *J Frailty Aging*. 2021;10(2):132–8. [PubMed: 33575701]
 45. Tavassoli N, de Souto Barreto P, Berbon C, Mathieu C, de Kerimel J, Lafont C, et al. Implementation of the WHO integrated care for older people (ICOPE) programme in clinical practice: a prospective study. *The Lancet Healthy Longevity*. 2022 Jun;3(6):e394–404. [PubMed: 36098317]
 46. Jack CR, Bennett DA, Blennow K, Carrillo MC, Dunn B, Haeberlein SB, et al. NIA-AA Research Framework: Toward a biological definition of Alzheimer's disease. *Alzheimers Dement*. 2018;14(4):535–62. [PubMed: 29653606]
 47. Hlavka JP, Mattke S, Liu JL. Assessing the Preparedness of the Health Care System Infrastructure in Six European Countries for an Alzheimer's Treatment. *Rand Health Q*. 2019 May;8(3):2. [PubMed: 31205802]

48. Mattke S, Cho SK, Bittner T, Hlávka J, Hanson M. Blood-based biomarkers for Alzheimer's pathology and the diagnostic process for a disease-modifying treatment: Projecting the impact on the cost and wait times. *Alzheimers Dement (Amst)*. 2020;12(1):e12081. [PubMed: 32832590]
49. Onega T, Tosteson TD, Wang Q, Hillner BE, Song Y, Siegel BA, et al. Geographic and sociodemographic disparities in PET use by Medicare beneficiaries with cancer. *J Am Coll Radiol*. 2012 Sep;9(9):635–42. [PubMed: 22954545]
50. Mozersky J, Sankar P, Harkins K, Hachey S, Karlawish J. Comprehension of an Elevated Amyloid Positron Emission Tomography Biomarker Result by Cognitively Normal Older Adults. *JAMA Neurol*. 2018 Jan 1;75(1):44–50. [PubMed: 29059270]
51. Erickson CM, Chin NA, Johnson SC, Gleason CE, Clark LR. Disclosure of preclinical Alzheimer's disease biomarker results in research and clinical settings: Why, how, and what we still need to know. *Alzheimers Dement (Amst)*. 2021;13(1):e12150. [PubMed: 33665341]
52. Lingler JH, Sereika SM, Butters MA, Cohen AD, Klunk WE, Knox ML, et al. A randomized controlled trial of amyloid positron emission tomography results disclosure in mild cognitive impairment. *Alzheimers Dement*. 2020 Sep;16(9):1330–7. [PubMed: 32588971]
53. Largent EA, Harkins K, van Dyck CH, Hachey S, Sankar P, Karlawish J. Cognitively unimpaired adults' reactions to disclosure of amyloid PET scan results. *PLoS One*. 2020;15(2):e0229137. [PubMed: 32053667]

Table 1.

Some examples of blood-based markers use in clinical trials

Study	Clinicaltrial.gov identifier	Phase	Population	Therapy	Blood biomarker	Role
AHEAD 3-45	NCT04468659	III	Preclinical AD	Lecanemab	Aβ42/40 ratio	Prescreening
AUTONOMY	NCT04619420	II	Early symptomatic AD	JNJ-63733657	p-tau217	Prescreening
BAN2401-G000-201 Core and Open Label Extension	NCT01767311	II	Early symptomatic AD	Lecanemab	Aβ42/40 ratio p-tau 181	Pharmacodynamic
DIAN-TU	NCT04623242	II/III	Preclinical and early symptomatic AD	Gantenerumab	Aβ42/40 ratio p-tau181	Pharmacodynamic
EMERGE	NCT02484547	III	Early symptomatic AD	Aducanumab	p-tau 181	Pharmacodynamic
ENGAGE	NCT02477800	III	Early symptomatic AD	Aducanumab	p-tau 181	Pharmacodynamic
EVOKE	NCT04777396	III	Early symptomatic AD	Semaglutide	p-tau181, NfL, GFAB	Pharmacodynamic
EVOKE-PLUS	NCT04777409	III	Early symptomatic AD	Semaglutide	p-tau181, NfL, GFAB	Pharmacodynamic
INVOLVE-2	NCT04592874	II	Early symptomatic AD	AL002	PredixyAD™ (algorithm derived from Aβ42/40 ratio, APOE genotype and age)	Prescreening
PROSPECT-ALZ	NCT05063539	II	Early symptomatic AD	LX3372689	p-tau217	Prescreening
TRAILBLAZER-ALZ 2	NCT044437511	III	Early symptomatic AD	Donanemab	p-tau181 p-tau217	Exploratory endpoint
TRAILBLAZER-ALZ 3	NCT05026866	III	Preclinical AD	Donanemab	p-tau217	Exploratory endpoint

Aβ, Amyloid-beta; AD, Alzheimer's Disease; APOE, Apolipoprotein E; NfL, Neurofilament Light chain; P-tau, phosphorylated tau.

Table 2.

Potential roles for blood-based markers use in clinical practice

Potential role		Where?
Risk/probability	To detect persons at risk for a disease or condition	Primary care
Diagnosis	To confirm the presence of a disease/condition	Primary care and specialists in memory disorders
Prognosis	To determine risk of worsening and/or progression from MCI to AD	Primary care and specialists in memory disorders
Monitoring	To assess changes in disease severity or the effect of a treatment	Specialists in memory disorders
Pharmacodynamic	To detect changes in response to treatment	Specialists in memory disorders and Clinical trials
Predictive	To predict a favorable or unfavorable effect of a treatment	Specialists in memory disorders and Clinical trials
Safety	To detect an adverse event	Specialists in memory disorders and Clinical trials

Table 3.

Appropriate and inappropriate uses of blood-based markers in clinical practice

	Appropriate use	Inappropriate use
What blood-based markers?	<ol style="list-style-type: none"> 1. p-tau, Aβ42/40 alone or in combination with other biomarkers, in individuals with typical amnesic presentation 2. NFL to explore neurodegenerative process 	<ol style="list-style-type: none"> 1. Any biomarker quantified in an unregulated, non-certified, non-accredited laboratory
When to use blood-based markers?	<ol style="list-style-type: none"> 1. In individuals with objective cognitive impairment (possible or probable AD, MCI/dementia) 2. If suspicion of AD, as part of the initial diagnostic workup 3. If any contraindication or patient aversion to LP (CSF biomarkers) 	<ol style="list-style-type: none"> 1. Instead of the cognitive testing 2. In cognitively unimpaired individuals, except context of clinical research 3. Use to determine disease severity in patients having already received a diagnosis of AD 4. APOE4 carriers with no cognitive impairment
Where to use blood-based markers?	<ol style="list-style-type: none"> 1. In primary care to help PCP referring patients to specialists in memory disorders 2. In primary and specialty care to aid in diagnosis of AD (positive biomarkers along with classical cognitive presentation). 3. In clinical trials (Research Setting) 	<ol style="list-style-type: none"> 1. In any facility in the absence of trained physicians
How to interpretate blood-based markers?	<ol style="list-style-type: none"> 1. Holistic approach, model combining blood-based biomarkers and cognitive performance 2. Need to perform CSF biomarkers or PET if clinical presentation, structural imaging or other evaluative tests conflict with the blood-based biomarker test result 	<ol style="list-style-type: none"> 1. Interpretation of biomarkers without considering the history, clinical exam, cognitive testing, and patient autonomy

Aβ, Amyloid-beta; AD, Alzheimer Disease; CSF, Cerebrospinal Fluid; LP, Lumbar Puncture; PCP, Primary Care Physician; NFL, Neurofilament light Chain; P-tau, phosphorylated tau.